# BMJ Best Practice Getting started guide



## How to access



Visit bestpractice.bmj.com

Access instantly from your institution's network and click 'Sign in'\* \*If you need help setting up BMJ Best Practice, contact your local library.

If you are a new user, you will need to now register for a BMJ Best Practice user profile. Use your user profile to download the app for access anywhere, anytime, even offline.

BMJ Best Practice

Indates III



### Download the BMJ Best Practice app

Search for 'BMJ Best Practice', click download and enter your user profile details to log in.

Our app works offline, so you can access clinical content wherever and whenever you like!

> Need help? If you have any questions, please do not hesitate to contact our support team at: support@bmj.com | +44 (0) 20 7111 1105



# Your expert guide to diagnosis, prognosis, treatment and prevention, anytime, anywhere



Search thousands of clinical topics and over 6,000 clinical guidelines



Watch animated videos of common medical and surgical procedures



500+ patient leaflets which provide concise, user-friendly summaries



Stay current with Important Updates and Cochrane Clinical Answers highlighted within each topic Ensure your ongoing professional development, with automatic CME/CPD tracking





Get step-by-step guidance via our unique treatment algorithm tables



Access anywhere, online and offline with the BMJ Best Practice app



### Comorbidities Manager available within BMJ Best Practice

The Comorbidities Manager supports the management of the whole patient by providing guidance on the treatment of acute conditions alongside their pre-existing comorbidities.

You can select from a range of common, high-cost comorbidities to generate an instant treatment plan tailored to the unique needs of the patient. The Comorbidities Manager makes it easy to quickly treat and stabilise patients with comorbidities. This effective management of the patient increases efficiencies and quality of care, ultimately leading to better clinical outcomes, shorter hospital stays and fewer readmissions.

				Claim 2.62 CM	E/CPD c	redits	) English	~	
BMJ Be	st Practio	ce			Search o	conditions, syr	nptoms		C
/haťs new 🗸	Specialties	Calculators	Patient	leaflets Mu	ultimedi	ia 🗸 🕠	About us 🗸	Your p	rofil
Bacterial r	neningitis in	adults						J. View	v PDI
OVERVIEW	THEORY	DIAGN	IOSIS	MANAGEMENT		FOLLOW UP	R	ESOURCES	
Summary	Epidemiology Aetiology Case history	Histor	mmendations ry and exam tigations entials	Recommendation Treatment algorit Emerging Prevention Patient discussion	hm C	Monitoring Complication Prognosis	s li F F	Buidelines mages and vid teferences l'atient leaflets Calculators Evidence	eos
100000	ey disease (CKD)	not select comorb Heart failure Stroke			algorithm		DTHER CONS Suspected Current sn	IDERATIONS	^
If your patient is     Diabetes     Chronic kidne     Hypertension     Coronary arter	s pregnant or a child, do C ay disease (CKD) C	Heart failure Stroke COPD Asthma	idities using this !	tool. Use the standard a	algorithm		THER CONS	IDERATIONS	^
If your patient in  Diabetes  Chronic kidn  Hypertension  Coronary arts  The following cc  Look out for this  INITIAL  Suspected bi	s pregnant or a child, do ay disease (CKD) C ay disease CKD) C an ary disease C morbidites have been loon: C for treatme	Hoart failure     Heart failure     Stroke     COPD     Astma	idities using this I re affected, or a INITIAL   suspe immunocompe	tool. Use the standard of Depression Dementia Indeed, as a result of rected bacterial menini	your pa	tient's comor	bidities.	IDERATIONS I frailty noker	~ ~
If your patient in     Diabotes     Chronic kidne     Hypertension     Coronary arts     The following cc      Look out for this      INITIAL     suspected bi presenting in	s pregnant or a child, do ay disease (CKD) C ay disease CKD) C an ary disease C morbidites have been loon: C for treatme	Heart select comorb     Heart failure     Stroke     COPD     Astma	idities using this I	tool. Use the standard in Depression Dementia	your pa	tient's comor	bidities.	IDERATIONS I frailty noker	
If your patient in     Diabotes     Chronic kidne     Hypertension     Coronary arts     The following cc     Look out for this     INITIAL     suspected by     presenting in	a pregnant or a child; de ay disease (CKD) E ary disease (CKD) E ary disease E comorbidites have been loon: Cror treatme acterial meningitis: h bospital d immunocompetent	not select comoth } Heart failure Stroke COPD Asthma added: OMAREE added: OMAREE A	differ using this i affected, or a INITIAL   suspe- immunocompe PLUB Corticost Treatment recomm Give empirical in	tool: Use the standard Depression Derrentia Indeed, as a result of Indeed bacterial menin tent tent tertion tert	your par ngitis: p	tient's comor resenting in I spatient group all adults with	bidities.	In frailing noker	<
If your patient in     If your patient in     Diabotes     Chronic kidn.     Hypertansion     Coronary arti     The following co     Look out for this     INITIAL     Suspected by     resenting	a program or a child, dd y y disease (CKD) [ y y disease (CKD) [ y ry disease ] Incon: (or treatment Incon: (or tr	n of select convolve ☐ Heart failure ☐ Stroke ☐ COPD ☐ Addmail added: ©ANIEST and options that a	diffee using this i e affected, or a INITIAL   suspe immunocompe Pus Corticost Treatment recomm Give empirical in hour of presenta	tool: Use the standard Depression Dementia	your par ngitis: p n selected asone to	tient's cornor resenting in I all adults with Cochrane 1 Library	bidities.	In Frailing socker	e ithin

#### Follow these steps to use

- 1 Click on 'Your Profile' and select 'Comorbidities Manager' from the list.
- 2 Select your chosen acute topic from the list. We are adding more topics to this list all the time, so keep checking back for updates.
- 3 Navigate to the 'Treatment algorithm' section of your chosen acute topic. The pink C+ symbol highlights that the Comorbidities Manager is available for this topic.
- Select from a list of common comorbidities. You can select as many comorbidities as you like. You will see the pink C+ symbol when a treatment option may be impacted by the comorbidities that have been selected.