

Patient information from BMJ

Last published: Aug 05, 2021

Atrial fibrillation: what is it?

Atrial fibrillation can be alarming, especially the first time it happens. Your heart beats very fast. It may go away on its own, or you may need treatment to make your heart beat smoothly again.

What happens?

Atrial fibrillation is when your heart doesn't beat regularly. For some people it lasts only a short time - from a few minutes to a day or two. For others it comes and goes and can last a long time.

Your heartbeat is controlled by electrical signals from a 'pacemaker' in your heart. If you have atrial fibrillation the pacemaker sends out too many signals at once. Your heart rate will probably be fast and not have its regular pattern.

Men are more likely than women to get atrial fibrillation. It's also more common if you:

- are older
- have high blood pressure, or
- have heart problems.

Atrial fibrillation can also sometimes be linked to drinking a lot of alcohol, or to medical problems such as:

- diabetes
- lung diseases, and
- problems with your thyroid gland.

Atrial fibrillation can make you more likely to have a stroke. But there are good treatments to help prevent this.

What are the symptoms?

The most common symptom is feeling your heart racing.

Atrial fibrillation: what is it?

But this doesn't happen to everyone. Some people only find out that they have atrial fibrillation after they have a stroke. Signs of a stroke include feeling numb down one side of your body, problems with your vision, and having trouble speaking or swallowing.

If you have atrial fibrillation you might also feel chest pain, get breathless, and feel dizzy. If you get these symptoms you should call an ambulance or go to hospital straight away. They could also be signs of a heart attack.

Tests for atrial fibrillation include:

- An ECG (electrocardiogram), which shows how your heart is beating
- An echocardiogram, which is an ultrasound scan to produce a picture of your heart.

What will happen to me?

More than half the people who have short-term atrial fibrillation get their normal heart rhythm back in less than two days without any treatment. But for some people it's a long-term condition.

Atrial fibrillation increases your risk of having a stroke. But having treatment can help reduce this risk. To learn more, see *Atrial fibrillation: what treatments work?*

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group