

Patient information from BMJ

Last published: Jun 02, 2023

Catheterisation (male): having a catheter fitted

Sometimes it's necessary to have a catheter fitted to help you to pass urine - often because you are having surgery. This leaflet tells you how and why it's done, how it helps, and if there are any possible problems.

You can use our information to talk with your doctor about your treatment.

What is a urethral catheter?

A urethral catheter is a thin rubber tube that's placed into your urethra to help you urinate. Your **urethra** is the tube that carries urine from your bladder down through your penis and out of your body when you go to the toilet.

The catheter goes all the way along the urethra and into the bladder and allows urine to pass easily out of the body.

Why might I need a urethral catheter?

There are several reasons why you might need to have a urethral catheter fitted. They include:

- having surgery in the area of your abdomen (tummy). You always need to have an empty bladder if you are having any kind of surgery around your abdomen. A full bladder can get in the way of your operation, and can easily be damaged if the surgeon is operating on nearby tissue
- if you suddenly find it hard to urinate, even though your bladder is full. This is called **acute urinary retention**. This can have several causes, including damage to nerves or to the bladder itself
- when your doctor needs to measure accurately how much urine your body is making. This is usually because you are ill. Healthy urine production is one way of checking that your body's systems are working properly
- bladder irrigation. This is when the bladder is rinsed out, to clear away material such as blood clots

Catheterisation (male): having a catheter fitted

- bladder instillation. Medications are pumped gently into the bladder through the catheter to help treat the symptoms of cystitis, such as pain and burning
- a bladder obstruction. Several medical conditions can cause a bladder obstruction. This tends to be more common in men.

What will happen?

To have the catheter fitted, you need to be undressed and lying on your back.

Before fitting the catheter your doctor will give you a local anaesthetic so that you don't feel any pain. This is done by pumping anaesthetic gel into the urethra, at the top of your penis. It's done with a syringe, but there's no needle and no puncture of the skin.

The doctor will wait for a minute or so, for the anaesthetic to start working. He or she will then:

- insert the tip of the catheter into the end of your penis
- guide the catheter into the urethra with steady, gentle pressure
- wait until urine appears at the end of the catheter. When this happens, the doctor knows that the catheter is in the bladder. He or she will guide it a little farther in, so that it's as secure as possible
- pump a small amount of sterile water into the catheter. This inflates a small balloon at the end of the catheter that's in your bladder, which stops it from falling out
- attach a bag to the end of the catheter to collect the urine that you pass.

Some people need to have catheters fitted to help with long-term medical conditions. But in most cases the catheter will be removed as soon as you no longer need it, to reduce the chance of an infection.

If you need to have your catheter in place for more than a short time, and you are able to walk, the collection bag can be simply attached to your leg, so that you can walk around.

If you need to have a long-term catheter, it will need to be changed every few weeks.

What are the risks?

Having a catheter fitted shouldn't hurt, because of the local anaesthetic. But it can feel a little uncomfortable. There are several other possible complications that your doctor should discuss with you before the procedure, including:

- urinary tract infection (UTI). All the equipment your doctor uses will be sterile. But infections do still happen sometimes. If you get a UTI you might need antibiotics to treat it
- bleeding. A small amount of bleeding is common after having a catheter fitted. It usually stops by itself. But sometimes the bleeding can be heavier and take longer to stop
- a blocked catheter. Material from inside the bladder or urethra, such as a blood clot, can sometimes block the catheter. If this happens, you will need to have a new catheter fitted.

Catheterisation (male): having a catheter fitted

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group