

Patient information from BMJ

Last published: Apr 17, 2023

BCG vaccination

BCG is a vaccination that protects against infection with tuberculosis (TB). BCG stands for Bacillus Calmette-Guérin.

The BCG vaccine used to be part of the standard vaccination programmes in most countries. But it is no longer routinely used in many countries, including the US.

The vaccine is still widely used in countries with high rates of TB, including those in Africa and some Asian countries.

What is tuberculosis?

Tuberculosis is a bacterial infection that mainly damages the lungs. It is one of the top ten causes of death worldwide.

The disease is spread from person to person in droplets of fluid when someone coughs or sneezes.

TB doesn't spread as easily as some diseases, such as colds and flu. To become infected, you usually have to spend a long time in close company with someone who has TB.

The infection spreads most easily in places with cramped, overcrowded housing. Most deaths from TB happen in Africa, the Indian subcontinent, China, and other parts of Asia, including South-East Asia.

TB is especially dangerous in areas with high rates of **HIV**, as people with weakened immune systems find it harder to fight the infection.

TB can be treated with **antibiotics**. But the disease can be hard to cure, and some people need to take antibiotics for many months. Even then, some strains of TB are **drug resistant**. This means that some antibiotics don't work on these strains.

Symptoms of TB include:

- severe coughing, which can include coughing up blood
- chest pain
- fever and night sweats

BCG vaccination

- loss of appetite, and
- weakness and fatigue.

If it is not treated, TB can affect other parts of the body, including the bones and kidneys.

TB is not common in developed countries. But it can be present in communities:

- where people have family who came from countries where TB is more common, and
- where people have recently arrived from countries where there are high levels of the disease.

Who needs the BCG vaccination?

Children at high risk

The BCG vaccine is usually given to children who are at risk of being infected with TB.

In the US, children are usually only offered the BCG vaccine if they live with adults who:

- have TB and who might not have been properly treated for the infection, or
- have certain strains of drug resistant TB.

Young people and adults

The BCG vaccine is not usually given to people aged over 16 years, because it doesn't work very well in adults.

But some adults might be offered this vaccine if they are thought to have an increased chance of getting TB.

For example, healthcare workers might be offered the vaccine if they are likely to come into contact with people who have drug resistant TB.

If you wish to have a BCG vaccination, or if you wish your child to have it, you will need to speak to your doctor, who will assess whether you need the vaccine.

Is there anyone who should not have the vaccine?

People who have already had TB, or who are currently infected with the disease, should **not** have the vaccine. This is because:

- it won't help, and
- it can cause unpleasant side effects.

This means that, before being given the vaccine, you will need to have a test called the **tuberculin skin test** or **Mantoux test**.

This involves injecting a small amount of a substance similar to TB just under the skin. If your skin reacts to the injection, it's likely that you have been exposed to TB before, and you should not have the vaccine.

BCG vaccination

If the test is negative you can have the vaccine.

Other people who should not have the BCG vaccine include:

- pregnant women, and
- people with a weakened immune system (this is called being immunocompromised), either because of a medical condition, or because they are taking medication that can weaken (suppress) the immune system.

How is the vaccination given?

The BCG vaccine is given in the same way as many other vaccines, as an injection in the upper arm.

Is it safe?

The most common side effect of the BCG injection is a small scar at the injection site in the upper arm.

Over several weeks after the injection you might notice a blister appear. This might then ooze and form a crusty scab. You should leave it uncovered and allow it to heal. If you or your child has a more severe skin reaction, tell your doctor.

Other common temporary side effects of the BCG vaccine can include:

- a headache
- fever, and
- swollen glands.

More serious side effects include abscesses and swelling in some bones. But these are rare.

Allergic reactions to the vaccine

Some people have an allergic reaction to the BCG vaccine. But the medical staff who give you the vaccine will be trained in treating allergic reactions.

Even so, you should tell the medical staff before you have the vaccination if you or your child has any allergies.

How well does the vaccine work?

The vaccine is about 70% to 80% effective against the most severe types of TB, including TB meningitis that affects children. This means that it works for between 70 and 80 in every 100 children.

It is less effective against TB that affects the lungs, which is the most common type of TB in adults. And it is less effective in adults in general, against all types of TB.

BCG vaccination

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group