

Patient information from BMJ

Last published: Mar 05, 2021

Depression in adults: questions to ask your doctor

If you've been diagnosed with depression, you may want to talk to your doctor or psychiatrist to find out more.

Here are some questions you may want to ask. You may find it helpful to have someone with you, to help you find out everything you want to know.

- I feel very down. Is this just a normal feeling, or do I have depression?
- Do I need treatment?
- What treatment do you recommend?
- Does the treatment have any side effects?
- When will I start to feel better?
- How long will I need to have this treatment?
- What will happen to me if I don't have this treatment?
- What kind of talking treatments (psychotherapies) are available? Would any of them help me?
- Are there any alternative therapies or complementary therapies that may help?
- Is there anything I can do to avoid getting depressed again?
- Are there any support groups for people with depression in my area?
- What should I do if I have thoughts about harming myself?

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2024. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group