

Patient information from BMJ

Last published: Sep 14, 2022

Ischemic bowel disease: what are the treatment options?

Ischemic bowel disease can happen when the blood supply to your intestines (bowels) becomes restricted. It can come on suddenly or develop over time. Treatment can help many people, but it is a serious condition that can be life threatening.

What treatments are available?

Like all the body's organs, the digestive system needs a healthy supply of blood to work properly. Blood brings oxygen and nutrients and carries away waste products. If the blood flow to the intestines is restricted for any reason it can cause ischemic bowel disease. The word "ischemia" means restricted blood supply.

Ischemic bowel disease is not one specific condition. The name covers any problems that can be caused by reduced blood supply to the bowel. For more background information see our leaflet *Ischemic bowel disease: what is it?*

The treatment you need will depend on which blood vessels are affected and on how severe the problem is. But many people need surgery.

Mild ischemic bowel disease

If the blood supply to your bowel is only slightly restricted, and if there are no signs of infection, you probably won't need surgery. Some types of mild ischemic bowel disease get better by themselves.

But even if your ischemia is mild you will need treatment to make you comfortable and stable and to prevent the problem getting worse.

First treatments

The first treatments you will be given will be to help make you stable. They may include oxygen via a mask and intravenous (IV) fluids. You may need a tube inserted into your stomach or treatment to correct an irregular heart rhythm.

You will also be given antibiotics. This is because any damage to the bowels can cause bacteria to spread quickly around the body. If you have signs of heart failure you will also be given medication to help your heart pump more strongly.

Ischemic bowel disease: what are the treatment options?

Surgery

Many people with ischemic bowel disease need surgery. This is usually done either to remove a blockage from a blood vessel or to bypass a blocked or badly damaged portion of a blood vessel.

Both these methods can help restore a healthy flow of blood. The type of surgery you need depends on which blood vessels are affected and how badly.

You might also need surgery if the lining of your abdomen is inflamed (this swelling is called peritonitis), or if there is a tear in your bowel. To begin with you will need what's called exploratory surgery.

This means that the surgeon makes a cut in the abdomen and is then able to see clearly any swelling and damage. He or she can then remove any badly damaged section of bowel and repair the more healthy parts.

After surgery

If you have surgery you will need drug treatment to help your recovery. You will probably need to take at least one of the following drug treatments. You may need to take some of them for some weeks or months:

- Antibiotics to target any specific infection that has been identified
- Drugs called anticoagulants to stop your blood from clotting too easily and to help prevent more blockages while you recover
- Other drugs to help dilate (open up) narrow blood vessels to improve blood flow. You only take these for a few days.

If you have inflamed blood vessels you may need to take powerful anti-inflammatories called corticosteroids to help relieve the swelling.

What to expect in the future

The follow-up care that you need will depend on whether you needed surgery.

If you didn't have surgery you may need to have tests and procedures so that your doctor can check for any damage to your digestive system.

If you had surgery you will need regular check-ups to make sure you are healing well. But, whether you had surgery or not, you will need check-ups so that your doctor can look out for signs of anything that might cause another episode of ischemia. You will also need treatment for any other problems that might have caused the ischemia.

You will need to be aware of the symptoms of bowel ischemia returning. If you have any abdominal pain, bloating, bloody stools, nausea, or vomiting, seek medical help right away. And it's vital that you take your follow-up medications as instructed by your doctor.

Unfortunately an episode of bowel ischemia can be fatal, especially if it was diagnosed late and there is already a lot of damage. The chances of survival also depend on which blood vessels are involved and how bad the blockage is.

Ischemic bowel disease: what are the treatment options?

The outlook is better for people with chronic ischemic bowel disease. With treatment, between 70 and 90 in 100 people survive. For people whose large intestine is affected, rather than their small intestine, the outlook is even better.

But acute ischemic bowel disease tends to be more serious. More than half of people with this type of ischemia will die.

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2025. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.



BMJ Group