
BMJ Best Practice

Assessment of abdominal pain in children

Straight to the point of care



Last updated: Dec 12, 2024

Table of Contents

Overview	3
Summary	3
Theory	4
Aetiology	4
Emergencies	30
Urgent considerations	30
Diagnosis	34
Approach	34
Differentials overview	50
Differentials	52
Guidelines	101
Evidence tables	103
References	105
Images	122
Disclaimer	143

Summary

Paediatric abdominal pain is a common problem that often poses a diagnostic dilemma. The vast majority of episodes are benign and self-limiting, but persistent abdominal pain may signify an underlying pathology requiring urgent intervention. Acute abdominal pain can also be caused by a wide range of conditions, some of which are emergencies (e.g., acute appendicitis). Timely assessment and intervention are critical in children presenting with abdominal pain.

A broad spectrum of aetiologies can cause abdominal pain, and diagnosis can be challenging. In most cases, a thorough history and physical examination can narrow the differential diagnoses. Additional investigations may be required to confirm the diagnosis.

Furthermore, even with the assistance of parents or guardians, a comprehensive history is often difficult to obtain, and diagnosis therefore relies heavily on the clinical acumen of the practitioner.

Aetiology

The differential diagnoses for paediatric abdominal pain are broad and encompass almost every organ system. In addition, distinguishing acute from chronic abdominal pain may be particularly difficult in children. Although the most common aetiologies are not immediately life-threatening, the ability to diagnose urgent pathology remains paramount. A thorough history and physical examination, and an understanding of the more common diseases affecting the child's age group, are essential.

Gastrointestinal

Gastrointestinal (GI) sources are the most common aetiology of abdominal pain in children, encompassing infectious, congenital, functional, and mechanical causes.

Constipation

- A common condition, with a reported pooled prevalence of 9.5%.[1]
- Childhood constipation is typically characterised by infrequent bowel evacuations, large stools, and difficult or painful defecation.[2] [3]
- Symptoms usually result from low-fibre, poor-nutrient intake, and too little water, which leads to high
 levels of colonic reabsorption of water and hardening of the stool. Additional risk factors include
 genetic predisposition, infection, stress, obesity, low birth weight, cerebral palsy, spina bifida, and
 learning difficulties.
- Constipation starts as an acute problem but can progress to faecal impaction and chronic constipation.
- It tends to develop during three stages of childhood: weaning (infants), toilet training (toddlers), starting school (older children).


Infantile colic

- Characterised by paroxysms of uncontrollable crying in an otherwise healthy and well-fed infant aged
 5 months. The duration of crying is >3 hours per day, and >3 days per week, for at least 3 weeks.[4]
- The crying typically starts in the first weeks of life and ends by 4-5 months of age.
- · Food allergy may play a role in the pathogenesis.
- Colic occurs equally in both male and female infants.[5] Infants with colic tend to have siblings who
 also have this condition.


Appendicitis

- Develops when the appendiceal lumen becomes obstructed by stool, barium, food, or parasites.
- Can occur in all age groups, but is rare in infants. A cohort study in Sweden found that 2.5% of children had had appendicitis by age 18 years.[6]
- If left untreated, acute appendicitis may progress to ischaemia, necrosis, and eventually perforation.

 The overall rate of perforation is about 30%.[7]
- Validated paediatric clinical prediction rules may help to quantify risk for appendicitis, but guidelines recommend against making a diagnosis of appendicitis in children based on clinical scores alone.[8]
 [9] [10] [11] [12] [13]
- Although the risk of perforation increases if appendectomy is substantially delayed, surgery within 24 hours does not increase the risk of perforation for children with uncomplicated acute appendicitis.[13]
 [14] [15]



Necrotic appendix
From the collection of Dr KuoJen Tsao; used with permission



CT scan demonstrating faecalith (white arrow) outside the lumen of the appendix consistent with perforated appendix


• Stump appendicitis may occur after an appendectomy when the residual stump is >0.5 cm. It can occur after both open and laparoscopic surgery.[16]

Gastroenteritis

- May be due to acute or chronic viral infection (especially rotavirus), or bacterial or parasitic infection.
- Causes vague, cramping abdominal pain in association with fever, vomiting, and diarrhoea.
- Eosinophilic gastroenteritis, defined as a condition affecting the GI tract with eosinophil-rich inflammation without a known cause for the eosinophilia, can result in significant abdominal pain.[17]
- Haemolytic uraemic syndrome, characterised by microangiopathic haemolytic anaemia, thrombocytopenia, and nephropathy, can occur as a complication of gastroenteritis caused by verotoxin-producing *Escherichia coli*. Abdominal pain is a common presenting symptom.[18]

Intussusception

• Occurs when a proximal segment of the intestine telescopes into the lumen of an immediately distal segment. In most cases, the intussusception is in the ileocaecal area.




Intussusception: blood vessels become trapped between layers of intestine, leading to reduced blood supply, oedema, strangulation of bowel, and gangrene. Sepsis, shock, and death may eventually occur Created by the BMJ Knowledge Centre

- Usually occurs in infants between 3 and 12 months of age. Peak incidence is 5-7 months of age.[19]
- Intussusception should be suspected in an infant in this age group presenting with colicky abdominal pain, flexing of the legs, fever, lethargy, and vomiting.

- In infants <2 years of age, episodes of intussusception are most likely to be caused by mesenteric lymphadenopathy secondary to an associated illness (e.g., viral gastroenteritis). In older children, mesenteric lymphadenopathy is still the most likely cause, but other aetiologies should be considered (e.g., intestinal lymphomas, Meckel's diverticulum). Therefore, children ≥6 years or with jejuno-jejunal or ileo-ileal intussusception should be assessed for a pathological lead point.
- Ileo-ileal intussusception may also be indicative of Henoch-Schonlein purpura (HSP).[20] [21] HSP is a vasculitis that affects small veins and primarily occurs in children <11 years of age.[22]
- Occasionally, an ultrasound examination will show small-bowel to small-bowel intussusception. If there
 are no obstructive symptoms, and the wall of the bowel is normal, with length <3.5 cm, and normal
 vascularity with no colonic involvement, then these findings are usually transient and require no further
 intervention.[23]

Meckel's diverticulum

• A finger-like projection located in the distal ileum arising from the anti-mesenteric border; usually 40-60 cm from the ileocaecal valve, measuring 1-10 cm long and 2 cm wide.



Intraoperative photo of Meckel's diverticulum

From the collection of Dr Kuojen Tsao; used with permission

- The majority of symptomatic patients present before the age of 2 years.
- Meckel's diverticulum is a common cause of paediatric gastrointestinal bleeding and should be considered if there is painless rectal bleeding.
- The prevalence is estimated to be up to 3%.[24]
- Intestinal obstruction is a known complication and may be observed in as many as 40% of all symptomatic Meckel's diverticula (according to some series).[25] [26]

Mesenteric adenitis

- Refers to inflammation of the mesenteric lymph nodes. This process may be acute or chronic.
- It is often mistaken for other diagnoses, such as appendicitis; up to 23% of patients undergoing negative appendectomy have been found to have non-specific mesenteric adenitis.[27]
- One retrospective study reported that, compared with children who have appendicitis, patients who have mesenteric adenitis are more likely to have high fever (above 39°C) and dysuria, and are less likely to have migratory pain, vomiting, or typical abdominal signs of appendicitis on examination.[28]

Hirschsprung's disease

- Most commonly diagnosed in the first year of life but can present later in childhood; higher male preponderance in the most common anatomical variant (rectosigmoid disease).[29]
- Congenital condition characterised by partial or complete colonic obstruction associated with the
 absence of intramural ganglion cells. Because of the aganglionosis, the lumen is tonically contracted,
 causing a functional obstruction. The aganglionic portion of the colon is always located distally, but the
 length of the segment varies.



Abdominal x-ray of a neonate with abnormal stooling pattern and constipation. The dilated transverse and descending colon is suggestive of Hirschsprung's disease

- Hirschprung-associated enterocolitis is the main cause of serious morbidity and death in patients with Hirschprung's disease if not recognised and treated promptly.[30] The classic presenting features include abdominal distention, fever, and diarrhoea; however, other signs or symptoms may include vomiting, rectal bleeding, lethargy, loose stools, and obstipation.[30] [31]
- May be associated with Down syndrome and multiple endocrine neoplasia type IIA.

Intestinal obstruction

- Small or large bowel obstruction may be the result of various aetiologies and can occur at any age.
 Abdominal pain may not occur until the obstruction has progressed to include extensive abdominal distension or intestinal ischaemia. Intestinal obstruction may mimic intestinal ileus, which usually does not require surgical intervention.
- Intestinal obstruction in a child without a history of prior surgery is traditionally considered an indication for surgery.

- The aetiology of intestinal obstruction can be congenital or acquired. Congenital causes include atresias or stenosis, which present in the newborn period. Acquired causes include small bowel adhesions, strangulated or incarcerated hernias, and tumours.
- · Congenital causes:
 - Duodenal atresia or stenosis may cause complete or partial obstruction of the duodenum
 as a result of failed re-canalisation during development. This results in either stenosis with
 incomplete obstruction of the duodenal lumen (allowing some but not all gas and liquid to pass)
 or an atresia where the duodenum ends blindly causing a true complete obstruction.
 - Jejuno-ileal atresia or stenosis is a complete or partial obstruction of any part of the jejunum or ileum. Although uncertain, it is believed to result from a vascular accident during development. Jejunal stenosis may still have bowel lumen continuity with a narrowed lumen and thickened muscular layer. There are four types of atretic bowel: type I, an obstructing web or septum with intact bowel wall and mesentery; type II, an atretic cord of remnant bowel with an intact mesentery; type IIIa, a missing atretic segment with a mesenteric defect; type IIIb, atresia with the distal bowel coiled ('apple peel') around a distal mesenteric vessel; and type IV, multiple segmental atresias ('sausage-links').
 - Hernias may be internal or external and congenital (most commonly in children) or acquired.
 - Colonic atresia is a rare complete obstruction of any part of the colon, although it usually occurs near the splenic flexure. Like jejuno-ileal atresia, it is thought to occur as a result of a vascular event.




Abdominal x-ray demonstrating double bubble gas pattern consistent with duodenal atresia

From the collection of Dr KuoJen Tsao; used with permission

- Meconium ileus is an important cause of intestinal obstruction in the neonatal period; cystic fibrosis should be suspected as an associated disease. There may be associated pancreatic abnormalities.
- Duplication cysts occur most commonly in the small intestine; they may serve as a lead point
 for volvulus and intussusception and can also result in obstruction. In the presence of duodenal
 duplication cysts, peptic ulcer disease, haemorrhage, or perforation may result secondary to
 ectopic gastric mucosa.

Acquired causes:

- May occur at any age.
- Tumours may be intraluminal or extra-intestinal.
- Hernias may be internal or external and congenital or acquired (e.g., prior operation, incisional, traumatic).



Infant with right groin bulge consistent with incarcerated inguinal hernia. The lack of overlying skin oedema and erythema does not rule out strangulation of the small intestine From the collection of Dr KuoJen Tsao; used with permission

- A history of previous intra-abdominal surgery or inflammation (such as necrotising enterocolitis) should prompt concern for adhesive small bowel obstruction.
- Bowel obstruction without previous surgical or inflammatory history in the paediatric population should raise concern for aetiologies such as a mass or cyst that requires intervention.
- Omental cysts, although rare, can present with intestinal obstruction; may be confused with ovarian cysts on ultrasound.
- In patients with cystic fibrosis, partial bowel obstruction may sometimes be referred to as
 distal intestinal obstruction syndrome (DIOS, formerly known as meconium ileus-equivalent
 syndrome). DIOS is not related to meconium. It refers to a distal small bowel obstruction caused
 by impacted bowel contents that typically occurs in adolescents and adults with cystic fibrosis.

Volvulus

- This can occur in any age group but is most common in children <1 year old; at least 60% of children present before 1 month of age.[32] Mid-gut volvulus is the most common type.
- Green (bilious) vomiting is a cardinal symptom of duodenal obstruction secondary to mid-gut volvulus.[32]
- Intestinal malrotation is a term used to encompass the entire spectrum of anatomical arrangements that result from incomplete rotation of the gut during embryonic development. Volvulus of the entire small bowel and part of the colon is only possible, unless in extremely preterm infants, when malrotation exists.[33]
- In malrotation, the most significant pathological concerns are a lack of gut fixation to the retroperitoneum and narrow mid-gut mesenteric base that predisposes patients to mid-gut volvulus, which occurs when the duodenum or colon twist around this mesenteric base.
- Colonic volvulus (sigmoid or caecal) is rare.[34] It usually occurs in children with a history of chronic constipation, in dysmotility disorders, or in patients who have cognitive deficits with limited mobility.

Necrotising enterocolitis

- A disease primarily of premature infants, particularly those weighing less than 1500 g.[35] The
 pathogenesis is multifactorial and not well understood, although ischaemia, reperfusion injury,
 microbiome changes, and infectious pathogens may play a role.
- Typical symptoms are feeding intolerance, abdominal distension, and bloody diarrhoea at approximately 1-2 weeks of age.[36] Other signs and symptoms include apnoea, lethargy, abdominal tenderness, abdominal wall erythema, and bradycardia.

Peptic ulcer disease

- Gastric and duodenal ulcers are uncommon among the paediatric population.[37] When they occur, they are classified as primary or secondary peptic ulcers.
- Primary ulcers occur without predisposing factors and are most commonly located in the duodenum
 or pyloric channel. They manifest most often in older children and adolescents with a positive family
 history. Rarely, primary peptic ulcers can occur in the first month of life, presenting with bleeding
 and possible perforation. Most are located in the stomach. Primary ulcers may be associated with
 Helicobacter pylori.
- Secondary ulcers are usually associated with stress, burns, trauma, infection, neonatal hypoxia, chronic illness, and ulcerogenic medications or lifestyle habits (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], salicylates, corticosteroids, smoking, or intake of caffeine, nicotine, or alcohol).[37]
 It is important to treat the predisposing condition. Exacerbations and remissions can last for weeks to months.

Inflammatory bowel disease (IBD)

- This category includes ulcerative colitis, Crohn's disease, and indeterminate colitis (up to 30% of paediatric cases of IBD in retrospective cohorts, but initial diagnosis may be reclassified during followup).[38] [39]
- Ulcerative colitis affects the rectum and extends proximally; it is characterised by diffuse inflammation
 of the colonic mucosa and a relapsing, remitting course. Ulcerative colitis is uncommon in children, but
 prevalence is increasing.[40]
- Crohn's disease may involve any or all parts of the entire GI tract from mouth to perianal area. Unlike ulcerative colitis, Crohn's disease is characterised by skip lesions. The transmural inflammation often leads to fibrosis, causing intestinal obstruction. The inflammation can also result in sinus tracts that

- burrow through and penetrate the serosa, giving rise to perforations and fistulas. Onset of Crohn's disease typically occurs in the second to fourth decade of life.[41] [42] [43]
- Ulcerative colitis often presents with bloody diarrhoea, whereas this is an unusual presentation in Crohn's disease. Both conditions cause cramping abdominal pain, anorexia, and weight loss when they present late in the course of the disease. Depending on the intestinal location of Crohn's disease, it may mimic other disease processes such as acute appendicitis.
- IBD in the paediatric population can start with very subtle signs that are difficult to interpret. IBD should be considered for any vague, ongoing chronic abdominal pain combined with a slowing of the patient's normal growth curve.

Coeliac disease

- Systemic autoimmune disease triggered by dietary gluten peptides found in wheat, rye, barley, and related grains.
- Immune activation in the small intestine leads to villous atrophy, hypertrophy of the intestinal crypts, and increased numbers of lymphocytes in the epithelium and lamina propria. Locally these changes lead to GI symptoms and malabsorption.
- Coeliac disease is a common disorder in the US and in Europe. A relatively uniform prevalence has been found in many countries, with pooled global seroprevalence and biopsy-confirmed prevalence of 1.4% and 0.7%, respectively.[44]
- Patients may present with recurrent abdominal pain, cramping, or distension.[45] Other common symptoms include bloating, weight loss, vomiting and diarrhoea.[46] Dermatitis herpetiformis, an intensely pruritic papulovesicular rash that affects the extensor limb surfaces, almost universally occurs in association with coeliac disease.[46]

Cholelithiasis/cholecystitis

 Cholelithiasis describes the entity of stones in the gallbladder (usually asymptomatic or an incidental finding). Biliary colic refers to the classic description of intermittent, recurrent right upper quadrant (RUQ) pain that resolves without intervention. This is usually caused by intermittent obstruction of the cystic duct due to cholelithiasis and contraction of a distended gallbladder.



Gallbladder ultrasound demonstrating cholelithiasis with characteristic shadowing



Abdominal x-ray with opacities in the RUQ consistent with gallstones

Cholecystitis refers to inflammation of the gallbladder precipitated by obstruction of bile through
the cystic duct. Symptoms do not usually resolve spontaneously, and there are specific findings on
diagnostic imaging. Cholecystitis may be acalculous (without stones) or calculous (with stones).
 Choledocholithiasis is the term describing a gallstone(s) in the common bile duct.

Biliary dyskinesia


- Characterised by symptoms of biliary colic (intermittent, recurrent RUQ pain that resolves without intervention) in the absence of documented stones in the gallbladder; the diagnosis should be considered in those with symptoms suggestive of biliary colic but with negative laboratory tests and ultrasound in their work-up for symptomatic cholelithiasis.
- Caused by abnormal or altered contraction of the gallbladder resulting in biliary colic. Patients have
 frequently gone through a comprehensive work-up prior to being diagnosed with this entity; increasing
 recognition and testing for the disease has led to more frequent diagnosis in children.[47]

Viral hepatitis

- The viral hepatitides include A, B, C, D, and E.
- Hepatitis A virus remains a significant cause of acute viral hepatitis and jaundice, particularly in developing countries, in travellers to those countries, and in sporadic food-borne outbreaks in developed countries.
- Hepatitis B virus (HBV) frequently causes acute hepatitis and is the most common cause of chronic hepatitis in Africa and the Far East.
- Hepatitis C virus (HCV) represents the leading cause of chronic viral hepatitis in developed countries.
- Hepatitis D virus is a defective virus that needs the presence of hepatitis B to cause clinically recognisable disease.
- Hepatitis E virus represents a major cause of mortality in developing countries, especially among pregnant females.

Acute pancreatitis

- Refers to inflammation of the pancreas; it does not necessarily imply that infection is present.
- Paediatric acute pancreatitis is classified into mild, moderately severe and severe.[48]
- Pancreatitis in children is often due to drugs, infection, anatomical abnormalities, or trauma.[49]
 Corticosteroids, adrenocorticotrophic hormones, oestrogens including contraceptives, azathioprine, asparaginase, tetracycline, chlorothiazides, and valproic acid may induce pancreatitis. Congenital causes include choledochal cyst causing abnormal pancreas and bile drainage and pancreas divisum. Infectious causes include mumps and infectious mononucleosis.
- Excessive alcohol and gallstones are the most common causes of pancreatitis in adults; these causes are relatively less common in children, although they may still occur. Paediatric pancreatitis is rare, but the growing population of children with gallstones is likely to increase future incidence.




CT scan of teenage girl presenting with mid-epigastric abdominal pain as a result of gallstone pancreatitis. The large fluid collection in the pancreatic bed (white arrow) and lack of pancreatic enhancement suggest liquefactive necrosis of the pancreas

Splenic infarction and cysts

Cysts are classified as either primary or secondary (acquired). Primary cysts are usually congenital
and have a true epithelial lining. Eighty percent of splenic cysts are pseudocysts related to infection,
infarction, or trauma.[50] Most cysts are incidental diagnoses, although some patients may present
with dull, left-sided abdominal pain. In paediatric patients, the most common splenic masses are
congenital and/or acquired cysts.[51]



CT scan demonstrating fluid-filled cyst within the spleen



Intraoperative photo of large splenic cyst

Splenic infarction occurs when there is occlusion of the splenic blood supply. It may affect the whole
organ or only a portion of the spleen, depending on the blood vessels involved. The incidence of
splenic infarction is difficult to assess.

Abdominal trauma

- A multi-centre prospective study found that abdominal trauma accounted for 3% of admissions to paediatric trauma units.[52]
- · Generally classified as penetrating or blunt.
- Occult blunt abdominal trauma should always be considered in the setting of vague or inconsistent history. The liver, spleen, and kidneys are the most commonly injured intra-abdominal organs in blunt trauma. Most cases of blunt injury to the liver and spleen are managed non-operatively.
- It is important to exclude duodenal and/or pancreatic injuries with bicycle handlebar injuries and/or direct blows to the abdomen. Hollow viscus injuries (e.g., stomach and intestines) are more common with penetrating trauma.
- It is essential to consider child abuse/non-accidental trauma in this patient population.

Genitourinary

Urinary tract infection (UTI)

- Infection may arise along any part of the urinary tract including the urethra, bladder, ureter, and kidney.
 Diagnosis and treatment is paramount to prevent potential long-term adverse effects, including renal or urinary tract scarring and hypertension.
- Estimates of the true incidence of UTI depend on rates of diagnosis and investigation. UTI is more common in girls. UTIs affect approximately 4% and 10% of children by ages 1 year and 6 years, respectively.[53]
- Bacterial infections are the most common cause, particularly Escherichia coli infection.

Primary dysmenorrhoea

- Dysmenorrhoea, or painful menstruation, is one of the most common gynaecological conditions affecting females of reproductive age.[54]
- Primary dysmenorrhoea is characterised by menstrual pain in the absence of pelvic pathology.

Nephrolithiasis

- Refers to stones that may be located anywhere in the genitourinary tract; the majority of stones are noted in the kidneys, followed by the bladder and ureter.
- Most patients have a predisposing factor, such as a family history of nephrolithiasis, high-risk diet (e.g., high oxalate intake), or chronic disease (e.g., renal tubular acidosis).
- Stones less than 5 mm in diameter will generally pass spontaneously.

Testicular torsion

• A urological emergency caused by the twisting of the testicle on the spermatic cord, leading to constriction of the vascular supply and time-sensitive ischaemia and/or necrosis of testicular tissue.



Young boy with right testicular pain. The testicle is swollen, tender, and erythematous as a result of torsion of the appendix testes. The clinical signs and symptoms mimic those of testicular torsion



Infant boy with swollen, tender, and erythematous left testicle.

The testicle is retracted consistent with testicular torsion

From the collection of Dr KuoJen Tsao; used with permission

24



Torsion of an appendix testis resulting in acute infarction

- Has a bimodal distribution, with extravaginal testicular torsion affecting neonates in the perinatal period, and intravaginal testicular torsion affecting males of any age but most commonly adolescent boys.[55]
- Systemic symptoms such as nausea and vomiting are not usually present. The thin skin of the scrotum sometimes allows visualisation of the torsed appendage ('blue dot or black dot sign').
- The differential diagnosis includes pain from torsion of a testicular appendage; this may develop more gradually (over days to weeks) and frequently is pinpoint (superior pole of testes).
- Epididymitis can also mimic testicular torsion, but it is more gradual in onset with less severe symptoms.


Ruptured ovarian cyst

- Ovarian cyst rupture is rare and may occur in conjunction with torsion.
- Symptoms usually occur prior to the expected time of ovulation and may mimic ruptured ectopic pregnancy. Pain arises from local peritonitis secondary to haemorrhage.[56] [57] [58]


Ovarian torsion

- Although it can affect females of any age, it most commonly occurs in the early reproductive years. [59]
- In children, torsion of the ovary is often associated with the presence of an ovarian tumour, most commonly a teratoma.

- Twisting or torsion of the ovary compromises the arterial inflow and venous outflow, producing ischaemia, which, if not relieved promptly, can affect the viability of the ovary.
- Oophorectomy is infrequently indicated and most ovaries can be preserved.[60]



Intraoperative photo of ovarian mass that presented as ovarian torsion



CT scan of a young girl presenting with ovarian torsion. The large pelvic cystic lesion contains calcifications (white arrow) consistent with a teratoma or dermoid cyst

Pelvic inflammatory disease (PID)

- Represents a spectrum of upper genital tract infections that includes any combination of endometritis, salpingitis, pyosalpinx, tubo-ovarian abscess, and pelvic peritonitis; usually caused by *Neisseria* gonorrhoeae or *Chlamydia trachomatis* and less commonly by normal vaginal flora including streptococci, anaerobes, and enteric gram-negative rods.
- Adolescents are at higher risk of developing PID compared with older women.[61] Sexually transmitted infections (chlamydia and gonorrhoea) are a key risk factor.[62]
- PID is rare in the absence of sexual activity; PID in a young child should prompt work-up for possible sexual abuse.

Pregnancy complications

- Miscarriage and ectopic pregnancy should be a concern in any female of reproductive age presenting with lower abdominal pain, amenorrhoea, and vaginal bleeding.
- Miscarriage is an involuntary, spontaneous loss of a pregnancy before 20-24 completed weeks.
 The gestational threshold for the definition varies between countries: in the US it is usually 20 weeks (but may vary in different states), whereas in the UK, the Royal College of Obstetricians and Gynaecologists defines it as 24 weeks.[63] [64] The majority of spontaneous miscarriages occur in the first trimester.[65]
- Ectopic pregnancy occurs when a fertilised ovum implants and matures outside the uterine endometrial cavity, with the most common sites being the fallopian tube (97%), the ovary (3.2%), and the abdomen (1.3%).[66] Use of oral contraceptives before age 16 years is associated with increased risk of ectopic pregnancy.[67] The classic presentation includes lower abdominal pain, amenorrhoea, and vaginal bleeding. Haemorrhage from a ruptured ectopic pregnancy can be fatal.

Pulmonary

Primary respiratory illnesses such as pneumonia or empyema may present as abdominal pain in the paediatric population.[68] lleus following a thoracic procedure can lead to abdominal distension, GI symptoms, and discomfort.

Recurrent pneumonia in children is usually the result of a particular susceptibility, such as disorders of immunity and leukocyte function, disorders of ciliary function, anatomical abnormalities, or specific genetic disorders such as cystic fibrosis.[69]

Functional abdominal pain

Functional abdominal pain is also referred to as non-specific abdominal pain; pain is usually chronic or recurrent. Visceral hyperalgesia is the final outcome of sensitising medical and psychosocial events, on a background of genetic predisposition.[70]

Functional abdominal pain disorders are classified according to Rome IV criteria, which describe functional dyspepsia, irritable bowel syndrome, abdominal migraine, and functional abdominal pain - not otherwise specified. [70] [71] [72]

- Typically affects children between 5 and 14 years of age.
- Prevalence estimates vary from 10% to 30% in samples of school students, to 87% in some gastroenterology clinics.[73]
- Family history of functional disorder common (irritable bowel syndrome, mental illness, migraine, anxiety).
- Clarifying the type of functional disorder is important to determine which treatments are most likely to improve symptoms.

Functional dyspepsia

 Defined as one or more of the following bothersome symptoms on at least 4 days per month: postprandial fullness, early satiation, epigastric pain, or burning not associated with defecation. After appropriate evaluation the symptoms cannot be fully explained by another medical condition.[70]

Irritable bowel syndrome

Three criteria must be fulfilled for 2 months prior to diagnosis:[70]

- 1. Abdominal pain at least 4 days per month associated with one or more of:
 - · Related to defecation
 - Change in stool frequency
 - · Change in stool form.
- 2. In children with constipation, the pain does not resolve with resolution of constipation.
- 3. After appropriate evaluation the symptoms cannot be fully explained by another medical condition.

Abdominal migraine

All of the following criteria must be fulfilled for at least 6 months prior to diagnosis and on at least two occasions:[70]

- 1. Paroxysmal episodes of intense, acute periumbilical, midline, or diffuse abdominal pain lasting at least 1 hour. The abdominal pain must be the most severe and distressing symptom.
- 2. Episodes separated by weeks or months.
- 3. Pain is incapacitating and interferes with normal activities.
- 4. Stereotypical pattern and symptoms in the individual.
- 5. Pain associated with 2 or more of:
 - Anorexia
 - Nausea
 - Vomiting
 - Headache
 - Photophobia
 - · Pallor.
- 6. After appropriate evaluation the symptoms cannot be fully explained by another medical condition.

Functional abdominal pain - not otherwise specified

Three diagnostic criteria must be fulfilled at least four times per month, for 2 months prior to diagnosis:[70]

- 1. Episodic or continuous abdominal pain that does not occur solely during physiological events (e.g., eating, menstruation)
- 2. Insufficient criteria for irritable bowel syndrome, functional dyspepsia, or abdominal migraine diagnosis
- 3. After appropriate evaluation the symptoms cannot be fully explained by another medical condition.

Alarm features in children with chronic abdominal pain, which may indicate an organic or motility-related rather than a functional cause, include:[70] [74] [75]

- · Family history of inflammatory bowel disease, coeliac disease, or peptic ulcer disease
- Persistent right upper or right lower quadrant pain
- · Dysphagia
- Odynophagia
- · Persistent vomiting
- Gastrointestinal bleeding
- Nocturnal diarrhoea
- Arthritis
- · Peri-rectal disease
- · Involuntary weight loss
- · Deceleration of linear growth
- Delayed puberty
- · Unexplained fever.

Urgent considerations

(See **Differentials** for more details)

Although the causes of abdominal pain in children are frequently benign (e.g., constipation), there is always the potential for life- or organ-threatening conditions, which require urgent intervention.

Absent bowel sounds, bilious vomiting, bloody diarrhoea or occult blood in stool, fever (≥38.0 °C [≥100.4 °F]), rebound tenderness, rigidity, and guarding indicate a possible need for surgery.

Gastrointestinal (GI) emergencies

Acute appendicitis resulting in perforation

- Untreated acute appendicitis may progress to ischaemia, necrosis, and eventually perforation. The
 clinician may encounter a range of presentations. Patients often complain of abdominal pain localised
 to the right lower quadrant; in more severe cases the pain may be diffuse (e.g., if perforation results in
 generalised peritonitis).
- Perforation should be considered when a patient presents with a longer duration of symptoms and/ or suspected appendicitis with marked systemic signs of illness (e.g., high fever [>38.3°C, >101°F]), tachycardia, and anorexia), localised or generalised peritonitis with guarding, distended abdomen with absent bowel sounds. There may be a palpable mass if a peri-appendiceal abscess is present due to perforation.
- Refer all children with suspected appendicitis to the paediatric surgery team on call, if available.
 Where no paediatric surgery team is available, joint care should be managed between paediatrics and surgical teams.
- Ultrasound (US) is usually the preferred initial diagnostic study (due to low cost and lack of radiation exposure).[76] [77] [78] In children, specificity and sensitivity of ultrasound is similar to that of computed tomography (CT) (specificity 92.0% vs. 94.2%; sensitivity 90.3% vs. 93.0%); for MRI the specificity is 96.5% and sensitivity 96.9%.[78] However, all imaging modalities have limitations in differentiating perforated from non-perforated appendicitis.[76] [78][79]
- Treatment (antibiotics and possibly surgery) of acute or perforated appendicitis should be instituted promptly, but the condition is more urgent than emergent. Once antibiotics are administered, appendectomy for out-of-hours cases selected for operative management can be safely deferred until the following day.[13] [14]
- Appendectomy can be done with an open approach or laparoscopically. Laparoscopic appendectomy
 is associated with lower postoperative pain, lower incidence of surgical site infection, and higher
 quality of life in children.[13] Referral to a children's hospital or a paediatric surgeon should be
 considered for children younger than 5 years of age.[80]

Intestinal obstruction

- Urgency of intervention is dependent on the clinical severity of the obstruction.
- Strangulated obstructions are usually complete obstructions in which the blood supply to the bowel is cut off as a result of oedema, twisting of the bowel, or adhesions. These usually demonstrate diffuse or local peritonitis, fever, and leukocytosis. Untreated, they progress to intestinal necrosis and/or perforation. Urgent surgical treatment is mandatory.
- Non-strangulated obstructions involve a loop of bowel that is partially or completely obstructed but has an adequate blood supply and is not necrotic. This type of obstruction is usually not associated with peritonitis, fever, or leukocytosis, but may be associated with abdominal distension, nausea, and

vomiting. Although surgical intervention may be necessary, it is usually not urgent. However, prolonged delay may progress to strangulation.

Intussusception

- May lead to venous obstruction and bowel-wall oedema and can progress, if untreated, to bowel
 necrosis, perforation, and, rarely, death.[81] [82] Treatment should be initiated at the time of diagnosis.
 The goal is correction of hypovolaemia and electrolyte abnormalities, and antibiotic administration,
 followed by urgent reduction.
- Reduction can be accomplished with a radiographic enema study (air is preferred over contrast reagent) or by surgery.[83]
- There is a risk of recurrence of intussusception after reduction, especially within the first 48 hours, so prompt re-evaluation is necessary if symptoms recur.[84]

Volvulus

- Malrotation with mid-gut volvulus is a surgical emergency, and bilious vomiting in any child should prompt concern for this condition until confirmed otherwise.
- With a corresponding history and physical examination (bilious vomiting and feeding difficulty, especially in infants during the first month of life), no further diagnostic intervention is necessary, and prompt surgical exploration is recommended.
- Ambiguous cases may proceed to an upper GI contrast study (the gold standard test). However, this should not preclude surgical intervention if clinical suspicion is high.

Incarcerated hernia

- Prompt attention should be paid to an incarcerated inguinal or umbilical hernia due to the danger of bowel strangulation (compromise of blood flow to the bowel with consequent bowel ischaemia and gangrene). Incarceration, with or without strangulation, occurs if intra-abdominal contents become trapped in the protruding hernia sac.
- Clinically, the hernia is irreducible and tender. Associated symptoms may include nausea, vomiting, and generalised abdominal pain. In severe cases, fever, abdominal distension, and skin changes may be present.
- Peritonitis is a contraindication to attempted non-operative reduction.
- If strangulation is evident, surgery is required urgently to resect the gangrenous segment of bowel.

Necrotising enterocolitis

- The most common medical/surgical emergency affecting neonates, particularly premature infants, especially those weighing less than 1500 g.[35]
- Signs and symptoms include feeding intolerance, apnoea, lethargy, bloody stools, abdominal distension, tenderness, abdominal wall erythema, and bradycardia.
- Early intervention is mandatory to prevent morbidity and mortality due to multiple organ impairment.

 Treatment may be medical or surgical, and is determined by severity of the clinical presentation.

Abdominal trauma

- Most cases of blunt injury to the liver and spleen are managed non-operatively.
- Indications for urgent surgery include haemodynamic instability despite adequate resuscitation, free air in the abdomen, penetrating injuries with fascial penetration, and peritonitis.[85] [86]

Genitourinary emergencies

Ruptured ectopic pregnancy

- If undiagnosed or incorrectly managed, a ruptured ectopic pregnancy may lead to maternal death due to rupture of the implantation site and intraperitoneal haemorrhage.
- The classic presentation includes lower abdominal pain, amenorrhoea, and vaginal bleeding. Patients with a positive urine pregnancy test and the absence of an intrauterine pregnancy on transvaginal ultrasound are considered to have an ectopic pregnancy until confirmed otherwise.
- A quick and focused ultrasonographic examination to assess for the presence of free fluid or blood may be helpful when this diagnosis is suspected, but this should not delay other care.
- Haemodynamic instability associated with a ruptured ectopic pregnancy results from severe
 hypovolaemia secondary to blood loss. As such, the management of these patients involves
 stabilisation with emergency fluid resuscitation and immediate transfer to theatre. Rapid volume
 repletion with isotonic solution and blood products is of paramount importance to avoid ischaemic
 injury and multi-organ damage.
- Urgent laparoscopy with salpingectomy or salpingostomy is performed for a ruptured ectopic pregnancy.

Ovarian torsion

- Twisting or torsion of the ovary compromises the arterial inflow and venous outflow, producing ischaemia, which, if not relieved, can affect the viability of the ovary.
- It presents with acute-onset lower abdominal pain and, frequently, nausea and vomiting. Symptoms may be intermittent and fluctuate in severity.
- Doppler ultrasound is of only variable reliability in the diagnosis of ovarian torsion.[87] [88]
- It is not known how long an ovary can withstand ischaemia without permanent damage (it may be up to 72 hours or even longer), but definitive operative intervention should be undertaken as soon as possible.[89] [90] A delay in the timing of surgery is associated with a reduction in the possibility of ovarian salvage.[91]
- Conservative management with detorsion is highly recommended regardless of appearance of ovary.[60] [92] [93]

Testicular torsion

- Should be ruled out in any male child presenting with abdominal pain. The twisting of the testis and spermatic cord causes obstruction of arterial inflow and venous drainage from the testis.
- It typically presents with sudden-onset testicular pain; however, younger boys may only complain of abdominal tenderness, nausea, and/or vomiting.
- Physical findings suggestive of testicular torsion include loss of the cremasteric reflex, diffuse testicular tenderness, raised testes, and a horizontal rather than vertical position of the testes.
- Prompt recognition and early surgical intervention are necessary to prevent testicular loss. Manual detorsion may be attempted while preparations for surgery are being made.
- Duplex Doppler ultrasound is the definitive test for testicular torsion and should be obtained unless the
 test would unnecessarily delay intervention in a child for whom there is a high index of suspicion.[94]
 [95]

Sepsis

Sepsis and septic shock

- Patients with intra-abdominal sepsis may present with abdominal pain. [96]
- Sepsis is a spectrum of disease where there is a systemic and dysregulated host response to an infection.[97]
- Presentation ranges from subtle, non-specific symptoms (e.g., feeling unwell with a normal temperature) to severe symptoms with evidence of multi-organ dysfunction and septic shock. Patients may have signs of tachycardia, tachypnoea, hypotension, fever or hypothermia, poor capillary refill, mottled or ashen skin, cyanosis, newly altered mental state, or reduced urine output.[98]
- · Sepsis and septic shock are medical emergencies.
- In children, risk factors for sepsis include: age under 1 year, impaired immunity (due to illness or drugs), recent surgery or other invasive procedures, any breach of skin integrity (e.g., cuts, burns), and indwelling lines or catheters.[98]
- Early recognition of sepsis is essential because early treatment improves outcomes.[98] [99] [Evidence C] However, detection can be challenging because the clinical presentation of sepsis can be subtle and non-specific. A low threshold for suspecting sepsis is therefore important.
- The key to early recognition is the systematic identification of any patient who has signs or symptoms suggestive of infection and is at risk of deterioration due to organ dysfunction. Criteria to identify sepsis and septic shock in children and young people under the age of 18 years have been developed.[100] Several other risk stratification approaches exist. All rely on a structured clinical assessment and recording of the patient's vital signs.[98] [100][101] [102][103] It is important to check local guidance for information on which approach your institution recommends.
- The timeline of ensuing investigations and treatment should be guided by this early assessment.[103]
- Treatment guidelines have been produced by the Surviving Sepsis Campaign and remain the most widely accepted standards.[99] Within the first hour:[99]
 - Follow institutional protocols for management of sepsis/septic shock in children; these improve
 the speed and reliability of care.
 - Obtain blood cultures before administering antibiotics (provided this does not substantially delay antibiotic administration).
 - Administer broad-spectrum antibiotics.
 - Administer crystalloid fluids titrated to clinical signs of cardiac output and stopped if there is evidence of volume overload. Consult local protocols.
 - Use trends in blood lactate levels to guide resuscitation. If the child's hypotension is refractory to fluid resuscitation, consider use of vasopressors.

For more information on sepsis, please see our topic Sepsis in children .

Approach

Assessment of paediatric abdominal pain can prove a diagnostic challenge. Children may be limited in their ability to give an accurate history. Parents or guardians may also have difficulty interpreting the complaints of small children. In many cases, the causes are benign with few long-term sequelae. However, some require rapid diagnosis and treatment in order to prevent significant morbidity or mortality. Consideration of the child's age helps narrow the differential diagnoses.[75]

History

The clinician should determine early on whether the abdominal pain is acute or chronic in nature, as this will help indicate the urgency of treatment. Acute abdominal pain is usually a single episode that typically lasts from hours to days. The pain may vary in severity over time and is often localised and described as sharp and/or stabbing in nature. Conversely, chronic abdominal pain typically lasts days to weeks to months, and is usually dull, diffuse, and poorly localised. There may be pain-free intervals of variable duration, and when it recurs the pain may vary in intensity.

In addition, the history should cover the following:

Age of the child

The age of the child affects some of the likely differential diagnoses:[19] [104]

- Neonates, infants, and toddlers: may present with abdominal pain caused by congenital causes or diseases linked to prematurity, such as necrotising enterocolitis, Meckel's diverticulum, Hirschsprung's disease, volvulus, intestinal obstruction caused by congenital atresia, or stenosis. Intussusception is also common in this age group.
- School-age children: idiopathic constipation and infectious causes of pain are most common in this
 age group. Functional abdominal pain and abdominal migraine may also present in school-age
 children
- Adolescents: conditions related to menstruation, sexually transmitted infections, and pregnancy should be considered. Testicular torsion, ovarian torsion, irritable bowel syndrome, and inflammatory bowel disease are more common in this age group.

Pain history

- Useful mnemonics for abdominal pain are 'CLAIR PRADER': Characteristics, Location, Aggravating factors, Intensity, Relieving factors, Periodicity, Rhythmicity, Associations, Duration, Exacerbating factors, and Radiation; or 'SOCRATES': Site, Onset, Character, Radiation, Associated Symptoms, Time course, Exacerbating or relieving factors, Severity.
- Onset, frequency, timing, and duration of the pain episode.
- Whether the pain is localised or diffuse: right lower quadrant (RLQ) pain suggests appendicitis; epigastric pain suggests peptic ulcer disease; diffuse pain may indicate perforation or peritonitis.
 Poorly localised pain that improves with movement is likely to be visceral pain from hollow organs. Well localised pain that worsens with movement is likely to arise from parietal peritoneum, parietal muscle, or skin.[105]
- Whether the pain radiates or migrates between areas of the abdomen: abdominal pain radiating to the back is suggestive of cholecystitis or pancreatitis; pain that begins centrally and migrates to the RLQ is typical of appendicitis.

- Any factors that make the pain better or worse, such as movement, defecation, food, or medication: in cholelithiasis/cholecystitis pain often occurs after eating (particularly fatty foods); epigastric pain due to peptic ulcer disease is usually related to eating meals; painful defecation may indicate constipation.
- The character of the pain: pain associated with peptic ulcer disease is dull rather than burning in nature; sharp or stabbing pain is typical of appendicitis.
- An acute exacerbation with a history of chronic pain can confound the diagnosis. It is important to determine if the nature of the acute pain is different (in character, location, or other factors) from the background, chronic pain. It is easy to miss or overlook an unrelated acute pathology in a patient with a chronic pain history of any sort (e.g., acute appendicitis in children with long-standing functional bowel complaints). The clinician must be wary of this trap, and there is no substitute for experience.

Associated symptoms

The clinician should enquire about the presence and severity of:

- Fever, nausea, vomiting, anorexia (gastroenteritis, mesenteric adenitis)
- Diarrhoea (gastroenteritis)
- Fatigue or jaundice (viral hepatitis)
- · Lethargy, headache, photophobia (abdominal migraine)
- Cough, shortness of breath (pneumonia or empyema)
- Pain elsewhere (e.g., sudden-onset testicular pain suggests testicular torsion)
- Blood in stool (ulcerative colitis, necrotising enterocolitis, dysentery, haemolytic uraemic syndrome) or mucus in stool (suggests bacterial or parasitic infection)
- Blood or bile in vomitus (small bowel obstruction)
- Genitourinary symptoms: dysuria, frequency of micturition, and haematuria suggest a urinary tract infection (UTI); vaginal discharge is suggestive of pelvic inflammatory disease; current menstruation may be indicative of dysmenorrhoea.

Trauma history

• If there is a history of trauma, ascertain whether it is blunt or penetrating, and accidental or non-accidental.

Travel history

- Travel to a developing country increases risk of viral hepatitis infection and infectious gastroenteritis.
- Stooling patterns and dietary history
 - The presence of bilious vomiting without passage of stool/flatus suggests complete small bowel obstruction. Bilious vomiting with passage of stool or flatus suggests partial small bowel obstruction.
 - Infrequent bowel action or faecal incontinence is suggestive of constipation. Infants may extend their legs and squeeze anal and buttock muscles to prevent stooling; toddlers often rise up on their toes, shift back and forth, and stiffen their legs and buttocks.
 - In addition to daily stooling pattern, evaluation of constipation includes pertinent medical history, onset
 of constipation symptoms, severity, and any previous treatments. Dietary history, paying attention to
 fibre and fluid intake, is helpful.
 - New or unusual food intake may support the diagnosis of gastroenteritis.

Past medical history (including birth and development)

This should focus on previous operations, medication use, immunisations, allergies, and current comorbidities; for example:

- · Patients with sickle cell disease or cystic fibrosis are at higher risk of developing gallstones
- Patients with spina bifida, learning difficulties, or cerebral palsy are prone to constipation
- Splenic infarction may be a consequence of sickle cell disease
- Recent or current upper respiratory tract infection is suggestive of mesenteric adenitis or pulmonary cause
- · Failure to pass meconium in the first 36 hours of life is strongly suggestive of Hirschsprung's disease
- Hirschsprung's disease can be associated with Down syndrome
- Necrotising enterocolitis should be considered in a premature neonate weighing less than 1500 g.

Drug history

 Some medications, such as iron supplements, can cause constipation. Corticosteroids, adrenocorticotrophic hormones, oestrogens including contraceptives, azathioprine, asparaginase, tetracycline, chlorothiazides, and valproic acid may induce pancreatitis. Non-steroidal antiinflammatory drugs (NSAIDs), salicylates, and corticosteroids are associated with peptic ulcer development.

Family history

• Positive family history is a risk factor for inflammatory bowel disease, nephrolithiasis, and functional abdominal pain.

Social and psychiatric history

- This should include a discussion of family dynamics and may help determine if pain is functional or due to organic cause.
- Psychological factors (e.g., depression, abuse, attention deficit disorder, oppositional disorder), weaning, toilet training, start of schooling, or other causes of stress may play a role in constipation.
- History of nicotine, caffeine, or alcohol consumption may suggest peptic ulcer; excess alcohol consumption is a risk factor for pancreatitis.

Sexual history (in adolescents)

Adolescents may avoid answering sensitive questions regarding sexual history and drug use truthfully
in the presence of parents or guardians; therefore, it may be appropriate to conduct some parts of the
history with the adolescent alone.

Physical examination

This should be performed in a comfortable and non-threatening manner. Children may need to be distracted by parents or guardians in order to obtain an accurate examination. In younger children, localisation of the abdominal pain may be difficult.

Because paediatric abdominal pain may originate from other areas of the body (e.g., genitals or lungs), a comprehensive physical examination is necessary. Digital rectal examination is not routinely performed and is not needed to diagnose functional constipation.[106] UK guidelines recommend that digital rectal examination should only be performed by clinicians able to interpret features of anatomical abnormalities or

Hirschsprung's disease.[107] Pelvic examination is not routinely performed; history is usually a guide and digital examination should be reserved for adolescents who are sexually active.

All ages

Vital signs:

- Consideration of vital signs should be based on age-appropriate normal values.
- It is important to determine whether signs of volume depletion are present (tachycardia, hypotension, dry mucous membranes, poor capillary refill, sunken fontanelle in infants). Children with gastroenteritis may quickly become volume depleted.
- Fever may be the only presenting sign of a UTI, especially in the younger age group, and UTI should therefore be a top differential in children between 2 months and 2 years of age with fever.[108] High fever is suggestive of pyelonephritis.[108] [109]
- Cholecystitis, pancreatitis, and splenic infarction typically cause fever.
- Patients with appendicitis do not usually have significant changes in vital signs. Body temperature may be increased by 1°C (1.8°F).
- · Children with constipation are usually well with normal vital signs.

Abdominal examination:

- Patients presenting with central abdominal pain, with or without guarding and rigidity, that settles in
 the RLQ should arouse suspicion of appendicitis. Classic abdominal signs of appendicitis are RLQ
 abdominal tenderness (McBurney's sign) and localised rebound tenderness, if the appendix is anterior.
 Compressing the left lower quadrant (LLQ) may elicit pain in the RLQ (Rovsing's sign). Patients with
 appendicitis may lie still and try not to move, particularly in severe cases with significant peritoneal
 irritation.
- Acute mesenteric adenitis often resembles acute appendicitis; however, pain in the abdomen is usually diffuse with tenderness not localised to the RLQ. Guarding may be present but rigidity is usually absent. One retrospective study found that, compared with children who have appendicitis, patients who have mesenteric adenitis are more likely to have high fever (above 39°C) and dysuria, and are less likely to have migratory pain, vomiting, or typical abdominal signs of appendicitis on examination.[28]
- Patients with gastroenteritis usually exhibit diffuse abdominal pain without evidence of peritonitis (no guarding or rebound tenderness). Abdominal distension and hyperactive bowel sounds are common findings.
- The presence of abdominal distension and tenderness associated with decreased or absent bowel sounds is strongly suggestive of large bowel obstruction. In severe cases of constipation, abdominal distension may be present with a palpable faecal mass per abdomen or rectum.
- Flank pain or costovertebral angle tenderness may indicate pyelonephritis or nephrolithiasis.
- Epigastric pain may indicate peptic ulcer disease or pancreatitis. Patients with pancreatitis may lie with their knees and hips flexed and avoid moving. It is important to note that, in younger patients with pancreatitis (<3 years of age), abdominal tenderness may not be the main finding; these patients may demonstrate increased irritability and abdominal distension. With haemorrhagic pancreatitis, discoloration may be noted around the umbilical area (Cullen's sign) or in the flanks (Grey-Turner's sign) due to blood tracking along defined fascial planes.
- Tenderness in the right upper quadrant (RUQ) is a classical sign of gallbladder disease, as is Murphy's sign (cessation of inspiration during concurrent deep RUQ palpation). Patients with biliary dyskinesia usually present in a similar fashion to those with cholelithiasis and cholecystitis and may have RUQ tenderness on palpation.

- Patients with splenic infarction typically present with left-sided abdominal pain and fevers. Pain may also be reported in the left side of the chest or the left shoulder. Those with a splenic cyst are either asymptomatic or present with dull left-sided abdominal pain in the absence of fever.[110]
- In trauma patients, signs of accidental (e.g., seat belt mark suggesting a motor vehicle accident) and non-accidental injury (particularly if history is suspicious) should be sought (e.g., cigarette burns, subdural haemorrhages in an infant/young toddler). The presence of seat belt marks increases the likelihood that intra-abdominal injuries are present, particularly in the presence of lumbar fracture or persistent tachycardia.[111]
- If clinical findings are minimal and the child appears well, a diagnosis of functional abdominal pain should be considered. Diagnostic criteria for functional abdominal pain are symptom based, not physical examination or laboratory based.[71]
- Signs of peritonitis, such as absent bowel sounds, bilious vomiting, bloody diarrhoea or occult blood in stool, fever (≥38°C [≥100.4°F]), rebound tenderness, rigidity, and guarding indicate a possible need for surgery.

External genital and perineal examination:

- A neonatal examination must evaluate for the presence of an anus along with the proper location within the sphincter complex.
- Testicular torsion is likely in any male child with abdominal tenderness plus loss of the cremasteric reflex, diffuse testicular tenderness, raised testes, and a horizontal rather than vertical position of the testes on exam.
- Blood at the urethral meatus, or haematuria, after trauma may suggest urinary tract or kidney injury.
- The presence of an anal fissure and/or haemorrhoids (rare in children; may be mistaken for skin tags from Crohn's disease), imperforate anus, or anal stenosis (particularly in a neonate or infant) on inspection of the perianal skin may provide further diagnostic clues.

General examination:

- Generalised lymphadenopathy is common and signs of an upper respiratory tract infection may be present in children with mesenteric adenitis (e.g., hyperaemic pharynx or oropharynx suggesting pharyngitis).
- Jaundice is rare with cholelithiasis or acute cholecystitis and, if present, suggests an obstruction of the common bile duct. Jaundice associated with abdominal tenderness, hepatomegaly (splenomegaly may also be present), and lymphadenopathy, particularly in a child of school age, should arouse suspicion of viral hepatitis (commonly hepatitis A).
- Children with spina bifida, learning difficulties, and cerebral palsy are prone to constipation, and features of these conditions may be obvious on examination (e.g., sacral dimples or pits and/or tags/tufts indicative of abnormality of spinal cord).
- Henoch-Schonlein purpura (HSP) may be the initiating factor in an older child with abdominal pain (usually <11 years of age), and therefore signs of HSP should be sought (rash of palpable purpura, blood in the stools).
- Extra-intestinal manifestations of inflammatory bowel disease may be evident (e.g., iritis, arthritis, sacroiliitis, erythema nodosum, pyoderma gangrenosum).
- The presence of cyanosis, tachypnoea, decreased breath sounds on auscultation, dullness on percussion (indicates consolidation), and abdominal tenderness and distension without guarding or rebound should arouse suspicion of a pulmonary cause such as pneumonia or empyema.

Infants and toddlers

- In a neonate, the triad of abdominal distension, delayed passage of meconium (not occurring in the first 36 hours of life), and vomiting is highly suggestive of Hirschsprung's disease.
- Necrotising enterocolitis should be considered in a premature neonate weighing less than 1500 g.
 Early signs may include inability to tolerate feeds, abdominal distension and tenderness, blood in the stool, and abdominal wall erythema. In severe cases, systemic signs of sepsis may be present.
- A neonate presenting with bilious vomiting, with (partial obstruction) or without (complete obstruction)
 the passage of meconium, is highly suggestive of small bowel obstruction. Causes such as meconium
 ileus, intestinal atresia, and mid-gut volvulus should be excluded with further investigations.
- Any abdominal examination of an infant and toddler must include a check for inguinal hernias; if a nappy is worn, remove it, as otherwise a hernia may be missed.
- Meckel's diverticulum should be considered in a child <2 years old with abdominal tenderness
 (Meckel's diverticulitis); haematochezia, typically dark red, maroon, or red-brick-coloured jelly-like
 stools (indicates intestinal bleeding as they contain heterotopic gastric tissue); or signs of obstruction
 such as nausea, vomiting, and constipation (intussusception, volvulus, or herniation can result).

Reproductive age

- Ectopic pregnancy and miscarriage should be suspected in any female of reproductive age presenting
 with lower abdominal pain, amenorrhoea, and vaginal bleeding. Pelvic examination may reveal a
 mass, eliciting cervical motion tenderness if haemoperitoneum is present; tubal rupture can cause
 haemodynamic instability.
- Clinical features of a ruptured ovarian cyst usually occur prior to the expected time of ovulation and may mimic ectopic pregnancy. Pain arises from local peritonitis secondary to haemorrhage.[56] [57]
 [58] Signs of peritonitis may be present in the lower abdomen and pelvis; adnexal size is unremarkable due to collapsed cyst.
- The presence of a tender pelvic mass associated with nausea and vomiting may suggest ovarian torsion. In addition, in patients old enough to undergo pelvic examination, cervical motion tenderness may be elicited; typically no vaginal discharge is present, but there may be some mild to moderate vaginal bleeding.
- Physical findings of pelvic inflammatory disease (PID) vary widely and may include lower abdominal tenderness, adnexal tenderness, and cervical motion tenderness.[62] Fever and cervical or vaginal discharge may also be present. PID is rare in the absence of sexual activity; PID in a young child should prompt work-up for possible sexual abuse.
- Patients with PID may also present with RUQ pain resulting from inflammation of the liver capsule or diaphragm, referred to as Fitz-Hugh-Curtis syndrome. This is secondary to an ascending infection.
 Referred pain to the right shoulder may result from irritation of the diaphragm.[112]
- Primary dysmenorrhoea should be considered if lower abdominal tenderness is associated with current menstruation.

Laboratory tests

Challenges in the clinical assessment of abdominal pain in the paediatric patient mean laboratory and imaging studies can play an important role.

Urinalysis is essential to exclude underlying UTI or haematuria (associated with nephrolithiasis, UTI, haemolytic uraemic syndrome, urinary tract or kidney injury) and should be performed in children of all ages presenting with abdominal pain. For females of reproductive age a urine pregnancy test and/or serum human chorionic gonadotrophin (beta-hCG) is necessary to exclude miscarriage and ectopic pregnancy.

Initial blood and stool tests may include:

- FBC, recommended in all patients (useful in assessing infection and inflammation).
- Complete chemistry panel, recommended in all patients (electrolyte disturbances associated with GI causes are common).
- Blood type and screen and rhesus status when ectopic pregnancy is suspected.
- Liver function tests (LFTs) are helpful baseline investigations when considering a hepatobiliary or pancreatic cause. In abdominal trauma, results from one retrospective study suggest that LFTs may be used as a screening tool to determine need for computed tomography (CT) scan; in a haemodynamically normal patient with normal AST/ALT, it may be possible to avoid CT scan.[113]
- Serum lipase, or amylase if lipase is unavailable, is indicated if pancreatitis is suspected. [114]
- Inflammatory markers. Although non-specific, erythrocyte sedimentation rate and C-reactive protein may suggest underlying infection or inflammation. Furthermore, these inflammatory markers correlate closely with disease activity in cases of inflammatory bowel disease.
- A positive faecal occult blood test can support a suspicion of intussusception. However, a negative test cannot reliably rule out the diagnosis.[115]
- Faecal calprotectin is used in the diagnosis and monitoring of inflammatory bowel disease and to distinguish it from functional gastrointestinal disorders.[116]
- A coagulation profile, including prothrombin time and international normalised ratio (INR), is usually necessary in cases of suspected viral hepatitis to measure liver synthetic function.

Microbiological investigations

- Stool microscopy and culture may be helpful in determining an infectious aetiology of gastroenteritis.
 Risk factors and features of the clinical presentation help guide the choice of tests for specific pathogens. Guidelines recommend that when there is fever or bloody diarrhoea, investigations for enteropathogens for which antimicrobial agents may confer clinical benefit (including Salmonella enterica subspecies, Shigella, and Campylobacter) should be done.[117]
- Blood cultures are indicated when sepsis is a concern. Blood cultures are recommended: in children with infectious diarrhoea who are <3 months of age or who are immunocompromised; when enteric fever is suspected (including travel to enteric fever-endemic areas, or contact with travellers from enteric fever-endemic areas who have a febrile illness of unknown aetiology); when there are systemic manifestations of infection; and with high-risk conditions such as haemolytic anaemia.[117] [118]
- Urine culture is necessary if urinalysis is suggestive of a UTI.
- Sputum culture is indicated in patients with suspected pneumonia. Aspiration of frank pus on thoracentesis is diagnostic of empyema.
- In cases of patients with suspected peptic ulcer disease, *Helicobacter pylori* breath test or stool antigen test may be helpful.[119] [120]
- Serological markers (perinuclear anti-neutrophil cytoplasmic antibody and anti-saccharomyces cerevisiae antibody) may be particularly useful for differentiating between Crohn's disease and ulcerative colitis in the paediatric population.[121] [122]
- Polymorphonuclear leukocytes (PMNs) seen on wet mount of vaginal secretions confirms vaginal
 infection in cases of PID. All women who receive a diagnosis of PID should be tested for gonorrhoea,
 chlamydia, HIV, and syphilis.[62] Hepatitis studies may be considered.
- In patients with suspected exposure to or symptoms of hepatitis A, B, C, D, and E, the following laboratory tests are warranted: hepatitis A antibody IgM, hepatitis B surface, core, and e antigen or viral load, hepatitis C serology or viral load, hepatitis D and E serologies.

Imaging and other investigations

Imaging studies are guided by history and physical examination findings.

Abdominal and chest x-ray

Plain abdominal x-rays are often non-specific but may suggest the presence of an obstruction; faecal impaction and duodenal atresia can be detected on x-ray. In addition, if sufficiently radiopaque, it may be possible to identify gallstones or urinary stones on a plain abdominal film. This is often the initial test, as it can be performed quickly. Supine and upright films are usually requested. Free air under the diaphragm suggests perforation and requires immediate surgical assessment. Chest x-ray should be ordered if perforation is suspected or a respiratory cause such as pneumonia or empyema is likely.

Ultrasound

Ultrasound scans avoid radiation exposure and are typically better tolerated than other imaging modalities such as computed tomography (CT) scan. An experienced paediatric radiologist may be able to utilise ultrasound in all regions of the body (abdomen, chest, testicles). Abnormalities detected by ultrasound may not be causally related to the patient's abdominal pain.[74]

Ultrasound is usually recommended as the first imaging test for suspected appendicitis.[76] [78] Ultrasound scans are considered particularly useful in assessing pain in the RUQ (i.e., gallbladder disease), and lower abdominal pain (pelvic pain) in females.

Colour Doppler may be helpful in determining ovarian blood flow in cases of suspected ovarian torsion, but is of only variable reliability in the diagnosis.[87] [88] Testicular ultrasound (using duplex Doppler ultrasound) should be able to diagnose testicular torsion and provide information on the vascular integrity of the testis.

Ultrasound of the urinary tract (including kidneys) is of benefit when wanting to exclude anatomical abnormalities (e.g., when UTI is present) or nephrolithiasis and associated complications such as hydronephrosis.

Point-of-care ultrasound may be useful for detecting intussusception at an early stage.[123] [124]

Focused abdominal sonography for trauma (FAST) may be useful in children with blunt abdominal trauma who are haemodynamically unstable.[125] The presence of large amounts of free fluid indicates a need for immediate operative intervention. FAST in haemodynamically stable patients has a less certain impact, as a negative scan does not preclude injury. A large multi-institutional study at paediatric trauma centres demonstrated poor FAST sensitivity (28.6%) in normotensive patients.[126] Though specificity was high (91.1%), results of FAST rarely changed management. A randomised trial comparing FAST with standard care in haemodynamically stable children and adolescents with blunt torso trauma also found no significant difference in the proportion of abdominal CT scans, missed intra-abdominal injuries, length of stay in the accident and emergency department, and median hospital costs.[127]

Abdominal CT scan with intravenous contrast is the diagnostic test of choice for the identification of solid organ injuries, especially to the liver, kidney, and/or spleen. Oral contrast is usually not necessary when scanning a patient for trauma.

CT and magnetic resonance imaging (MRI)

CT with contrast may be appropriate in investigating suspected appendicitis if ultrasound is non-diagnostic; however, radiation exposure with its risk of malignancy should be considered.[78] [128] [129] MRI avoids

ionising radiation and has similar or better sensitivity and specificity than CT, though availability and longer scan times may limit its utility.[78] [130] [131] Increasingly, MR enterography (MRE) is being used to evaluate paediatric Crohn's disease patients. This tool can evaluate extraluminal and extra-intestinal manifestations of Crohn's disease as well as the status of the bowel wall and presence of possible stricture formation.[132]

In general, for investigation of abdominal pain in children, CT scan of the abdomen and pelvis, with or without contrast, may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children.

GI contrast studies (upper-GI or barium enema studies) are routinely used in place of CT scans, especially in infants. In assessment of intestinal obstruction, clinical suspicion should direct which contrast study should be performed first and will be most informative.

Endoscopy

Endoscopy (oesophagogastroduodenoscopy or colonoscopy) with biopsy may be required to assess mucosa-based diseases, such as peptic ulcer disease and inflammatory bowel disease. Rectal biopsy and anorectal manometry help confirm the diagnosis of Hirschsprung's disease. These procedures may be particularly challenging when performed in a newborn.

Nuclear medicine scans

Diagnosis of biliary dyskinesia is confirmed by hepatobiliary iminodiacetic acid (HIDA) scan with an ejection fraction <35%. It is important to note that the cut-off of <35% is not universally accepted; some consider an ejection fraction <15% to be more predictive of success of surgical treatment (i.e., cholecystectomy).[134] Hyperkinetic results (very high ejection fractions) are sometimes considered an indication for cholecystectomy, as are normal scans (at some institutions). Outcomes vary widely among paediatric biliary dyskinesia patients. In retrospective studies, overall rates of symptom resolution post operation ranged from 34% to 100%.[135] [136] The absence of reliable factors to predict which patients will benefit is a source of frustration to patients, families, and physicians. HIDA scan is considered a useful adjunct in the diagnosis of cholecystitis.

Technetium-99m pertechnetate scan is considered the most useful method to diagnose a suspected Meckel's diverticulum; this scan identifies ectopic gastric mucosa as tracer is taken up by gastric mucin-producing cells.

Voiding cystourethrogram

The American Academy of Pediatrics (AAP) recommends that a voiding cystourethrogram (VCUG) is indicated in children between 2 and 24 months of age following an initial UTI if renal and bladder ultrasonography reveals hydronephrosis, scarring, or other findings that would suggest either high-grade vesicoureteral reflux or obstructive uropathy, as well as in other atypical or complex clinical circumstances.[137] Further evaluation is required if there is a recurrence of febrile UTI.[137] [138][139]

Full skeletal x-rays

Skeletal survey identifies previous skeletal injuries and should be obtained if there is a high suspicion of non-accidental trauma. This should only be performed to provide adjunctive diagnoses to support an initial injury and suspicion of abuse.

Diagnostic Iaparoscopy

Laparoscopy may be necessary to confirm diagnosis of pelvic inflammatory disease as it allows direct visualisation of the gynaecological and abdominal structures.



Abdominal x-ray of a neonate with abnormal stooling pattern and constipation. The dilated transverse and descending colon is suggestive of Hirschsprung's disease

From the collection of Dr KuoJen Tsao; used with permission



Abdominal x-ray demonstrating double bubble gas pattern consistent with duodenal atresia


From the collection of Dr KuoJen Tsao; used with permission




Abdominal x-ray with opacities in the RUQ consistent with gallstones From the collection of Dr KuoJen Tsao; used with permission



Gallbladder ultrasound demonstrating cholelithiasis with characteristic shadowing



CT scan demonstrating faecalith (white arrow) outside the lumen of the appendix consistent with perforated appendix From the collection of Dr KuoJen Tsao; used with permission



CT scan of teenage girl presenting with mid-epigastric abdominal pain as a result of gallstone pancreatitis. The large fluid collection in the pancreatic bed (white arrow) and lack of pancreatic enhancement suggest liquefactive necrosis of the pancreas

From the collection of Dr Kuojen Tsao; used with permission



Contrast enema demonstrating ileocolic intussusception (black arrow)
From the collection of Dr KuoJen Tsao; used with permission

Differentials overview

Common
Constipation
Acute appendicitis
Gastroenteritis
Urinary tract infection
Abdominal trauma (blunt or penetrating)
Cholelithiasis/cholecystitis
Primary dysmenorrhoea
Pneumonia
Functional abdominal pain
Infantile colic
Uncommon
Uncommon Intussusception
Intussusception
Intussusception Meckel's diverticulum
Intussusception Meckel's diverticulum Mesenteric adenitis
Intussusception Meckel's diverticulum Mesenteric adenitis Hirschsprung's disease
Intussusception Meckel's diverticulum Mesenteric adenitis Hirschsprung's disease Ulcerative colitis
Intussusception Meckel's diverticulum Mesenteric adenitis Hirschsprung's disease Ulcerative colitis Crohn's disease
Intussusception Meckel's diverticulum Mesenteric adenitis Hirschsprung's disease Ulcerative colitis Crohn's disease Small bowel obstruction

Uncommon
Peptic ulcer disease
Coeliac disease
Viral hepatitis
Biliary dyskinesia
Acute pancreatitis
Splenic infarction/cysts
Nephrolithiasis
Sepsis
Testicular torsion
Ovarian torsion
Ruptured ovarian cyst
Pelvic inflammatory disease (PID)
Pregnancy complications
Empyema

Differentials

Common

♦ Constipation

History

poor diet and fluid intake; history of cerebral palsy, learning difficulties, or spinal cord problems; psychological factors (e.g., depression, abuse, ADHD, autism, oppositional disorder), weaning, toilet training, start of schooling or other causes of stress may be present; vague abdominal pain, painful defecation (infants may extend their legs and squeeze anal and buttock muscles to prevent stooling; toddlers often rise up on their toes, shift back and forth, and stiffen their legs and buttocks), faecal incontinence; medication with known constipating agents (e.g., iron supplements); obesity, low birth weight

Exam

examination findings may be minimal (mild abdominal tenderness. stool in rectum): abdominal distension in severe cases or in small children; faecal mass palpable on abdominal examination; absence of peritonitis (guarding or rebound tenderness); sacral dimples or pits and/or tags/tufts indicative of spinal cord abnormality (i.e., spina bifida); anal fissure, haemorrhoids (rare in children; may be mistaken for skin tags from Crohn's disease); imperforate anus or anal stenosis

1st Test

"none: clinical diagnosis Characteristic history and examination findings are often sufficient to diagnose the condition.

Other tests

»abdominal x-ray: stool visible throughout colon Stool throughout the colon does not rule out other aetiologies of abdominal pain.



Abdominal x-ray of a young boy with acute, severe abdominal pain, demonstrating stool throughout the colon and rectum

From the collection of Dr KuoJen Tsao; used with permission

»radiopaque marker colonic transit study: slow colonic transit, as measured by delay in marker movement, confirms constipation

History

r-Acute appendiciti

history of sharp or stabbing periumbilical pain that migrates to the right lower quadrant (RLQ); anorexia, fever, vomiting, and/ or diarrhoea may be present; occurs in all age groups but is rare in infants

Exam

patient lies still, tries not to move (especially in severe cases with significant peritoneal irritation); positive McBurney's sign (RLQ pain and tenderness to palpation at a point two-thirds along a line from the umbilicus to the anterior superior iliac spine); positive Rovsing's sign (pain in the RLQ in response to left-sided palpation, suggesting peritoneal irritation); positive psoas sign (pain in the RLQ when child placed on left side and right hip gently hyperextended, suggesting irritation to the psoas fascia and muscle); positive obturator sign (RLQ pain on internal rotation of the flexed right thigh); rectal tenderness and/or palpable abscess in **RLQ**

1st Test

- »FBC: normal or raised WBC
 May see leukocytosis with neutrophilia.
 However, a normal WBC count does not exclude appendicitis.
- »C-reactive protein (CRP): likely to be raised CRP ≥10 mg/L (≥1 mg/ dL) is a strong predictor of acute appendicitis in children <6 years old.[142]
- »urinalysis: normal If positive for red cells, white cells, or nitrates an alternative diagnosis such as renal colic or urinary tract infection should be considered.

The specific gravity of urine can sometimes suggest volume status.

»urine pregnancy test: negative In adolescent girls, a pregnancy test should be performed to exclude ectopic pregnancy.

Other tests

- »abdominal ultrasound: dilated appendix, free fluid; appendicolith may be present Ultrasound is the preferred method of assessment.[76] In children, specificity of ultrasound is similar to that of CT (0.92 vs. 0.94).[78] It may elicit a sonographic McBurney's sign (pain on compression of the appendix that is visualised on ultrasound). A negative ultrasound does not necessarily rule out appendicitis as a cause of abdominal pain. Subsequent investigations for paediatric patients with an equivocal ultrasound include CT or MRI.[78] [129]
- »CT scan abdomen and pelvis: dilated appendix, free fluid, mesenteric stranding, or appendicolith; abscess or phlegmon consistent with perforated appendicitis Useful when body habitus of patient makes ultrasound difficult or when ultrasound and/or clinical assessment

PAcute appendicitis

History	Exam	1st Test	Other tests
			are inconclusive. CT has greater sensitivity than ultrasound (0.93 vs. 0.90).[78] Radiation exposure and contrast burden should be considered.[128] [133] However, use of IV contrast, when feasible, is recommended for patients undergoing CT scan.[129] Sedation or general anaesthesia may be required in some children.
			CT scan
			demonstrating
			faecalith (white
			arrow) outside
			the lumen of
			the appendix
			consistent with
			perforated appendix
			From the collection
			of Dr KuoJen Tsao;
			used with permission

PAcute appendicitis

History	Exam	1st Test	Other tests
			CT scan
			demonstrating intra-
			abdominal abscess
			consistent with
			perforated appendix
			From the collection
			of Dr KuoJen Tsao;
			used with permission
			»MRI scan abdomen and pelvis: dilated appendix; hyperintensity of the luminal contents of the appendix, periappendiceal tissue and thickened wall In children, MRI has excellent sensitivity and specificity (0.97); however, there are fewer studies evaluating its test characteristics.[78] [131][143] It is increasingly used due to the potential hazards of ionising radiation from CT.

History	Exam	1st Test	Other tests
vague abdominal pain with nausea and vomiting; diarrhoea	diffuse abdominal pain without evidence of peritonitis (no	»none: clinical diagnosis	»serum electrolytes: normal or low sodium and potassium

History	Exam	1st Test	Other tests
with or without mucus in stool; recent travel or contact with sick individual(s) or ingestion of suspected food and drink; >10 days suggests parasitic or non-infectious cause; fever, chills, myalgia, rhinorrhoea, upper respiratory symptoms	guarding or rebound tenderness); abdominal distension; hyperactive bowel sounds; mucus in stool (bacterial or parasitic); signs of volume depletion (tachycardia, hypotension, dry mucous membranes, poor capillary refill, sunken fontanelle in infants); low-grade fever, lethargy and/ or irritability, reduced response to noxious stimuli, abnormal temperature (raised or low)		Measurements of serum electrolytes, urea, and creatinine are usually not necessary to assess children with acute gastroenteritis, as the results do not change the therapeutic strategy. These tests should only be considered in a subset of patients with severe dehydration receiving intravenous therapy. "urea and creatinine: normal; may have evidence of renal failure in patients with haemolytic uraemic syndrome Measurements of serum electrolytes, urea, and creatinine are usually not necessary to assess children with acute gastroenteritis, as the results do not change the therapeutic strategy. These tests should only be considered in a subset of patients with severe dehydration receiving intravenous therapy. "stool microscopy and culture: faecal leukocytes; ova or parasites; culture positive for infectious agent in bacterial gastroenteritis

History	Exam	1st Test	Other tests
			Faecal leukocytes and stool culture may be helpful in children presenting with dysentery and to demonstrate invasive pathogens such as Clostridium difficile. Ova and parasite studies are useful if history suggests camping, travel to other countries, or long-term disease. In patients who have taken previous antibiotics or have contact with an individual with C difficile diarrhoea, it is important to test for C difficile toxin. Untreated C difficile -related diarrhoea can have serious consequences (e.g., toxic megacolon).
			»urine dipstick: may detect presence of albumin or blood in haemolytic uraemic syndrome
			»FBC: variable Eosinophilia with viral or parasitic aetiology, peripheral eosinophilia suggests eosinophilic
			gastroenteritis; bandaemia with bacterial aetiology; anaemia and/or
			thrombocytopenia if

History	Exam	1st Test	Other tests
			haemolytic uraemic
			syndrome.
			Ordered to help
			differentiate alternativ
			disease processes.
			However, in true
			gastroenteritis, only
			indicated in patients
			with systemic infectio
			»blood culture:
			may be positive for
			infectious agent in presence of sepsis
			Important to order if
			concern for sepsis.
			Guidelines recommer
			blood cultures:
			in children with
			infectious diarrhoea
			who are <3 months
			of age or who are
			immunocompromised
			when enteric fever is
			suspected (including
			travel to enteric fever
			endemic areas, or
			contact with travellers
			from enteric fever-
			endemic areas who
			have a febrile illness
			of unknown aetiology
			when there are
			systemic manifestation
			of infection; and with
			high-risk conditions
			such as haemolytic
			anaemia.[117] [118]
			»endoscopy with

♦ Gastroenteritis

History	Exam	1st Test	Other tests
			Indicated only when eosinophilic gastroenteritis is a concern (condition affecting the gastrointestinal tract with eosinophil-rich inflammation without a known cause of the eosinophilia).[17]

□ Urinary tract infection

History	Exam	1st Test	Other tests
neonates and infants: fever, vomiting, lethargy, irritability, and poor feeding; older children: dysuria, urinary frequency and urgency, back pain if pyelonephritis	variable; fever >39°C (>102.2°F); suprapubic and/or costovertebral angle tenderness; irritability; foul-smelling urine; gross haematuria	»urine dipstick: positive leukocyte esterase and/or positive nitrite Urinalysis should be performed within 1-2 hours of obtaining specimen.[144] First morning voids may be best for yielding a positive nitrite test. Positive leukocyte esterase alone: sensitivity 84%, specificity 77%, likelihood ratio (LR)+ 5:5, LR- 0:26.[145] [146] Positive nitrite alone: sensitivity 58%, specificity 99%, LR+ 15:9, LR- 0:51.[145] [146] This test has a high positive predictive value.	»renal ultrasound: abnormalities may be present such as dilatation of the renal pelvis or ureters, or distension of thick- walled bladder; renal abscess: area of radiolucency to the renal parenchyma with local hypoperfusion on colour Doppler; perinephric abscess: hypoechoic fluid Initially performed to look for any anatomical abnormalities of the urinary tract. Also may be performed to look for evidence of a renal or perinephric abscess when the urinalysis and culture are negative but abdominal pain and fever persist. »voiding cystourethrogram

PUrinary tract infection

History	Exam	1st Test	Other tests
		Positive for either leukocyte esterase or nitrite: sensitivity 92%, LR- 0:2.[145] [146] This test is best at ruling out disease. Positive for leukocyte esterase and nitrite: LR+ 28:2.[146] This test is best at ruling in disease. **wrine microscopy: >4 WBC per high-power field or any bacteria Microscopic analysis is more accurate; however, a urine dipstick analysis may suffice. UK guidelines recommend sending a sample for urine microscopy from all infants under 3 months, before antibiotics are given.[147] Pyuria (the presence of WBC): sensitivity 78%, specificity 87%; LR- 0:27.[145] [146] Bacteriuria: sensitivity 88%, specificity 93% LR+ 14:7, LR- 0:19.[145] [146] Centrifugation reduces the specificity of these tests. **wrine culture: suprapubic aspirate:	(VCUG): if vesicoureteral reflux is present: contrast seen ascending out of the bladder into the upper urinary tract The American Academy of Pediatrics (AAP) recommends a VCUG in children between 2 and 24 months of age following an initial urinary tract infection (UTI) if renal and bladder ultrasonography reveals hydronephrosis scarring, or other findings that would suggest either high- grade vesicoureteral reflux or obstructive uropathy, as well as in other atypical or complex clinical circumstances.[137] Further evaluation is required if there is a recurrence of febrile UTI.[137][138] [139]

□ Urinary tract infection

History	Exam	1st Test	Other tests
		>1000 colony-forming units (CFU)/mL; catheter: >10,000 CFU/mL; clean-catch midstream: >100,000 CFU/mL Urine collected from urinary bags is likely to be contaminated.	

Abdominal trauma (blunt or penetrating)

1st Test Other tests History Exam history of trauma: abdominal tenderness: »FBC: may be normal »chest x-ray: may have multiple skin marks reflecting or show decreased may be normal or complaints; history may mechanism of injury haematocrit and show compatible suggest child abuse or (e.g., seat belt mark); haemoglobin thoracic injury (e.g., referred left shoulder non-accidental trauma pulmonary contusion, Patients with acute-(e.g., inconsistent pain (due to splenic pneumothorax); free onset haemorrhage or changing history, injury); blood at the air under diaphragm may have normal history not consistent urethral meatus, or (suggests perforation) haematocrit and with injuries/ haematuria (indicate »abdominal examination) urinary tract or kidney haemoglobin values. ultrasound: variable; injury); signs of nonmay show free fluid in accidental trauma »abdominal CT scan abdominal cavity may be present with intravenous Focused abdominal (e.g., cigarette burns, contrast: variable sonography for trauma subdural haemorrhages This is the diagnostic in an infant/young (FAST) may be useful test of choice for toddler) in patients with blunt the identification of abdominal trauma who solid organ injuries, are haemodynamically especially to liver, unstable.[125] kidney, and/or spleen. Presence of large In abdominal trauma. amounts of free fluid results from one indicates need for retrospective study immediate operative suggest that LFTs may intervention. FAST in be used as a screening haemodynamically tool to determine stable patients with need for CT scan; in blunt abdominal trauma a haemodynamically has a less certain normal patient with impact as a negative normal AST/ALT, it may

► Abdominal trauma (blunt or penetrating)

History	Exam	1st Test	Other tests
		be possible to avoid CT scan.[113]	scan does not preclude injury.[126] »full skeletal x-rays: variable Skeletal survey identifies previous skeletal injuries and should be obtained if there is a high suspicion of non-accidental trauma. This should only be performed to provide adjunctive diagnoses to support an initial injury and suspicion of abuse.

♦ Cholelithiasis/cholecystitis

History	Exam	1st Test	Other tests
recurrent, episodic right upper quadrant (RUQ) pain, may radiate to the back and is classically colicky in nature; often occurs after eating, particularly fatty foods; nausea, vomiting, and anorexia may be present; persistent pain and fever may signify acute cholecystitis; referred pain to right shoulder can occur; presence of risk factors (e.g., sickle cell disease, cystic fibrosis)	right subcostal region tenderness; positive Murphy's sign (during palpation, deep inspiration causes pain to suddenly become worse and produces inspiratory arrest); palpable distended, tender gallbladder; fever suggests acute cholecystitis; jaundice rare and suggests common bile duct obstruction	»RUQ ultrasound: gallstones; ductal dilation, thickened gallbladder wall (>4 mm); pericholecystic fluid; may also see ultrasonographic Murphy's sign »liver tests: may see raised alk phos, bilirubin, and aminotransferase May be normal or marginally increased; if significantly raised, other aetiologies (such as hepatitis, choledocholithiasis,	»abdominal x-ray: opacities in RUQ consistent with gallstones If sufficiently radiopaque, it may be possible to identify gallstones on a plain abdominal film.

♦ Cholelithiasis/cholecystitis

History	Exam	1st Test	Other tests
		or cholangitis) may be present. It is important to note the direct bilirubin concentration, which is the fraction that is raised with common duct obstruction. "FBC: normal WBC (suggests cholelithiasis) or leukocytosis (suggests acute cholecystitis) "C-reactive protein: normal (suggests acute cholelithiasis) or raised (suggests acute cholecystitis)	Abdominal x-ray with opacities in the RUQ consistent with gallstones From the collection of Dr KuoJen Tsao; used with permission *hepatobiliary iminodiacetic acid (HIDA) scan: non- filling gallbladder Considered a useful adjunct in the diagnosis of cholecystitis. The classical finding is non- filling of the gallbladder. Prolonged fasting may result in a false-positive test.

♦ Primary dysmenorrhoea

History	Exam	1st Test	Other tests
history of recurrent crampy abdominal pain associated with menstruation	lower abdominal tenderness; normal pelvic examination	»none: diagnosis is clinical	»abdominal/pelvic ultrasound: normal; however, useful to rule out other diagnoses

◊ Pneumonia

History	Exam	1st Test	Other tests
cough; purulent sputum production; upper respiratory tract symptoms (rhinorrhoea, sore throat, nasal congestion), shortness of breath, fever, and chills; splinting secondary to pain; vomiting, diarrhoea, anorexia	tachypnoea, cyanosis, decreased breath sounds, crackles/ rales on auscultation, dullness on percussion; abdominal tenderness and distension without guarding or rebound	»FBC: variable In patients with viral pneumonia, WBC count may be normal or decreased. »chest x-ray: infiltration, consolidation, effusion »sputum culture: growth of infecting organism	»chest ultrasound: localised fluid collection May consider before proceeding to CT scan due to the lack of radiation exposure. »CT scan chest with intravenous contrast: consolidation of lung parenchyma; extraparenchymal fluid with loculations suggests empyema CT scan may better differentiate between empyema and simple effusion. Radiation exposure should be considered. Sedation or general anaesthesia may be required in some children.

♦ Functional abdominal pain

History	Exam	1st Test	Other tests
history may be acute, chronic, or cyclic (frequently girls aged 8-12 years), complaint of vague, persistent, central abdominal pain common, may be associated nausea and vomiting, particularly in chronic cases; family history of functional disorders common (e.g., irritable bowel syndrome, anxiety, psychiatric disorders, and migraine); Rome IV criteria use symptoms for diagnosis	periumbilical tenderness, abdomen is soft, non-distended, no guarding or rebound tenderness; examination of other systems normal	»none: diagnosis is clinical after exclusion of possible organic causes	»FBC: normal »erythrocyte sedimentation rate: normal »urinalysis: normal »stool microscopy: normal

♦ Infantile colic

History	Exam	1st Test	Other tests
paroxysms of uncontrollable crying in otherwise healthy and well-fed infant aged <5 months; duration of crying is >3 hours per day, and >3 days per week, for at least 3 weeks; sibling(s) may have history of infantile colic	examination findings may be minimal; infant typically well and thriving	»none: clinical diagnosis Characteristic history and examination findings are often sufficient to diagnose the condition.	"urinalysis: normal Rarely required. Ordered only when the diagnosis is in doubt and urinary tract infection is suspected, when it may show pyuria and bacteriuria. "urine culture: normal Rarely required. Ordered only when the diagnosis is in doubt and urinary tract infection is suspected, when it may be positive.

Uncommon

PIntussusception

History	Exam	1st Test	Other tests
usually infant between 3 months and 12 months of age presenting with colicky abdominal pain, flexing of the legs, fever, lethargy, and vomiting; Henoch-Schonlein purpura (HSP) may be initiating factor in an older child (usually <11 years of age); vague abdominal complaints; severe, cramp-like abdominal pain; child may be inconsolable	may see gross or occult blood that may be mixed with mucus and have red-brick-coloured jelly-like appearance, abdominal tenderness, and palpable abdominal mass; signs of HSP may be present in older child (rash of palpable purpura, blood in the stools)	"abdominal ultrasound: tubular mass in longitudinal view; and a doughnut or target lesion in transverse view Point-of-care ultrasound may be useful for detecting intussusception at an early stage.[123] [124] Ultrasound can also be used to monitor success of treatment with enema, or if recurrence suspected after reduction.	»CT scan abdomen and pelvis: target lesion: intraluminal soft-tissue density mass with an eccentrically placed fatty area; reniform mass: high attenuation peripherally and lower attenuation centrally; sausage-shaped mass: alternating areas of low and high attenuation representing closely spaced bowel wall, mesenteric fat and/or intestinal fluid and gas CT scan should include intravenous and oral contrast for best yield

PIntussusception

History	Exam	1st Test	Other tests
		wdiagnostic#nema: meniscus sign; coiled spring sign Contraindicated if peritonitis, shock, perforation, or an unstable clinical condition is present. Should perforation ensue, the use of barium would lead to barium peritonitis; therefore, it would be problematic. Contrast enema (air or contrast reagent) is the long-standing most specific and sensitive diagnostic test. Contrast enema demonstrating ileocolic intussusception (black arrow) From the collection of Dr KuoJen Tsao; used with permission	and be performed only in cases in which peritonitis is not present. In general, for investigation of abdominal pain in children, CT scan of th abdomen and pelvis, with or without contras may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children. "FBC: may show raised WBC (suggests intestinal ischaemia) "faecal occult blood test: positive test can support suspicion of intussusception Negative test cannot reliably rule out the diagnosis.[115]

♦ Meckel's diverticulum

History	Exam	1st Test	Other tests
typically aged <2 years; may present with abdominal pain (may be intermittent or mimic acute appendicitis), and/or painless passage of bright red blood per rectum (haematochezia); often asymptomatic	painless dark red, maroon, or red-brick-coloured jelly-like stools; abdominal tenderness with guarding and rebound (may suggest diverticulitis); palpable abdominal mass (may suggest intussusception)	wabdominal ultrasound: tubular mass in longitudinal views and a doughnut or target appearance in transverse views suggests intussusception Good initial test if there is concern for intussusception. Point-of-care ultrasound may be useful for detecting intussusception at an early stage.[123] Sensitivity is user-dependent. *technetium-99m pertechnetate scan: positive Identifies ectopic gastric mucosa as tracer is taken up by mucin-secreting cells. A positive scan shows immediate tracer localisation in the stomach and in the right lower quadrant; an area of 1.8 cm² of ectopic gastric mucosa in a Meckel's diverticulum is required to produce a positive result. In the paediatric population, the specificity has been reported to be 95%, with an accuracy of 90%.[148]	»CT scan abdomen and pelvis: may show intussusception, Meckel's diverticulitis, and/or dilated bowel consistent with bowel obstruction Not routinely ordered unless concerned for alternative disease process or assessing for complications. May be considered if technetium-99m pertechnetate scan negative, but clinical suspicion remains.

♦ Mesenteric adenitis

History	Exam	1st Test	Other tests
diffuse abdominal pain; history of recent or current upper respiratory tract infection	fever, abdominal tenderness not localised to right lower quadrant, rhinorrhoea, hyperaemic pharynx or oropharynx (pharyngitis), and/ or associated extramesenteric lymphadenopathy (usually cervical)	 »abdominal ultrasound: enlarged mesenteric lymph nodes Can also show mural and mesenteric thickening suggestive of enteritis. Usually first-line test because it is inexpensive, reproducible, and carries no radiation risk. 	»CT scan abdomen and pelvis: enlarged mesenteric lymph nodes Demonstrates number and size of mesenteric lymph nodes. Can assess entire abdomen in a single scan. In general, for investigation of abdominal pain in children, CT scan of the abdomen and pelvis, with or without contrast, may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children.

♦ Hirschsprung's disease

History	Exam	1st Test	Other tests
males affected more commonly, mainly presents in early infancy (prior to 6 months); failure to pass meconium in first 36 hours of life strongly suggestive; increased incidence in Down syndrome	abdominal distension, fullness in left lower quadrant; palpable faecal mass on abdomen examination; absence of peritonitis (no guarding or rebound tenderness); small rectum and absence of stool on rectal examination (should be performed by clinician able to interpret features of Hirschsprung's	»abdominal x-ray: stool visible throughout colon, decreased air in rectum; air-fluid levels may be present Stool throughout the colon does not rule out other aetiologies of abdominal pain.	»rectal biopsy: required for a definitive diagnosis; absence of ganglion cells and the presence of an excess of non- myelinated nerves; presence of increased acetylcholinesterase The specimen must be taken at least 1.5 cm above the pectinate line. Suction biopsy has gained wide

Hirschsprung's disease **History** Exam disease); dysmorphic features of Down syndrome may be present

1st Test



Abdominal xray of a neonate
with abnormal
stooling pattern and
constipation. The
dilated transverse
and descending
colon is suggestive
of Hirschsprung's
disease

From the collection of Dr KuoJen Tsao; used with permission

It is very difficult to differentiate between a distended colon and the small bowel; it is a nonspecific investigation.

»contrast enema: proximal dilation with narrowing of the distal colon, with a funnel-shaped transition zone Considered a classical finding. However, in children less than 1 month of age, the findings may be difficult to interpret.

Other tests

acceptance, as it is easily performed, is associated with virtually no risk of perforation, and does not require an anaesthetic.

»anorectal
manometry: absent
reflex
Normally, when the
rectum is distended
with a balloon, pressure
in the anal canal falls
because of internal
sphincter relaxation.
This reflex is absent
in patients with
aganglionosis.

This is the most sensitive and specific test. However, the test has several limitations including the technical difficulty of assessing the newborn.

♦ Hirschsprung's disease

History	Exam	1st Test	Other tests
		Performed with water-soluble contrast material. No bowel preparation required. The infant is placed in a lateral position, and a rectal tube is introduced to barely above the anal canal.	

♦ Ulcerative colitis					
History	Exam	1st Test	Other tests		
positive family history, bloody diarrhoea, cramping abdominal pain, anorexia, weight loss, fever, rash	evidence of weight loss, pallor, abdominal tenderness, abdominal mass, iritis (inflamed irritated eyes), arthritis, sacroiliitis, erythema nodosum, pyoderma gangrenosum	»faecal calprotectin: raised Increasingly used to distinguish between inflammatory bowel disease and non- inflammatory bowel disease, with a sensitivity of up to 90% and a specificity of up to 80%.[116] [149] »FBC: leukocytosis, anaemia, thrombocytosis Iron deficiency anaemia may be present, especially in those with chronic disease. Leukocytosis may also be caused by certain medications used to treat inflammatory bowel disease, such as corticosteroids. »colonoscopy with biopsy: continuous uniform rectal	»plain abdominal x-rays: dilated loops with air-fluid level secondary to ileus; free air is consistent with perforation; in toxic megacolon, the transverse colon is dilated to 6 cm or more in diameter This test gives an approximate estimate of the extent of disease because an ulcerated colon usually contains no solid faeces. Easy, inexpensive, and widely available; ordered when initial presentation or subsequent relapses are associated with signs and symptoms of an acute abdomen: "CT scan abdomen: thickened inflamed bowel mucosa, thumbprinting, intestinal dilation or evidence of stricture; inflamed		

♦ Ulcerative colitis

History	Exam	1st Test	Other tests
		involvement, loss of vascular marking, diffuse erythema, mucosal granularity and friability, mucosal oedema ulcers, fistulas (rarely seen), normal terminal ileum (or mild backwash ileitis in pancolitis) Biopsy of mucosa should be performed in multiple segments of the colon. Endoscopic findings may be highly suggestive of inflammatory bowel disease but the characteristics of the mucosa (including ulcers) cannot be used to distinguish between ulcerative colitis and Crohn's disease. **erythrocyte** sedimentation rate: raised Inflammatory markers correlate closely with disease activity. **C-reactive protein: raised Inflammatory markers correlate closely with disease activity.	mesentery; intra- abdominal abscesses Ordered when complications or other diagnoses are being considered. In general, for investigation of abdominal pain in children, CT scan of the abdomen and pelvis, with or without contrast, may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children. "serologic markers: perinuclear antineutrophil cytoplasmic antibody (pANCA) and anti- Saccharomyces cerevisiae antibody (ASCA): positive pANCA Require special laboratories and are expensive but may be particularly useful for differentiating between Crohn's disease and ulcerative colitis in the paediatric population; about 70% of patients with ulcerative colitis have positive pANCA; about 70% of patients with

Ulcerative colitis

History	Exam	1st Test	Other tests
			Crohn's disease have positive ASCA.[122]

♦ Crohn's disease

History Exam 1st Test aphthous ulcers, »faecal calprotectin: crampy abdominal pain, intermittent diarrhoea, evidence of weight may be raised bloody diarrhoea loss, pallor, abdominal Increasingly used to if colitis a feature tenderness, abdominal distinguish between mass, perianal fistula, (blood less common inflammatory bowel perirectal abscess, anal in Crohn's disease disease and nonthan in ulcerative fissure, perianal skin colitis), weight loss, tags; extraintestinal inflammatory bowel manifestations fatigue, family history disease, with a of inflammatory bowel including iritis, arthritis, sensitivity of up to 90% disease sacroiliitis, erythema and a specificity of up »upper nodosum, pyoderma gangrenosum to 80%.[116] [149] »FBC: leukocytosis, anaemia. thrombocytosis Iron deficiency anaemia may be present, strictures especially in those with chronic disease. Leukocytosis may also be caused by certain medications used to treat inflammatory bowel disease, such as corticosteroids. »C-reactive protein: raised fistulas Inflammatory markers

Other tests

- »plain abdominal x-rays: small bowel or colonic dilation: calcification: intraabdominal abscesses Suggestive of the diagnosis of Crohn's disease and useful to assess severity.
- gastrointestinal series with small bowel followthrough: oedema and ulceration of the mucosa with luminal narrowing and Demonstrates features suggestive of Crohn's disease and aids in defining its distribution and severity.
- »CT scan abdomen and pelvis: skip lesions, bowel wall thickening, surrounding inflammation, abscess, Ordered when complications or other diagnoses are being considered. In general, for investigation of abdominal pain in children, CT scan of the abdomen and pelvis,

correlate closely with

sedimentation rate:

disease activity.

»erythrocyte

raised

◊ Crohn's disease

History	Exam	1st Test	Other tests
		Inflammatory markers correlate closely with disease activity. "colonoscopy with biopsy: may demonstrate inflammation, friability, ulcer formation, and oedema Biopsy of mucosa should be performed in multiple segments of the bowel. Endoscopic findings may be highly suggestive of inflammatory bowel disease but the characteristics of the mucosa (including ulcers) cannot be used to distinguish between ulcerative colitis and Crohn's disease. "MR enterography: skip lesions, bowel wall thickening, surrounding inflammation, abscess, fistulas	with or without contrast, may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children. "serological markers: perinuclear antineutrophil cytoplasmic antibody (pANCA) and anti-Saccharomyces cerevisiae antibody (ASCA): positive ASCA Require special laboratories and are expensive, but may be particularly useful for differentiating between Crohn's disease and ulcerative colitis in the paediatric population; about 70% of patients with ulcerative colitis have positive pANCA; about 70% of patients with Crohn's disease have positive ASCA.[122]

™Small bowel obstruction

History	Exam	1st Test	Other tests
intolerant of feeding, with nausea and/ or bilious vomiting; abdominal pain may or	limited abdominal distension (with proximal obstructions in the duodenum or early	»abdominal x-ray: dilated small bowel loops, air-fluid levels throughout abdomen	»abdominal ultrasound: may demonstrate focal area causing obstruction

Small bowel obstruction

History

may not be a feature; history of previous abdominal surgery; history of cystic fibrosis may be present

Exam

jejunum); abdominal tenderness may or may not be present; rebound tenderness and guarding may occur if perforation, ischaemia, and peritonitis; hyperactive bowel sounds (early finding), hypoactive or absent bowel sounds (late finding); incarcerated femoral, obturator, umbilical or ventral hernia may be present

1st Test

Supine and upright films may reveal characteristic pattern of gas in bowel. In duodenal atresia may see double bubble sign.



Abdominal x-ray
demonstrating
double bubble gas
pattern consistent
with duodenal atresia
From the collection
of Dr KuoJen Tsao;
used with permission

Jejunal atresia shows dilated proximal small bowel with decompressed non-airfilled bowel distally.

Other tests

Can be used to diagnose intussusception; may be of particular value when diagnosing intussusception secondary to Henoch-Schonlein purpura, as it is typically ileo-ileal instead of ileocolic.[22] Point-of-care ultrasound may be useful for detecting intussusception at an early stage.[123]

Ultrasound with
Doppler blood flow
assessment may
assist in the diagnosis
of mid-gut volvulus
associated with
malrotation. In addition,
it may be useful in the
diagnosis of omental
and duplication cysts.

"upper gastrointestinal contrast study: dilated small intestine; may demonstrate a transition zone of obstruction Critical to rule out acute mid-gut volvulus and/or malrotation; water-soluble contrast challenge can be both diagnostic and therapeutic.[150]

™Small bowel obstruction

History	Exam	1st Test	Other tests
			Upper GI contrast
			study demonstrating
			malrotation with
			volvulus. The
			duodenum fails
			to develop the
			normal anatomical
			C-loop. There is
			failure of contrast
			to pass, resulting
			in a characteristic
			bird beak consistent
			with acute mid-
			gut volvulus
			From the collection
			of Dr KuoJen Tsao;
			used with permission
			»lower gastrointestinal contrast study: dilated small intestine; may demonstrate a transition zone of obstruction May be helpful in diagnosing meconium disease; hypertonic contrast medium may be helpful for clearing meconium.

PSmall bowel obstruction

History	Exam	1st Test	Other tests
			"CT scan abdomen: dilated small intestine; may demonstrate a transition zone of obstruction, mass, tumour, abscess Helps to determine diagnosis of underlying cause, extent, and location of the obstruction when x-rays are inconclusive.[151] CT scan should include intravenous and oral contrast for best yield and be performed only in cases in which peritonitis is not present.

PVolvulus

History	Exam	1st Test	Other tests
infant age group; history of bilious vomiting; pain usually manifests as notable transition to an inconsolable state	often diffuse abdominal distension and tenderness; faint or no bowel sounds, rigid abdomen, guarding, rebound tenderness, fever, or haematochezia	wupper gastrointestinal contrast study: bird beak sign of stricture at the site of the volvulus Standard diagnostic test for malrotation. Should be performed with anteroposterior and lateral projections if there is any history of bilious vomiting without previous abdominal surgery. Not necessary if diagnosis is strongly indicated from plain x-rays or there is	»CT scan abdomen: bowel obstruction with whirl pattern of mesentery Usually not necessary to make diagnosis.

PVolvulus

History	Exam	1st Test	Other tests
		evidence of bowel	
		necrosis.	
		and the same of th	
		\$100 PM	
		The case of the case of	
		The second second	
		13.00	
		Upper GI contrast	1
		study demonstrating	
		malrotation with	
		volvulus. The	
		duodenum fails	
		to develop the	
		normal anatomical	
		C-loop. There is	
		failure of contrast	
		to pass, resulting	
		in a characteristic	
		bird beak consistent	
		with acute mid-	
		gut volvulus From the collection	
		of Dr KuoJen Tsao;	
		used with permission	
		adda war pormiodion	
		»abdominal x-ray:	
		partial or complete obstruction; dilated	
		bowel loops; air-fluid	
		levels; abdominal free	
		air with perforation	
		Images in both flat	
		(supine) and upright	
		(erect) position are	
		taken.	

™Volvulus

History

History	Exam	1st Test	Other tests
		»FBC: raised WBC (suggests intestinal ischaemia)	

Large bowel obstruction

Exam

history of risk factors: neurodevelopmental problems, inflammatory bowel disease, diabetes, poor diet, previous colorectal resection, laxative misuse, megacolon, or previous abdominal surgery; change in bowel habit with partial or complete obstruction, or change in calibre of stool; colicky abdominal pain becoming more constant and worse with movement, coughing or deep breathing as bowel approaches perforation; intolerant of feeding, with nausea or vomiting

tympanic, distended abdomen; hyperactive bowel sounds that become absent in advanced stages; abdominal rebound, quarding, and/or rigidity if perforation or close to perforation; empty rectum; incarcerated femoral, obturator, umbilical, or ventral hernia may be present

»abdominal x-ray: gaseous distension of large bowel; volvulus suggested by kidneybean-shape bowel loop Diagnosis confirmed by colonic dilation. Level of obstruction may be determined by a cut-off beyond which the colon or rectum is empty of

gas.

1st Test

Intramural gas (pneumatosis) may suggest colonic ischaemia.

Other tests

- »abdominal ultrasound: may demonstrate focal area causing obstruction (e.g., intussusception) Point-of-care ultrasound may be useful for detecting intussusception at an early stage.[123]
- »lower gastrointestinal (GI) contrast study: may indicate site of obstruction Used to diagnose meconium disease or other cause of lower GI intestinal obstruction such as intussusception.

Hypertonic contrast medium may be helpful for clearing meconium.

»CT scan abdomen and pelvis: gaseous distension of large bowel; may demonstrate a transition zone of obstruction May also reveal underlying cause and provide more

PLarge bowel obstruction

History	Exam	1st Test	Other tests
			diagnostic information
			than contrast enema.
			Intravenous and oral
			contrast should be
			used for best diagnostic
			yield. Radiation
			exposure should be
			considered.[133]
			Sedation or general
			anaesthesia may
			be required in some
			children.
			»flexible/rigid sigmoidoscopy: flood of stool and mucus upon passing and decompressing apex of volvulus Useful if sigmoid volvulus is suspected, as sigmoidoscopy may be potentially therapeutic. Must exercise extreme caution if ischaemia is suspected, as there is an increased chance of perforation.

™Necrotising enterocolitis

History	Exam	1st Test	Other tests
premature neonate weighing less than 1500 g; feeding intolerance, apnoea, lethargy, bloody stools	abdominal distension, tenderness, abdominal wall erythema, haematochezia, bradycardia	»FBC: leukocytosis or leukopenia, anaemia, thrombocytopenia Decreased WBC count may be more ominous than leukocytosis. Thrombocytopenia develops in 50% to	 »abdominal ultrasound: fluid collections, ascites Useful for diagnosing fluid collections that may represent perforation or abscess. A few centres use

™ Necrotising enterocolitis

History	Exam	1st Test	Other tests
		95% of all infants with necrotising enterocolitis within 24-72 hours of receiving the diagnosis.[152] Anaemia may result with significant haematochezia. **blood culture:* negative Bacterial sepsis should be excluded. **serum electrolyte panel:* hyponatraemia **abdominal x-*	abdominal ultrasound for early diagnosis and serial follow-up; very sensitive but still not widely used.[153]
		ray: dilated loops of bowel, pneumatosis intestinalis, portal venous gas, free air, fixed loop of bowel, lack of normal intestinal gas pattern Important to obtain both anteroposterior and left lateral decubitus views.	

Peptic ulcer disease

History	Exam	1st Test	Other tests
non-steroidal anti- inflammatory drug use; family history of peptic ulcer disease; weight loss, vomiting, anorexia, and intermittent epigastric pain, usually related to eating meals; pain often nocturnal and usually relieved by antacids; melaena and/or haematemesis	unremarkable or epigastric tenderness, melaena, or occult bleeding on stool haemoccult test	»FBC: normal or leukocytosis; anaemia present if sustained blood loss »erect chest x-ray: usually normal Ordered to rule out gastric or duodenal perforation (demonstrated by free air under the diaphragm).	»Helicobacter pylori breath test or stool antigen test: positive result if Helicobacter pylori present Generally not necessary if endoscopy and biopsy are being performed; however, in places where endoscopy is not readily available, and

Peptic ulcer disease

History	Exam	1st Test	Other tests
if erosion into blood vessel		pastrointestinal series with water-soluble contrast: mucosal defect(s) consistent with ulcer or free intraperitoneal contrast consistent with perforation Gastric and duodenal ulcers can sometimes be visualised with x-ray imaging. Contrasted imaging studies can indicate location and depth of ulcers. This may serve as an initial test prior to the more definitive endoscopy: "upper gastrointestinal endoscopy: mucosal inflammation, ulceration, and haemorrhage Endoscopy should be performed urgently if bleeding is suspected (prior to any contrast radiological studies). Contrast may interfere with the assessment of the mucosa. Biopsies and testing for Helicobacter pylori may be performed.	peptic ulcer disease is suspected, <i>H pylori</i> breath test or stool antigen test may be helpful, and may also be performed 4 weeks after starting treatment, to assess treatment success.[119] [120] A blood test for <i>H pylori</i> is not useful in children.

♦ Coeliac disease

History	Exam	1st Test	Other tests
recurrent abdominal pain, cramping, or distension; bloating and	generalised abdominal pain or bloating; underweight or failing	»immunoglobulin A-tissue transglutaminase	<pre>»endomysial antibody (EMA): raised titre</pre>

♦ Coeliac disease

History diarrhoea: dermatitis herpetiformis, an intensely pruritic papulovesicular rash that affects the extensor limb surfaces, almost universally occurs in association with coeliac disease; may be a history of immunoglobulin A deficiency, type 1 diabetes, autoimmune thyroid disease, Down syndrome, Sjogren syndrome, inflammatory bowel disease, or primary biliary cholangitis; may be a family history of coeliac disease

Exam

to thrive; aphthous stomatitis; dermatitis herpetiformis

1st Test

(IgA-tTG): titre above normal range for laboratory IgA anti-tissue transglutaminase antibody is the preferred single test for detection of coeliac disease in individuals over the age of 2 years.[154] When screening children aged <2 years for coeliac disease, the IgA-tTG test should be combined with deamidated gliadin peptide (IgA and IgG).[154]

»quantitative immunoglobulin A (IgA): titre normal or below normal range for laboratory In patients with low total IgA concentrations, an IgG-based test (e.g., deamidated gliadin peptide) should be

»FBC: may show iron deficiency anaemia

performed.[155]

»endoscopy and small bowel biopsy: presence of intraepithelial lymphocytes, villous atrophy, and crypt hyperplasia Small-bowel histology is essential and the gold-standard test to confirm the diagnosis.

Other tests

EMA is a more expensive alternative to IgA-tTG with greater specificity but lower sensitivity.

Perform initially if IgAtTG is unavailable.

»human leukocyte antigen (HLA) typing: positive HLA-DQ2 or HLA-DQ8
This genetic test is useful to rule out coeliac disease in patients already on a gluten-free diet or in patients with an idiopathic coeliac-like enteropathy.[46]

Paediatric patients with symptoms consistent with coeliac disease and a high IgA-tTG titre (≥10 times the upper limit of normal) may go on to have confirmatory EMA and HLA-DQ2/-DQ8 testing. If both of these are positive, coeliac disease may be considered confirmed without a small intestinal biopsy.[155] Children with lower IgA-tTG titres (<10 times upper limit of normal) should undergo biopsies to decrease

♦ Coeliac disease

History	Exam	1st Test	Other tests
		Two biopsies of the	the risk of false positive
		bulb and at least four	diagnosis.[155]
		biopsies of the distal	
		duodenum should	
		be submitted for	
		histological analysis.	
		Paediatric patients with	
		symptoms consistent	
		with coeliac disease	
		and a high IgA-tTG titre	
		(≥10 times the upper	
		limit of normal) may go	
		on to have confirmatory	
		EMA and HLA-DQ2/-	
		DQ8 testing. If both	
		of these are positive,	
		coeliac disease may be	
		considered confirmed	
		without a small	
		intestinal biopsy.[155]	
		Children with lower	
		IgA-tTG titres (<10	
		times upper limit of	
		normal) should undergo	
		biopsies to decrease	
		the risk of false positive	
		diagnosis.[155]	

™Viral hepatitis

History	Exam	1st Test	Other tests
birth or residence in endemic area, antenatal exposure, family history of chronic viral hepatitis, multiple sexual partners, sexual intercourse with infected individuals (hepatitis B and/or C), travel to developing	jaundice; early disease: tender hepatosplenomegaly, lymphadenopathy; late disease: generalised wasting, cachexia, gynaecomastia, ascites, altered sensorium, asterixis, or decreased deep	»serum LFTs: high direct bilirubin, AST, ALT, alk phos and gamma-GT Although transaminases may be raised, the level is not a good correlate for	 »FBC: low or normal platelet count A low platelet count is suggestive of portal hypertension. »coagulation profile (prothrombin time [PT], INR): May be raised or normal

History

countries, pregnant (hepatitis E); early disease: malaise, muscle and joint aches, fever, nausea, vomiting, diarrhoea, headache, anorexia, dark urine, pale stool, abdominal pain; late disease: weight loss, easy bruising and bleeding tendencies

Exam

tendon reflexes, caput medusa, ascites, hepatosplenomegaly, congestion secondary to right heart failure

1st Test

severity of infection; it indicates liver damage.

- »serum IgM anti-HAV: positive if acute hepatitis A infection
- »serum hepatitis
 B surface antigen
 (HBsAg): positive if
 hepatitis B infection
 Positive result appears
 in serum 1 to 10 weeks
 after an acute exposure
 and disappears when
 infection abates.
- »serum hepatitis
 B core antigen
 (HBcAg): positive if
 hepatitis B infection
 An intracellular antigen
 that is found in infected
 hepatocytes.
- »serum hepatitis B e antigen (HBeAg): positive if hepatitis B infection Patients with a positive result are considered highly infective for hepatitis B.
- »serum HCV RNA:
 positive if hepatitis C
 infection
 Identifies acute or
 chronic infection with
 hepatitis C virus.
 Progression to chronic
 disease after acute
 infection is highly
 likely. Significant risk
 for hepatocellular
 carcinoma.

Other tests

In cases of severe hepatitis, liver synthetic function as measured by coagulation parameters may be compromised, reflected by increased PT and INR.

History	Exam	1st Test	Other tests
		Patients should be at	
		least 1 year old prior to	
		testing for hepatitis C.	
		»serum total (IgM and IgG) anti-HDV antibodies: positive if hepatitis D infection	
		Performed by enzyme-	
		linked or radio-	
		immunoassay. Hepatitis	
		D is a defective virus	
		and can only replicate	
		in the presence	
		of co-infection or	
		superinfection with	
		HBV.	
		»serum anti-HEV IgM antibodies: positive if acute hepatitis E infection	

♦ Biliary dyskinesia

History	Exam	1st Test	Other tests
history of previous negative work-up for cholelithiasis common; recurrent right upper quadrant (RUQ) pain; nausea and vomiting; symptoms may or may not be associated with eating	may be equivocal; RUQ tenderness	»LFTs: normal aspartate aminotransferase, alanine aminotransferase, alk phos, and bilirubin »RUQ ultrasound: normal Gallstones should not be demonstrated in patients with biliary dyskinesia.	»hepatobiliary iminodiacetic acid (HIDA) scan: decreased (<35%) gallbladder ejection fraction Considered the diagnostic test of choice. The ejection fraction (EF) of the gallbladder is measured after the administration of cholecystokinin (CCK); definition of dyskinesia widened at some centres to include

♦ Biliary dyskinesia

History	Exam	1st Test	Other tests
			high EFs (85% to 90% +) and even normal results. No consistent predictive factors for subset of children that will have mediumto long-term relief of symptoms following cholecystectomy.

			cholecystectomy.		
P Acute pancreatitis					
History	Exam	1st Test	Other tests		
nausea, vomiting, epigastric pain radiating to back; acute-onset abdominal pain	epigastric or upper abdominal tenderness; tachycardia and hypotension in severe cases; discoloration around the umbilicus (positive Cullen's sign) or flanks (positive Grey-Turner's sign) in cases of haemorrhagic pancreatitis; small children may demonstrate increased irritability and abdominal distension only	"lipase: at least 3 times upper limit of normal range; can be raised if amylase normal Use serum lipase in preference to serum amylase, if available.[114] Serum lipase levels remain raised for longer (up to 14 days after symptom onset vs. 5 days for amylase), providing a higher likelihood of picking up the diagnosis in patients with a delayed presentation.[156] "amylase: at least 3 times upper limit of normal range Use serum lipase in preference to serum amylase, if available.[114] Serum lipase levels remain raised for longer	"abdominal ultrasound: may appear normal early in disease course; enlargement of the pancreatic oedema; dilated pancreatic duct; may show underlying biliary disease Ultrasound is a useful and commonly employed modality for the diagnosis of pancreatic disease in children, as there is no radiation exposure involved. Permits concurrent assessment of the biliary system for choledithiasis and/or choledocholithiasis that may be the underlying cause of pancreatitis. "CT scan abdomen with intravenous contrast:		

PAcute pancreatitis

History	Exam	1st Test	Other tests
		(up to 14 days after symptom onset vs. 5 days for amylase), providing a higher likelihood of picking up the diagnosis in patients with a delayed presentation.[156] *bilirubin: normal or raised An raised bilirubin (specifically direct bilirubin) is suggestive of gallstone-induced pancreatitis. In addition, choledochal cysts are an important cause of pancreatitis in children.	peripancreatic inflammation (fat stranding); may show gallstones Although this involves exposure to radiation, it is an excellent tool in the diagnosis of early pancreatitis. Ordered if patient is not improving with standard treatment or if complicated disease is suspected. Should not be ordered for uncomplicated cases of pancreatitis in which the patient is improving. Allows better assessment of potential pancreatic necrosis (which would indicate more serious disease) and peripancreatic inflammation than is achievable with ultrasound.

PAcute pancreatitis

History	Exam	1st Test	Other tests
			CT scan of teenage
			girl presenting
			with mid-epigastric
			abdominal pain as
			a result of gallstone
			pancreatitis. The
			large fluid collection
			in the pancreatic bed
			(white arrow) and
			lack of pancreatic
			enhancement
			suggest liquefactive
			necrosis of
			the pancreas
			From the collection
			of Dr Kuojen Tsao;
			used with permission

PSplenic infarction/cysts

History	Exam	1st Test	Other tests
varied; may be history of trauma; cysts either asymptomatic or dull, left-sided abdominal pain; infarction typically causes fever as well as pain, but occasionally asymptomatic; left-sided shoulder and/or chest pain; presence of risk factors for splenic infarction (sickle cell disease, high altitude)	may be vague left upper quadrant tenderness	»Doppler ultrasound: infarction or cyst on spleen Splenic infarcts appear as wedge-shaped defects. Blood flow is absent, with absence of Doppler signal. Cysts appear anechoic.	»CT scan abdomen with intravenous contrast: infarction or cyst on spleen Splenic infarcts appear as wedge-shaped defects. Blood flow is absent, manifested as lack of contrast enhancement on CT scan.

PSplenic infarction/cysts

History	Exam	1st Test	Other tests
			Splenic cysts may also be visualised.
			MAPAY 197 Sci Landard Languages
			CT scan
			demonstrating
			fluid-filled cyst
			within the spleen
			From the collection
			of Dr KuoJen Tsao;
			used with permission

♦ Nephrolithiasis

History	Exam	1st Test	Other tests
family history of nephrolithiasis and/ or gout; intermittent, severe, colicky flank and/or abdominal pain; nausea and vomiting; gross or microscopic haematuria; urinary frequency/urgency; atypical presentation common in younger children	ipsilateral costovertebral angle and flank tenderness; tachycardia and hypotension in pain- controlled patient may suggest concurrent urosepsis	»ultrasound of the urinary tract: calcification seen within urinary tract; possible dilated proximal ureter and hydronephrosis Ultrasound should be the first modality. As well as diagnosing nephrolithiasis, it can provide information on potential complications of obstruction such as hydronephrosis. »urinalysis: may be normal or dipstick- positive for leukocytes, nitrites, blood; microscopic analysis positive for WBCs, red blood cells, or bacteria	»abdominal x-ray: radiopaque stones If sufficiently radiopaque, urinary stones may be visible on plain x-rays. Calcium oxalate and calcium phosphate stones are radiopaque, whereas pure uric acid and indinavir stones are radiolucent and cystine stones are partially radiolucent. »non-contrast CT scan abdomen and pelvis: calcification seen in renal collecting system or ureter; possible dilated proximal ureter and hydronephrosis

♦ Nephrolithiasis

History	Exam	1st Test	Other tests
		Patients with nephrolithiasis typically have blood in the urine. It is important to pay attention to other elements of the urine analysis, as the presence of white blood cells, leukocyte esterase, and nitrites may signify a concurrent urinary tract infection.	Highly sensitive for detection of nephrolithiasis and assessment of hydronephrosis. In addition, if stones are not seen, an alternative aetiology may be detected; frequently, nephrolithiasis detected on CT scan is a serendipitous finding for the investigation of other aetiologies of abdominal pain.

™Sepsis

History	Exam	1st Test	Other tests
may be history of decreased activity, carer concern that child is not behaving as normal; may not wake or may be difficult to rouse	fever may be present (although core temperature may also be normal or low), tachycardia, raised respiratory rate, may be delayed capillary refill time, oxygen saturations may be <92% in air, altered mental state, petechiae or purpura may be present	»FBC: WBC variable; may be thrombocytopenia Abnormal WBC count (i.e., above or below normal range for age or >10% immature white cells) is one of the core diagnostic criteria for systemic inflammatory response syndrome.[157] However, raised or depressed WBC count is not specific for the diagnosis of sepsis. »blood glucose: may be hypo- or hyperglycaemia »serum electrolytes: deranged	»Iumbar puncture: positive culture may reveal bacterial meningitis as infective source »procalcitonin: raised Raised procalcitonin levels may be better than traditional laboratory screening tests for detecting serious bacterial infections in febrile infants and young children without obvious sources of infection.[158] [159] [160] »CT scan of the chest and/or abdomen and pelvis:

™Sepsis

History	Exam	1st Test	Other tests
		»blood lactate: raised May be possible to measure lactate using the arterial blood gas sample. »arterial blood gas: hypoxaemia and/or hypercarbia; large base deficit »coagulation studies: may be abnormal »LFTs: may be abnormal »blood culture: may be growth of bacteria identifying pathogen »chest x-ray: may reveal evidence of pneumonia »urinalysis: may be positive for nitrites and leukocytes if urinary tract infection present »urine culture: may be positive if urinary tract infection present	may reveal focus of infection CT may play an important role in the evaluation of patients with suspected sepsis. The increased exposure to radiation with this investigation should be considered. CT scan can help to identify a hidden collection (e.g., an intra-peritoneal abscess or effusion) which may not be readily apparent on ultrasound or chest x-ray. CT can also be used to identify free air (perforation). "abdominal ultrasound scan: may reveal focus of infection Useful if an abdominal source of infection is suspected, or where the source of infection is not clear after the initial clinical examination and tests. Ultrasound has a reasonable falsenegative rate; absence of positive findings on ultrasound does not rule out any given infection source.

Testicular torsion

History 1st Test Other tests Exam tender, oedematous acute-onset testicular »Testicular Workup »urinalysis: usually pain: nausea, and testicle: affected testicle for Ischemia and normal, but can be **Suspected Torsion** abnormal in some vomiting; history of may appear higher recurrent episodes than unaffected testicle (TWIST) score: low cases suggests repeated with horizontal lie: risk (score 0 to 2), If abnormal, usually episodes of testicular associated scrotal intermediate risk (3 to suggests alternate torsion followed by erythema and oedema; 4), or high risk (5 to 7) diagnosis (e.g., spontaneous detorsion; absent cremasteric Sensitivity 98.4% in epididymitis or orchitis). history of trauma may reflex; usually no pain low-risk patients and be present relief with elevation of However, it is important specificity 97.5% in the scrotum; thin skin of to note that the high-risk patients.[161] the scrotum sometimes urinalysis may be allows visualisation of »duplex Doppler negative in cases of the torsed appendage ultrasound of ('blue dot or black dot epididymitis or orchitis scrotum: presence sign') and positive in the of fluid and the setting of testicular whirlpool sign (the swirling appearance torsion. of the spermatic cord from torsion as the ultrasound probe scans downwards perpendicular to the spermatic cord); absent or decreased blood flow in the affected testicle: decreased flow velocity in the intratesticular arteries, increased resistive indices in the intratesticular arteries Definitive test for torsion and should be obtained unless the test would unnecessarily delay intervention in a child for whom there is a high index of suspicion. Diagnostic studies should not

preclude operative intervention.[94] [95]

Duplex Doppler ultrasound includes grey scale, colour

Testicular torsion

History	Exam	1st Test Other tests
		Doppler, and spectral
		Doppler ultrasound.
		Power Doppler can
		be used as well as
		or instead of colour
		Doppler. Power Doppler
		is more sensitive for
		evaluating areas of low
		blood flow such as the
		prepubertal testis.[162]

POvarian torsion

History	Exam	1st Test	Other tests
acute onset of one- sided lower abdominal or pelvic pain; nausea and vomiting common; history of frequent, similar episodes; fever rare	tender pelvic mass (adnexal); in patients old enough to undergo pelvic examination, cervical motion tenderness may be elicited; typically no vaginal discharge, but may be some mild to moderate vaginal bleeding	»pelvic ultrasound: solid appearance of the ovary, unilateral ovarian enlargement, ovarian peripheral cystic structures, marked stromal oedema, fluid in the pouch of Douglas	"colour Doppler ultrasound: reduced or absent intraovarian blood flow Suggestive of ovarian torsion. Can help determine whether the flow in the ovary is impaired. Doppler ultrasound is of only variable reliability in the diagnosis of ovarian torsion.[87] [88] "CT scan abdomen and pelvis: enlarged, oedematous ovary with or without vascular enhancement; free fluid in pelvis CT scans can demonstrate adnexal masses and ischaemia of the ovary. Typically, ovarian torsion diagnosed by CT scan is a serendipitous finding during an

POvarian torsion

History	Exam	1st Test	Other tests
			attempt to identify other sources of abdominal pain. CT scan of a young girl presenting with ovarian torsion. The large pelvic cystic lesion contains calcifications (white arrow) consistent with a teratoma or dermoid cyst From the collection
			of Dr KuoJen Tsao; used with permission In general, for investigation of abdominal pain in children, CT scan of the abdomen and pelvis, with or without contrast, may provide a high yield of information, but radiation exposure should be considered.[133] Sedation or general anaesthesia may be required in some children.

♦ Ruptured ovarian cyst

History	Exam	1st Test	Other tests
rupture usually spontaneous, can follow history of trauma or sexual intercourse; mild chronic lower abdominal discomfort may suddenly intensify; may occur in conjunction with torsion	adnexal tenderness; adnexal size unremarkable due to collapsed cyst; peritonism may be present in lower abdomen and pelvis	»pelvic ultrasound: complex mass appearance; fluid in the pouch of Douglas A complex mass appearance suggests a ruptured haemorrhagic cyst surrounded by a haematoma. Fluid in the pouch of Douglas is also indicative of a ruptured cyst.	

♦ Pelvic inflammatory disease (PID)

History	Exam	1st Test	Other tests
sexually active; multiple partners; history may be suggestive of sexual abuse (particularly if young child); pain worse with sexual intercourse; dull, aching lower abdominal pain with or without dysuria; vaginal discharge, lowgrade fever	temperature >38.3°C (101°F); cervical motion tenderness, adnexal or uterine tenderness, vaginal or cervical mucopurulent discharge	»wet mount of vaginal secretions: polymorphonuclear leukocytes (PMNs) seen Presence of vaginal PMNs confirms vaginal infection. High predictive value but not specific for pathogens most likely to cause PID. »nucleic acid amplification test or culture of vaginal secretions for Neisseria gonorrhoeae and Chlamydia trachomatis: positive result indicates presence of organism Supports a diagnosis of PID. Specimens obtained from vaginal	»pelvic ultrasound: normal or may demonstrate endometritis, hydrosalpinx, pyosalpinx, tubo- ovarian abscess Test is useful in confirming an uncertain diagnosis. Ultrasound findings can be variable demonstrating any portion of the female reproductive organs. Most common findings include pyosalpinx and tubo-ovarian abscess. »HIV serology: positive or negative It is important to test for other sexually transmitted diseases during a work-up for PID, including HIV infection.

♦ Pelvic inflammatory disease (PID)

History	Exam	1st Test	Other tests
		examination can be sent for testing. Urine tests should not be the initial test ordered but may be necessary if a pelvic examination is not feasible (e.g., due to age of child).	»hepatitis studies: positive or negative Testing for hepatitis B and C may be considered during work- up for PID. »rapid plasma reagin (RPR): positive or negative It is important to test for other sexually transmitted diseases during a work-up for PID, including syphilis. »FBC: leukocytosis Result is not specific but may increase index of suspicion for PID. »C-reactive protein or erythrocyte sedimentation rate: raised Non-specific test for inflammatory process. »laparoscopy: normal or may demonstrate endometritis, hydrosalpinx, pyosalpinx, tubo- ovarian abscess Laparoscopy is the preferred invasive method of diagnosis allowing direct visualisation of the gynaecological and abdominal structures. In addition, it may aid the diagnosis of Fitz- Hugh-Curtis syndrome.

Pregnancy complications

Pregnancy col	inplications		
History	Exam	1st Test	Other tests
history of previous ectopic pregnancy or miscarriage, fallopian tube or pelvic surgery, pelvic inflammatory disease; lower abdominal pain, amenorrhoea, and vaginal bleeding	minimal abdominal tenderness and/or vaginal bleeding; pelvic examination may reveal a mass, eliciting cervical motion tenderness if haemoperitoneum is present; tubal rupture can cause haemodynamic instability	wurine pregnancy test: positive Confirms pregnancy; although false-negative tests, depending on kit used, are known. "quantitative serum beta-hCG: positive A normal intrauterine pregnancy is typically visualised at serum human chorionic gonadotrophin (beta-hCG) levels between 1500 and 2500 IU/L (1500 and 2500 mIU/mL). If not, the likely differential diagnosis is ectopic pregnancy or miscarriage. Serial beta-hCG measurement is usually used when the ultrasound is inconclusive and the initial beta-hCG is below the discriminatory zone (1500-2000 IU/L [1500-2000 mIU/mL]). An increase of at least 66% in 2 measurements over 48 hours is suggestive of a viable intrauterine pregnancy. An increase of <66% is suggestive of an ectopic pregnancy. A rapid and steady	»blood type and screen: variable Necessary to assess the rhesus status of the mother if not yet known. If Rh-negative, should receive anti-D immunoglobulin to prevent rhesus isoimmunisation, which may affect subsequent pregnancy outcomes.

Pregnancy complications

Pregnancy complications

History	Exam	1st Test	Other tests
		for an ultrasound. A quick and focused bedside ultrasound examination to assess for the presence of free fluid or blood may be helpful when ruptured ectopic pregnancy is suspected. **transvaginal ultrasound: presence or absence of intrauterine pregnancy Differentiates between different stages and types of miscarriage. Specific, fairly prognostic and would exclude other differential diagnoses of miscarriage.	
		Occasionally directly identifies ectopic pregnancy.	
		Usually used in conjunction with pelvic ultrasound in the setting of positive beta-hCG plus abdominal pain and/or vaginal bleeding.	

PEmpyema

History	Exam	1st Test	Other tests
recent pneumonia, fever, cough, chest pain; malaise, anorexia, weight loss, or fatigue may occur; presence of risk factors	febrile, toxic patient, dullness on percussion, absence of breath sounds over affected area; abdominal tenderness and	»FBC: raised WBC count»chest x-ray: blunting of costophrenic angle or effusion on	»blood culture: positive for specific pathogens Blood cultures may be positive even if the

Empyema

History Exam 1st Test Other tests (immunocompromise, distension without affected side, possible pleural fluid culture is comorbidities guarding or rebound consolidation, pleurally negative. predisposing to the based 'D' shape in development of empyema Should be taken pneumonia, pre-The presence of a before the initiation of existing lung disease, loculated effusion iatrogenic interventions antibiotics if the clinical suggests an empyema. in the pleural space, state of the patient There may be male sex) permits. associated pulmonary consolidation due to »chest ultrasound: localised fluid collection pneumonia. May consider before »thoracentesis: proceeding to CT frank pus in empyema, scan due to the lack of serous or cloudy radiation exposure. in complicated parapneumonic »CT scan chest effusions with intravenous Aspiration of frank contrast: consolidation pus is diagnostic of of lung parenchyma; empyema, and no extraparenchymal fluid with loculations other investigations are suggests empyema required to establish CT scan may better the diagnosis. differentiate between empyema and simple effusion.

Guidelines

Europe

Rome IV: functional gastrointestinal disorders - disorders of gut-brain interaction. 4th ed. (https://theromefoundation.org/rome-iv)

Published by: Rome Foundation

Last published: 2016

International

Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines (https://www.wses.org.uk/guidelines)

Published by: World Society of Emergency Surgery

Last published: 2020

The management of intra-abdominal infections from a global perspective (https://www.wses.org.uk/guidelines)

Published by: World Society of Emergency Surgery

Last published: 2017

North America

SAGES guideline for the diagnosis and treatment of appendicitis (https://www.sages.org/publications/guidelines)

Published by: Society of American Gastrointestinal and Endoscopic Surgeons

Last published: 2024

Clinical policy: critical issues in the evaluation and management of emergency department patients with suspected appendicitis (https://www.acep.org/patient-care/clinical-policies)

Published by: American College of Emergency Physicians

Last published: 2023

ACR appropriateness criteria: acute nonlocalized abdominal pain (https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria)

Published by: American College of Radiology

Last published: 2018

North America

ACR appropriateness criteria: suspected appendicitis - child (https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria)

Published by: American College of Radiology

Last published: 2018

ACR appropriateness criteria: fever without source or unknown origin - child (https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria)

Published by: American College of Radiology

Last published: 2015

Evidence tables

What are the effects of early versus late initiation of empiric antimicrobial treatment in children with or at risk of developing sepsis or severe sepsis?



This table is a summary of the analysis reported in a guideline (underpinned by a systematic review) that focuses on the above important clinical question.

View the full source guideline (https://www.nice.org.uk/guidance/ng51/evidence)

Evidence C *

Confidence in the evidence is very low or low where GRADE has been performed and the intervention may be more effective/beneficial than the comparison for key outcomes. However, this is uncertain and new evidence could change this in the future.

Population: Children with or at risk of developing sepsis or severe sepsis

Intervention: Early initiation of empiric antimicrobial treatment **Comparison:** Late initiation of empiric antimicrobial treatment

Outcome	Effectiveness (BMJ rating)	Confidence in evidence (GRADE) [‡]
<1 hour versus >1 hour a	'	'
Paediatric Intensive Care Unit (PICU) mortality	No statistically significant difference	Very Low
<2 hours versus >2 hours a	'	'
PICU mortality	No statistically significant difference	Very Low
<3 hours versus >3 hours a		1
PICU mortality	Favours intervention	Very Low
<4 hours versus >4 hours a	1	1
PICU mortality	Favours intervention	Very Low

Recommendations as stated in the source guideline

For children aged 5–11 years who have suspected sepsis and one or more high-risk criteria, give a broad-spectrum antimicrobial ^b at the maximum recommended dose without delay (within one hour of identifying that they meet any high-risk criteria in an acute hospital setting).

Note

The guideline group noted that the direct evidence in children came from one small (n=130), single-centre retrospective study of children in PICU with severe sepsis and septic shock. Therefore, they also

extrapolated from the indirect evidence in adults to make the same recommendation for all age groups (including children aged under 5 years and 5-11 years).

- ^a Time from sepsis recognition to initial treatment and first appropriate treatment.
- ^b See full guideline for more information.

* Evidence levels

The Evidence level is an internal rating applied by BMJ Best Practice. See the EBM Toolkit (https://bestpractice.bmj.com/info/evidence-tables/) for details.

Confidence in evidence

- A High or moderate to high
- **B** Moderate or low to moderate
- C Very low or low

† Effectiveness (BMJ rating)

Based on statistical significance, which demonstrates that the results are unlikely to be due to chance, but which does not necessarily translate to a clinical significance.

‡ Grade certainty ratings

High	The authors are very confident that the true effect is similar to the estimated effect.
Moderate	The authors are moderately confident that the true effect is likely to be close to the estimated effect.
Low	The authors have limited confidence in the effect estimate and the true effect may be substantially different.
Very Low	The authors have very little confidence in the effect estimate and the true effect is likely to be substantially different.

BMJ Best Practice EBM Toolkit: What is GRADE? (https://bestpractice.bmj.com/info/toolkit/learn-ebm/what-is-grade/)

Key articles

- Di Lorenzo C, Colletti RB, Lehmann HP, et al; AAP Subcommittee, NASPGHAN Committee on Chronic Abdominal Pain. Chronic abdominal pain in children: a technical report of the American Academy of Pediatrics and the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J Pediatr Gastroenterol Nutr. 2005 Mar;40(3):249-61. Full text (https://journals.lww.com/jpgn/ Fulltext/2005/03000/Chronic_Abdominal_Pain_In_Children__A_Technical.2.aspx) Abstract
- Di Lorenzo C, Colletti RB, Lehmann HP, et al; American Academy of Pediatrics Subcommittee on Chronic Abdominal Pain; NASPGHAN Committee on Abdominal Pain. Chronic abdominal pain in children: a clinical report of the American Academy of Pediatrics and the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J Pediatr Gastroenterol Nutr. 2005 Mar;40(3):245-8. Full text (https://journals.lww.com/jpgn/Fulltext/2005/03000/Chronic_Abdominal_Pain_in_Children__A_Clinical.1.aspx) Abstract
- Tabbers MM, DiLorenzo C, Berger MY, et al; European Society for Pediatric Gastroenterology,
 Hepatology, and Nutrition; North American Society for Pediatric Gastroenterology. Evaluation and
 treatment of functional constipation in infants and children: evidence-based recommendations from
 ESPGHAN and NASPGHAN. J Pediatr Gastroenterol Nutr. 2014 Feb;58(2):258-74. Full text (https://
 journals.lww.com/jpgn/Fulltext/2014/02000/Evaluation_and_Treatment_of_Functional.24.aspx)
 Abstract
- Diercks DB, Adkins EJ, Harrison N, et al. American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of emergency department patients with suspected appendicitis. Approved by ACEP Board of Directors February 1, 2023. Ann Emerg Med. 2023 Jun;81(6):e115-52. Full text (https://www.annemergmed.com/article/S0196-0644(23)00029-X/fulltext) Abstract
- Dahabreh IJ, Adam GP, Halladay CW, et al. Diagnosis of right lower quadrant pain and suspected acute appendicitis. Dec 2015 [internet publication]. Full text (https://www.ncbi.nlm.nih.gov/books/ NBK355441) Abstract

References

- Koppen IJ, Vriesman MH, Saps M, et al. Prevalence of functional defecation disorders in children: a systematic review and meta-analysis. J Pediatr. 2018 Jul;198:121-30.e6. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29656863?tool=bestpractice.bmj.com)
- Zeevenhooven J, Koppen IJ, Benninga MA. The new Rome IV criteria for functional gastrointestinal disorders in infants and toddlers. Pediatr Gastroenterol Hepatol Nutr. 2017 Mar;20(1):1-13. Full text (https://pghn.org/DOIx.php?id=10.5223/pghn.2017.20.1.1) Abstract (http://www.ncbi.nlm.nih.gov/ pubmed/28401050?tool=bestpractice.bmj.com)

- 3. Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders: neonate/toddler. Gastroenterology. 2016 Feb 15;150(6):P1443-55. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27144631?tool=bestpractice.bmj.com)
- Wessel MA, Cobb JC, Jackson EB, et al. Paroxysmal fussing in infancy, sometimes called colic. Pediatrics. 1954 Nov;14(5):421-35. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/13214956? tool=bestpractice.bmj.com)
- Lucassen PL, Assendelft WJ, van Eijk JT, et al. Systematic review of the occurrence of infantile colic in the community. Arch Dis Child. 2001 May;84(5):398-403. Full text (https://adc.bmj.com/content/84/5/398) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/11316682? tool=bestpractice.bmj.com)
- 6. Omling E, Salö M, Saluja S, et al. Nationwide study of appendicitis in children. Br J Surg. 2019
 Nov;106(12):1623-31. Full text (https://bjssjournals.onlinelibrary.wiley.com/doi/full/10.1002/bjs.11298)
 Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31386195?tool=bestpractice.bmj.com)
- 7. Howell EC, Dubina ED, Lee SL. Perforation risk in pediatric appendicitis: assessment and management. Pediatric Health Med Ther. 2018 Oct 26;9:135-45. Full text (https://www.dovepress.com/perforation-risk-in-pediatric-appendicitis-assessment-and-management-peer-reviewed-fulltext-article-PHMT) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30464677?tool=bestpractice.bmj.com)
- 8. Alvarado A. A practical score for the early diagnosis of acute appendicitis. Ann Emerg Med. 1986 May;15(5):557-64. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/3963537?tool=bestpractice.bmj.com)
- 9. Samuel M. Pediatric appendicitis score. J Pediatr Surg. 2002 Jun;37(6):877-81. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/12037754?tool=bestpractice.bmj.com)
- 10. Cotton DM, Vinson DR, Vazquez-Benitez G, et al; Clinical Research on Emergency Services and Treatments (CREST) Network. Validation of the Pediatric Appendicitis Risk Calculator (pARC) in a community emergency department setting. Ann Emerg Med. 2019 Oct;74(4):471-80. Full text (https://www.annemergmed.com/article/S0196-0644(19)30346-4/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31229394?tool=bestpractice.bmj.com)
- 11. Kharbanda AB, Vazquez-Benitez G, Ballard DW, et al. Development and validation of a novel Pediatric Appendicitis Risk Calculator (pARC). Pediatrics. 2018 Apr;141(4):e20172699. Full text (https://publications.aap.org/pediatrics/article/141/4/e20172699/37779/Development-and-Validation-of-a-Novel-Pediatric) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29535251? tool=bestpractice.bmj.com)
- 12. Mandeville K, Pottker T, Bulloch B, et al. Using appendicitis scores in the pediatric ED. Am J Emerg Med. 2011 Nov;29(9):972-7. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/20674221? tool=bestpractice.bmj.com)
- 13. Di Saverio S, Podda M, De Simone B, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. World J Emerg Surg. 2020 Apr 15;15(1):27. Full

- text (https://wjes.biomedcentral.com/articles/10.1186/s13017-020-00306-3) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32295644?tool=bestpractice.bmj.com)
- 14. Li J. Revisiting delayed appendectomy in patients with acute appendicitis. World J Clin Cases. 2021 Jul 16;9(20):5372-90. Full text (https://www.wjgnet.com/2307-8960/full/v9/i20/5372.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34307591?tool=bestpractice.bmj.com)
- 15. Papandria D, Goldstein SD, Rhee D, et al. Risk of perforation increases with delay in recognition and surgery for acute appendicitis. J Surg Res. 2013 Oct;184(2):723-9. Full text (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4398569) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23290595?tool=bestpractice.bmj.com)
- 16. Manatakis DK, Aheimastos V, Antonopoulou MI, et al. Unfinished business: a systematic review of stump appendicitis. World J Surg. 2019 Nov;43(11):2756-61. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31375871?tool=bestpractice.bmj.com)
- 17. Sunkara T, Rawla P, Yarlagadda KS, et al. Eosinophilic gastroenteritis: diagnosis and clinical perspectives. Clin Exp Gastroenterol. 2019 Jun 5;12:239-53. Full text (https://www.dovepress.com/eosinophilic-gastroenteritis-diagnosis-and-clinical-perspectives-peer-reviewed-fulltext-article-CEG)

 Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31239747?tool=bestpractice.bmj.com)
- Salvadori M, Bertoni E. Update on hemolytic uremic syndrome: diagnostic and therapeutic recommendations. World J Nephrol. 2013 Aug 6;2(3):56-76. Full text (https:// www.wjgnet.com/2220-6124/full/v2/i3/56.htm) Abstract (http://www.ncbi.nlm.nih.gov/ pubmed/24255888?tool=bestpractice.bmj.com)
- Jiang J, Jiang B, Parashar U, et al. Childhood intussusception: a literature review. PLoS One. 2013;8(7):e68482. Full text (https://journals.plos.org/plosone/article?id=10.1371/ journal.pone.0068482) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23894308? tool=bestpractice.bmj.com)
- 20. Saulsbury FT. Clinical update: Henoch-Schönlein purpura. Lancet. 2007 Mar 24;369(9566):976-8.
- 21. Zhao Q, Yang Y, He SW, et al. Risk factors for intussusception in children with Henoch-Schönlein purpura: a case-control study. World J Clin Cases. 2021 Aug 6;9(22):6244-53. Full text (https://www.wjgnet.com/2307-8960/full/v9/i22/6244.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34434991?tool=bestpractice.bmj.com)
- 22. Leung AKC, Barankin B, Leong KF. Henoch-Schönlein purpura in children: an updated review. Curr Pediatr Rev. 2020;16(4):265-76. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32384035? tool=bestpractice.bmj.com)
- 23. Mateen MA, Saleem S, Rao PC, et al. Transient small bowel intussusceptions: ultrasound findings and clinical significance. Abdom Imaging. 2006 Jul-Aug;31(4):410-6. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16944032?tool=bestpractice.bmj.com)
- Hansen CC, Søreide K. Systematic review of epidemiology, presentation, and management of Meckel's diverticulum in the 21st century. Medicine (Baltimore). 2018
 Aug;97(35):e12154. Full text (https://journals.lww.com/md-journal/Fulltext/2018/08310/

- Systematic_review_of_epidemiology,_presentation,.91.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30170459?tool=bestpractice.bmj.com)
- 25. Elsayes KM, Menias CO, Harvin HJ, et al. Imaging manifestations of Meckel's diverticulum. AJR Am J Roentgenol. 2007 Jul;189(1):81-8. Full text (https://www.ajronline.org/doi/full/10.2214/AJR.06.1257)

 Abstract (http://www.ncbi.nlm.nih.gov/pubmed/17579156?tool=bestpractice.bmj.com)
- 26. Lin XK, Huang XZ, Bao XZ, et al. Clinical characteristics of Meckel diverticulum in children: a retrospective review of a 15-year single-center experience. Medicine (Baltimore). 2017 Aug;96(32):e7760. Full text (https://journals.lww.com/md-journal/Fulltext/2017/08110/ Clinical_characteristics_of_Meckel_diverticulum_in.43.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28796070?tool=bestpractice.bmj.com)
- 27. Karabulut R, Sonmez K, Turkyilmaz Z, et al. Negative appendectomy experience in children. Ir J Med Sci. 2011 Mar;180(1):55-8. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/20658324? tool=bestpractice.bmj.com)
- 28. Gross I, Siedner-Weintraub Y, Stibbe S, et al. Characteristics of mesenteric lymphadenitis in comparison with those of acute appendicitis in children. Eur J Pediatr. 2017 Feb;176(2):199-205. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27987102?tool=bestpractice.bmj.com)
- 29. Kyrklund K, Sloots CEJ, de Blaauw I, et al. ERNICA guidelines for the management of rectosigmoid Hirschsprung's disease. Orphanet J Rare Dis. 2020 Jun 25;15(1):164. Full text (https://ojrd.biomedcentral.com/articles/10.1186/s13023-020-01362-3) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32586397?tool=bestpractice.bmj.com)
- 30. Gosain A, Frykman PK, Cowles RA, et al. Guidelines for the diagnosis and management of Hirschsprung-associated enterocolitis. Pediatr Surg Int. 2017 May;33(5):517-21. Full text (https://pmc.ncbi.nlm.nih.gov/articles/PMC5395325) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28154902?tool=bestpractice.bmj.com)
- 31. Gershon EM, Rodriguez L, Arbizu RA. Hirschsprung's disease associated enterocolitis: a comprehensive review. World J Clin Pediatr. 2023 Jun 9;12(3):68-76. Full text (https://www.wjgnet.com/2219-2808/full/v12/i3/68.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/37342453?tool=bestpractice.bmj.com)
- 32. Shalaby MS, Kuti K, Walker G. Intestinal malrotation and volvulus in infants and children. BMJ. 2013 Nov 26;347:f6949. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24285798? tool=bestpractice.bmj.com)
- 33. Maas C, Hammer S, Kirschner HJ, et al. Late-onset volvulus without malrotation in extremely preterm infants--a case-control-study. BMC Pediatr. 2014 Nov 12;14:287. Full text (https://bmcpediatr.biomedcentral.com/articles/10.1186/s12887-014-0287-2) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/25388806?tool=bestpractice.bmj.com)
- 34. Destro F, Maestri L, Meroni M, et al. Colonic volvulus in children: surgical management of a challenging condition. Children (Basel). 2021 Oct 30;8(11):982. Full text (https://

- www.mdpi.com/2227-9067/8/11/982/htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34828695? tool=bestpractice.bmj.com)
- 35. Battersby C, Santhalingam T, Costeloe K, et al. Incidence of neonatal necrotising enterocolitis in high-income countries: a systematic review. Arch Dis Child Fetal Neonatal Ed. 2018 Mar;103(2):F182-9. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29317459?tool=bestpractice.bmj.com)
- 36. Neu J, Walker WA. Necrotizing enterocolitis. N Engl J Med. 2011 Jan 20;364(3):255-64. Full text (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3628622) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/21247316?tool=bestpractice.bmj.com)
- 37. Sullivan PB. Symposium: gastroenterology. Peptic ulcer disease in children. Paediatr Child Health. 2010 Oct;20(10):462-4. Full text (https://www.paediatricsandchildhealthjournal.co.uk/article/S1751-7222(10)00093-4/abstract)
- 38. Rinawi F, Assa A, Eliakim R, et al. The natural history of pediatric-onset IBD-unclassified and prediction of Crohn's disease reclassification: a 27-year study. Scand J Gastroenterol. 2017 May;52(5):558-63. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28128677? tool=bestpractice.bmj.com)
- 39. Carvalho RS, Abadom V, Dilworth HP, et al. Indeterminate colitis: a significant subgroup of pediatric IBD. Inflamm Bowel Dis. 2006 Apr;12(4):258-62. Full text (https://academic.oup.com/ibdjournal/article/12/4/258/4682671) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16633047?tool=bestpractice.bmj.com)
- 40. Ye Y, Manne S, Treem WR, et al. Prevalence of inflammatory bowel disease in pediatric and adult populations: recent estimates from large national databases in the United States, 2007-2016. Inflamm Bowel Dis. 2020 Mar 4;26(4):619-25. Full text (https://academic.oup.com/ibdjournal/article/26/4/619/5554163) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31504515?tool=bestpractice.bmj.com)
- 41. Feuerstein JD, Cheifetz AS. Crohn disease: epidemiology, diagnosis, and management. Mayo Clin Proc. 2017 Jul;92(7):1088-103. Full text (https://www.mayoclinicproceedings.org/article/S0025-6196(17)30313-0/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28601423?tool=bestpractice.bmj.com)
- 42. Torres J, Mehandru S, Colombel JF, et al. Crohn's disease. Lancet. 2017 Apr 29;389(10080):1741-55. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27914655?tool=bestpractice.bmj.com)
- 43. Molodecky NA, Soon IS, Rabi DM, et al. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. Gastroenterology. 2012 Jan;142(1):46-54.e42. Full text (https://www.gastrojournal.org/article/S0016-5085(11)01378-3/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/22001864?tool=bestpractice.bmj.com)
- 44. Singh P, Arora A, Strand TA, et al. Global prevalence of celiac disease: systematic review and meta-analysis. Clin Gastroenterol Hepatol. 2018 Jun;16(6):823-36.e2. Full text (https://www.cghjournal.org/article/S1542-3565(17)30783-8/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29551598?tool=bestpractice.bmj.com)

- 45. National Institute for Health and Care Excellence. Coeliac disease: recognition, assessment and management. Sep 2015 [internet publication]. Full text (https://www.nice.org.uk/guidance/ng20)
- 46. Al-Toma A, Volta U, Auricchio R, et al. European Society for the Study of Coeliac Disease (ESsCD) guideline for coeliac disease and other gluten-related disorders. United European Gastroenterol J. 2019 Jun;7(5):583-613. Full text (https://onlinelibrary.wiley.com/doi/10.1177/2050640619844125) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31210940?tool=bestpractice.bmj.com)
- 47. Santucci NR, Hyman PE, Harmon CM, et al. Biliary dyskinesia in children: a systematic review. J Pediatr Gastroenterol Nutr. 2017 Feb;64(2):186-93. Full text (https://journals.lww.com/jpgn/Fulltext/2017/02000/Biliary_Dyskinesia_in_Children__A_Systematic.7.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27472474?tool=bestpractice.bmj.com)
- 48. Abu-El-Haija M, Kumar S, Szabo F, et al. Classification of acute pancreatitis in the pediatric population: clinical report from the NASPGHAN Pancreas Committee. J Pediatr Gastroenterol Nutr. 2017 Jun;64(6):984-90. Full text (hhttps://onlinelibrary.wiley.com/doi/10.1097/MPG.000000000001583) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28333771?tool=bestpractice.bmj.com)
- 49. Suzuki M, Sai JK, Shimizu T. Acute pancreatitis in children and adolescents. World J Gastrointest Pathophysiol. 2014 Nov 15;5(4):416-26. Full text (https://www.wjgnet.com/2150-5330/full/v5/i4/16.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/25400985?tool=bestpractice.bmj.com)
- 50. Burgener FA, Meyers SP, Tan RK, et al. Differential diagnosis in magnetic resonance imaging. New York: Thieme; 2002:530.
- 51. Aslam S, Sohaib A, Reznek RH. Reticuloendothelial disorders: the spleen. In: Adam A, Dixon A, eds. Grainger and Allison's Diagnostic radiology. 5th ed. Philadelphia: Churchill Livingstone; 2008:1759-70.
- 52. Bradshaw CJ, Bandi AS, Muktar Z, et al. International study of the epidemiology of paediatric trauma: PAPSA Research Study. World J Surg. 2018 Jun;42(6):1885-94. Full text (https://link.springer.com/article/10.1007%2Fs00268-017-4396-6) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29282513?tool=bestpractice.bmj.com)
- 53. Ladomenou F, Bitsori M, Galanakis E. Incidence and morbidity of urinary tract infection in a prospective cohort of children. Acta Paediatr. 2015 Jul;104(7):e324-9. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/25736706?tool=bestpractice.bmj.com)
- 54. De Sanctis V, Soliman A, Bernasconi S, et al. Primary dysmenorrhea in adolescents: prevalence, impact and recent knowledge. Pediatr Endocrinol Rev. 2015 Dec;13(2):512-20. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26841639?tool=bestpractice.bmj.com)
- 55. Zhao LC, Lautz TB, Meeks JJ, et al. Pediatric testicular torsion epidemiology using a national database: incidence, risk of orchiectomy and possible measures toward improving the quality of care. J Urol. 2011 Nov;186(5):2009-13. Full text (https://www.doi.org/10.1016/j.juro.2011.07.024) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/21944120?tool=bestpractice.bmj.com)

- 56. Katz VL. 18. Benign gynecologic lesions: vulva, vagina, cervix, uterus, oviduct, ovary. In: Katz VL, Lentz GM, Lobo RA, et al., eds. Comprehensive gynecology. 5th ed. Philadelphia: Mosby; 2007:419-72.
- 57. Boyle KJ, Torrealday S. Benign gynecologic conditions. Surg Clin North Am. 2008 Apr;88(2):245-64. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/18381112?tool=bestpractice.bmj.com)
- 58. Schultz KA, Ness KK, Nagarajan R, et al. Adnexal masses in infancy and childhood. Clin Obstet Gynecol. 2006 Sep;49(3):464-79. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16885654? tool=bestpractice.bmj.com)
- 59. Emeksiz HC, Derinöz O, Akkoyun EB, et al. Age-specific frequencies and characteristics of ovarian cysts in children and adolescents. J Clin Res Pediatr Endocrinol. 2017 Mar 1;9(1):58-62. Full text (http://cms.galenos.com.tr/Uploads/Article_15636/58-62.pdf) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28044991?tool=bestpractice.bmj.com)
- 60. Dasgupta R, Renaud E, Goldin AB, et al. Ovarian torsion in pediatric and adolescent patients: A systematic review. J Pediatr Surg. 2018 Jul;53(7):1387-91. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29153467?tool=bestpractice.bmj.com)
- 61. Trent M. Pelvic inflammatory disease. Pediatr Rev. 2013 Apr;34(4):163-72. Full text (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4530285) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23547062?tool=bestpractice.bmj.com)
- 62. Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. MMWR Recomm Rep. 2021 Jul 23;70(4):1-187. Full text (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8344968) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34292926? tool=bestpractice.bmj.com)
- 63. Centers for Disease Control and Prevention. Stillbirth. Oct 2024 [internet publication]. Full text (https://www.cdc.gov/stillbirth/about)
- 64. Royal College of Obstetricians and Gynaecologists. Recurrent miscarriage (green-top guideline no. 17). Jun 2023 [internet publication]. Full text (https://www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/recurrent-miscarriage-green-top-guideline-no-17)
- 65. American College of Obstetrics and Gynaecology. Early pregnancy loss. Practice bulletin 200. Nov 2018 [internet publication]. Full text (https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2018/11/early-pregnancy-loss)
- 66. Bouyer J, Coste J, Fernandez H, et al. Sites of ectopic pregnancy: a 10 year population-based study of 1800 cases. Hum Reprod. 2002 Dec;17(12):3224-30. Full text (https://academic.oup.com/humrep/article/17/12/3224/569616) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/12456628?tool=bestpractice.bmj.com)
- 67. Gaskins AJ, Missmer SA, Rich-Edwards JW, et al. Demographic, lifestyle, and reproductive risk factors for ectopic pregnancy. Fertil Steril. 2018 Dec;110(7):1328-37. Full text (https://www.fertstert.org/

article/S0015-0282(18)31831-4/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30503132? tool=bestpractice.bmj.com)

- 68. Kevat PM, Morpeth M, Graham H, et al. A systematic review of the clinical features of pneumonia in children aged 5-9 years: implications for guidelines and research. J Glob Health. 2022 Mar 26;12:10002. Full text (https://jogh.org/documents/2022/jogh-12-10002.pdf) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/35356655?tool=bestpractice.bmj.com)
- 69. Sectish TC, Prober CG. Pneumonia. In: Behrman RE, Kliegman RM, Jenson HB, eds. Nelson textbook of pediatrics. 18th ed. Philadelphia: WB Saunders; 2007:1795-800.
- 70. Hyams JS, Di Lorenzo C, Saps M, et al. Childhood functional gastrointestinal disorders: child/adolescent. Gastroenterology. 2016 May;150(6):P1456-68.e2. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27144632?tool=bestpractice.bmj.com)
- 71. Drossman DA, Chang L, Chey WD, et al. Rome IV: functional gastrointestinal disorders disorders of gut-brain interaction. 4th ed. Raleigh, NC: Rome Foundation; 2017.
- 72. Baaleman DF, Di Lorenzo C, Benninga MA, et al. The effects of the Rome IV criteria on pediatric gastrointestinal practice. Curr Gastroenterol Rep. 2020 Mar 19;22(5):21. Full text (https://link.springer.com/article/10.1007/s11894-020-00760-8) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32193635?tool=bestpractice.bmj.com)
- 73. Boronat AC, Ferreira-Maia AP, Matijasevich A, et al. Epidemiology of functional gastrointestinal disorders in children and adolescents: a systematic review. World J Gastroenterol. 2017 Jun 7;23(21):3915-27. Full text (https://www.wjgnet.com/1007-9327/full/v23/i21/3915.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28638232?tool=bestpractice.bmj.com)
- 74. Di Lorenzo C, Colletti RB, Lehmann HP, et al; AAP Subcommittee, NASPGHAN Committee on Chronic Abdominal Pain. Chronic abdominal pain in children: a technical report of the American Academy of Pediatrics and the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J Pediatr Gastroenterol Nutr. 2005 Mar;40(3):249-61. Full text (https://journals.lww.com/jpgn/Fulltext/2005/03000/Chronic_Abdominal_Pain_In_Children__A_Technical.2.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/15735476?tool=bestpractice.bmj.com)
- 75. Di Lorenzo C, Colletti RB, Lehmann HP, et al; American Academy of Pediatrics Subcommittee on Chronic Abdominal Pain; NASPGHAN Committee on Abdominal Pain. Chronic abdominal pain in children: a clinical report of the American Academy of Pediatrics and the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J Pediatr Gastroenterol Nutr. 2005 Mar;40(3):245-8. Full text (https://journals.lww.com/jpgn/Fulltext/2005/03000/Chronic_Abdominal_Pain_in_Children__A_Clinical.1.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/15735475?tool=bestpractice.bmj.com)
- 76. American College of Radiology. ACR appropriateness criteria: suspected appendicitis child. 2018 [internet publication]. Full text (https://acsearch.acr.org/docs/3105874/Narrative)
- 77. Cho SU, Oh SK. Accuracy of ultrasound for the diagnosis of acute appendicitis in the emergency department: a systematic review. Medicine (Baltimore). 2023 Mar 31;102(13):e33397. Full text (https://journals.lww.com/md-journal/fulltext/2023/03310/

- accuracy_of_ultrasound_for_the_diagnosis_of_acute.51.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/37000097?tool=bestpractice.bmj.com)
- 78. Kumar SS, Collings AT, Lamm R, et al. SAGES guideline for the diagnosis and treatment of appendicitis. Surg Endosc. 2024 Jun;38(6):2974-94. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/38740595?tool=bestpractice.bmj.com)
- 79. Fraser JD, Aguayo P, Sharp SW, et al. Accuracy of computed tomography in predicting appendiceal perforation. J Pediatr Surg. 2010 Jan;45(1):231-5. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/20105609?tool=bestpractice.bmj.com)
- 80. The Royal College of Surgeons of England; British Association of Paediatric Surgeons.

 Commissioning guide: paediatric emergency appendicectomy. Jul 2014 [internet publication]. Full text (https://www.rcseng.ac.uk/standards-and-research/commissioning/commissioning-guides/topics)
- 81. Hackam DJ, Upperman J, Grikscheit T, et al. Pediatric surgery: gastrointestinal tract. In: Schwartz's principles of surgery. 11th ed. New York: McGraw-Hill; 2019.
- 82. McCollough M, Sharieff GQ. Abdominal pain in children. Pediatr Clin North Am. 2006 Feb;53(1):107-37, vi. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16487787? tool=bestpractice.bmj.com)
- 83. Gluckman S, Karpelowsky J, Webster AC, et al. Management for intussusception in children.

 Cochrane Database Syst Rev. 2017 Jun 1;(6):CD006476. Full text (https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD006476.pub3/full) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28567798?tool=bestpractice.bmj.com)
- 84. Gray MP, Li SH, Hoffmann RG, et al. Recurrence rates after intussusception enema reduction: a meta-analysis. Pediatrics. 2014 Jul;134(1):110-9. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24935997? tool=bestpractice.bmj.com)
- 85. Cantor RM. Pediatric trauma. In: Marx JA, Hockberger RS, Walls RM, eds. Rosen's emergency medicine: concepts and clinical practice. 6th ed. Philadelphia: Mosby; 2006:267-81.
- 86. Dayan PS, Klein BL. Chapter 71: acute care of the victim of multiple trauma. In: Behrman RE, Kliegman RM, Jenson HB, eds. Nelson textbook of pediatrics. 18th ed. Philadelphia: WB Saunders; 2007.
- 87. Rey-Bellet Gasser C, Gehri M, Joseph JM, et al. Is it ovarian torsion? A systematic literature review and evaluation of prediction signs. Pediatr Emerg Care. 2016 Apr;32(4):256-61. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26855342?tool=bestpractice.bmj.com)
- 88. Grunau GL, Harris A, Buckley J, et al. Diagnosis of ovarian torsion: is it time to forget about Doppler? J Obstet Gynaecol Can. 2018 Jul;40(7):871-5. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29681508?tool=bestpractice.bmj.com)
- 89. Oelsner G, Shashar D. Adnexal torsion. Clin Obstet Gynecol. 2006 Sep;49(3):459-63. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16885653?tool=bestpractice.bmj.com)

- 90. Breech LL, Hillard PJ. Adnexal torsion in pediatric and adolescent girls. Curr Opin Obstet Gynecol. 2005 Oct;17(5):483-9. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16141762? tool=bestpractice.bmj.com)
- 91. Kives S, Gascon S, Dubuc É, et al. No. 341 diagnosis and management of adnexal torsion in children, adolescents, and adults. J Obstet Gynaecol Can. 2017 Feb;39(2):82-90. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28241927?tool=bestpractice.bmj.com)
- 92. American College of Obstetricians and Gynecologists. Adnexal torsion in adolescents: ACOG Committee Opinion No. 783. Obstet Gynecol. 2019 Aug;134(2):e56-63. Full text (https://journals.lww.com/greenjournal/Fulltext/2019/08000/Adnexal_Torsion_in_Adolescents__ACOG_Committee.45.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31348225?tool=bestpractice.bmj.com)
- 93. Geimanaite L, Trainavicius K. Ovarian torsion in children: management and outcomes. J Pediatr Surg. 2013 Sep;48(9):1946-53. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24074673? tool=bestpractice.bmj.com)
- 94. European Association of Urology. Paediatric urology. 2022 [internet publication]. Full text (https://uroweb.org/guidelines/paediatric-urology)
- 95. Altinkilic B, Pilatz A, Weidner W. Detection of normal intratesticular perfusion using color coded duplex sonography obviates need for scrotal exploration in patients with suspected testicular torsion. J Urol. 2013 May;189(5):1853-8. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23219539? tool=bestpractice.bmj.com)
- 96. Sartelli M, Chichom-Mefire A, Labricciosa FM, et al. The management of intra-abdominal infections from a global perspective: 2017 WSES guidelines for management of intra-abdominal infections. World J Emerg Surg. 2017 Jul 10;12:29. Full text (https://wjes.biomedcentral.com/articles/10.1186/s13017-017-0141-6) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28702076?tool=bestpractice.bmj.com)
- 97. Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). JAMA. 2016 Feb 23;315(8):801-10. Full text (https://jamanetwork.com/journals/jama/fullarticle/2492881) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26903338?tool=bestpractice.bmj.com)
- 98. National Institute for Health and Care Excellence. Suspected sepsis: recognition, diagnosis and early management. Mar 2024 [internet publication]. Full text (https://www.nice.org.uk/guidance/ng51)
- 99. Weiss SL, Peters MJ, Alhazzani W, et al. Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children. Intensive Care Med. 2020 Feb;46(suppl 1):10-67. Full text (https://link.springer.com/article/10.1007/s00134-019-05878-6) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32030529?tool=bestpractice.bmj.com)
- 100. Schlapbach LJ, Watson RS, Sorce LR, et al. International consensus criteria for pediatric sepsis and septic shock. JAMA. 2024 Feb 27;331(8):665-74. Full text (https://pmc.ncbi.nlm.nih.gov/articles/PMC10900966) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/38245889?tool=bestpractice.bmj.com)

- 101. Royal College of Physicians. National Early Warning Score (NEWS) 2. Dec 2017 [internet publication]. Full text (https://www.rcplondon.ac.uk/projects/outputs/national-early-warning-score-news-2)
- 102. American College of Emergency Physicians (ACEP) Expert Panel on Sepsis. DART: an evidence-driven tool to guide the early recognition and treatment of sepsis and septic shock [internet publication]. Full text (https://poctools.acep.org/POCTool/Sepsis(DART)/276ed0a9-f24d-45f1-8d0c-e908a2758e5a)
- 103. Academy of Medical Royal Colleges. Statement on the initial antimicrobial treatment of sepsis. May 2022 [internet publication]. Full text (https://www.aomrc.org.uk/wp-content/uploads/2022/05/ Statement_on_the_initial_antimicrobial_treatment_of_sepsis_0522.pdf)
- 104. Greear GM, Romano MF, Katz MH, et al. Testicular torsion: epidemiological risk factors for orchiectomy in pediatric and adult patients. Int J Impot Res. 2021 Mar;33(2):184-90. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32683416?tool=bestpractice.bmj.com)
- 105. Kim JS. Acute abdominal pain in children. Pediatr Gastroenterol Hepatol Nutr. 2013 Dec;16(4):219-24. Full text (https://www.doi.org/10.5223/pghn.2013.16.4.219) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24511517?tool=bestpractice.bmj.com)
- 106. Tabbers MM, DiLorenzo C, Berger MY, et al; European Society for Pediatric Gastroenterology, Hepatology, and Nutrition; North American Society for Pediatric Gastroenterology. Evaluation and treatment of functional constipation in infants and children: evidence-based recommendations from ESPGHAN and NASPGHAN. J Pediatr Gastroenterol Nutr. 2014 Feb;58(2):258-74. Full text (https://journals.lww.com/jpgn/Fulltext/2014/02000/Evaluation_and_Treatment_of_Functional.24.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24345831?tool=bestpractice.bmj.com)
- National Institute for Health and Care Excellence. Constipation in children and young people: diagnosis and management. Jul 2017 [internet publication]. Full text (https://www.nice.org.uk/guidance/CG99)
- 108. Mattoo TK, Shaikh N, Nelson CP. Contemporary management of urinary tract infection in children. Pediatrics. 2021 Feb;147(2). Full text (https://www.doi.org/10.1542/peds.2020-012138) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/33479164?tool=bestpractice.bmj.com)
- 109. Morello W, La Scola C, Alberici I, et al. Acute pyelonephritis in children. Pediatr Nephrol. 2016 Aug;31(8):1253-65. Full text (https://www.doi.org/10.1007/s00467-015-3168-5) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26238274?tool=bestpractice.bmj.com)
- 110. Birmole BJ, Kulkarni BK, Vaidya MM, et al. Splenic cyst. J. Postgrad Med. 1993 Jan-Mar;39(1):40-1. Full text (https://www.jpgmonline.com/article.asp? issn=0022-3859;year=1993;volume=39;issue=1;spage=40;epage=1;aulast=Birmole) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/8295148?tool=bestpractice.bmj.com)
- 111. American College of Surgeons. Advanced trauma life support student course manual, 10th edition. Chicago, IL; 2018. Full text (https://www.facs.org/quality-programs/trauma/education/advanced-trauma-life-support/?page=1)

- 112. Ekabe CJ, Kehbila J, Njim T, et al. Chlamydia trachomatis-induced Fitz-Hugh-Curtis syndrome: a case report. BMC Res Notes. 2017 Jan 3;10(1):10. Full text (https://bmcresnotes.biomedcentral.com/articles/10.1186/s13104-016-2357-z) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28057084? tool=bestpractice.bmj.com)
- 113. Zeeshan M, Hamidi M, O'Keeffe T, et al. Pediatric liver injury: physical examination, fast and serum transaminases can serve as a guide. J Surg Res. 2019 Oct;242:151-6. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31078899?tool=bestpractice.bmj.com)
- 114. Tenner S, Vege SS, Sheth SG, et al. American College of Gastroenterology guidelines: management of acute pancreatitis. Am J Gastroenterol. 2024 Mar 1;119(3):419-37. Full text (https://journals.lww.com/ajg/fulltext/2024/03000/american_college_of_gastroenterology_guidelines_.14.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/38857482?tool=bestpractice.bmj.com)
- 115. Chin A, De Wit K. Inability of a negative faecal occult blood test to rule out intussusception in children. Emerg Med J. 2017 Jul;34(7):489-90. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28652288? tool=bestpractice.bmj.com)
- 117. Shane AL, Mody RK, Crump JA, et al. 2017 Infectious Diseases Society of America clinical practice guidelines for the diagnosis and management of infectious diarrhea. Clin Infect Dis. 2017 Nov 29;65(12):e45-80. Full text (https://academic.oup.com/cid/article/65/12/e45/4557073) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29053792?tool=bestpractice.bmj.com)
- 118. Nabarro LE, McCann N, Herdman MT, et al. British infection association guidelines for the diagnosis and management of enteric fever in England. J Infect. 2022 Apr;84(4):469-89. Full text (https://www.journalofinfection.com/article/S0163-4453(22)00013-5/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/35038438?tool=bestpractice.bmj.com)
- 119. Broekaert IJ, Borrelli O, Dolinsek J, et al. An ESPGHAN position paper on the use of breath testing in paediatric gastroenterology. J Pediatr Gastroenterol Nutr. 2022 Jan 1;74(1):123-37. Full text (https://onlinelibrary.wiley.com/doi/10.1097/MPG.0000000000003245) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34292218?tool=bestpractice.bmj.com)
- 120. Jones NL, Koletzko S, Goodman K, et al. Joint ESPGHAN/NASPGHAN guidelines for the management of Helicobacter pylori in children and adolescents (update 2016). J Pediatr Gastroenterol Nutr. 2017 Jun;64(6):991-1003. Full text (https://onlinelibrary.wiley.com/doi/10.1097/MPG.000000000001594) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28541262? tool=bestpractice.bmj.com)
- 121. Mitsuyama K, Niwa M, Takedatsu H, et al. Antibody markers in the diagnosis of inflammatory bowel disease. World J Gastroenterol. 2016 Jan 21;22(3):1304-10. Full text (https://

- www.wjgnet.com/1007-9327/full/v22/i3/1304.htm) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26811667?tool=bestpractice.bmj.com)
- 122. Reese GE, Constantinides VA, Simillis C, et al. Diagnostic precision of anti-Saccharomyces cerevisiae antibodies and perinuclear antineutrophil cytoplasmic antibodies in inflammatory bowel disease. Am J Gastroenterol. 2006 Oct;101(10):2410-22. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/16952282? tool=bestpractice.bmj.com)
- 123. Lee JY, Kim JH, Choi SJ, et al. Point-of-care ultrasound may be useful for detecting pediatric intussusception at an early stage. BMC Pediatr. 2020 Apr 13;20(1):155. Full text (https://bmcpediatr.biomedcentral.com/articles/10.1186/s12887-020-02060-6) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32284038?tool=bestpractice.bmj.com)
- 124. Hom J, Kaplan C, Fowler S, et al. Evidence-based diagnostic test accuracy of history, physical examination, and imaging for intussusception: a systematic review and meta-analysis. Pediatr Emerg Care. 2022 Jan 1;38(1):e225-30. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32941364? tool=bestpractice.bmj.com)
- 125. Holmes JF, Brant WE, Bond WF, et al. Emergency department ultrasonography in the evaluation of hypotensive and normotensive children with blunt abdominal trauma. J Pediatr Surg. 2001 Jul;36(7):968-73. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/11431759?tool=bestpractice.bmj.com)
- 126. Calder BW, Vogel AM, Zhang J, et al. Focused assessment with sonography for trauma in children after blunt abdominal trauma: a multi-institutional analysis. J Trauma Acute Care Surg. 2017 Aug;83(2):218-24. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28590347? tool=bestpractice.bmj.com)
- 127. Holmes JF, Kelley KM, Wootton-Gorges SL, et al. Effect of abdominal ultrasound on clinical care, outcomes, and resource use among children with blunt torso trauma: a randomized clinical trial. JAMA. 2017 Jun 13;317(22):2290-6. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28609532? tool=bestpractice.bmj.com)
- 128. Lee KH, Lee S, Park JH, et al. Risk of hematologic malignant neoplasms from abdominopelvic computed tomographic radiation in patients who underwent appendectomy. JAMA Surg. 2021 Apr 1;156(4):343-51. Full text (https://jamanetwork.com/journals/jamasurgery/fullarticle/2775351) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/33471110?tool=bestpractice.bmj.com)
- 129. Diercks DB, Adkins EJ, Harrison N, et al. American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of emergency department patients with suspected appendicitis. Approved by ACEP Board of Directors February 1, 2023. Ann Emerg Med. 2023 Jun;81(6):e115-52. Full text (https://www.annemergmed.com/article/S0196-0644(23)00029-X/fulltext) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/37210169?tool=bestpractice.bmj.com)
- 130. Kulaylat AN, Moore MM, Engbrecht BW, et al. An implemented MRI program to eliminate radiation from the evaluation of pediatric appendicitis. J Pediatr Surg. 2015 Aug;50(8):1359-63. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/25783291?tool=bestpractice.bmj.com)
- 131. Mushtaq R, Desoky SM, Morello F, et al. First-line diagnostic evaluation with MRI of children suspected of having acute appendicitis. Radiology. 2019 Apr;291(1):170-7. Full text (https://

- pubs.rsna.org/doi/10.1148/radiol.2019181959) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30747595?tool=bestpractice.bmj.com)
- 132. Lee SM, Kim WS, Choi YH. Pediatric magnetic resonance enterography: focused on Crohn's disease. Pediatr Gastroenterol Hepatol Nutr. 2015 Sep;18(3):149-59. Full text (https://pghn.org/DOIx.php? id=10.5223/pghn.2015.18.3.149) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26473134? tool=bestpractice.bmj.com)
- 133. Brody AS, Frush DP, Huda W, et al; American Academy of Pediatrics Section on Radiology. Radiation risk to children from computed tomography. Pediatrics. 2007 Sep;120(3):677-82. Full text (https://publications.aap.org/pediatrics/article/120/3/677/71216/Radiation-Risk-to-Children-From-Computed)

 Abstract (http://www.ncbi.nlm.nih.gov/pubmed/17766543?tool=bestpractice.bmj.com)
- 134. Carney DE, Kokoska ER, Grosfeld JL, et al. Predictors of successful outcome after cholecystectomy for biliary dyskinesia. J Pediatr Surg. 2004 Jun;39(6):813-6. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/15185202?tool=bestpractice.bmj.com)
- Coluccio M, Claffey AJ, Rothstein DH. Biliary dyskinesia: fact or fiction? Semin Pediatr Surg. 2020 Aug;29(4):150947. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32861451? tool=bestpractice.bmj.com)
- 136. Nasri B, Glass T, Singh K, et al. Biliary hyperkinesia: an indication for cholecystectomy? Surg Endosc. 2021 Jul;35(7):3244-8. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32632487? tool=bestpractice.bmj.com)
- 137. Subcommittee on urinary tract infection. Reaffirmation of AAP clinical practice guideline: the diagnosis and management of the initial urinary tract infection in febrile infants and young children 2-24 months of age. Pediatrics. 2016 Dec;138(6):e20163026. Full text (https://publications.aap.org/pediatrics/article/138/6/e20163026/52686/Reaffirmation-of-AAP-Clinical-Practice-Guideline) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27940735?tool=bestpractice.bmj.com)
- 138. National Institute for Health and Care Excellence. Urinary tract infection in under 16s: diagnosis and management. July 2022 [internet publication]
- 139. Ammenti A, Alberici I, Brugnara M, et al; Italian Society of Pediatric Nephrology. Updated Italian recommendations for the diagnosis, treatment and follow-up of the first febrile urinary tract infection in young children. Acta Paediatr. 2020 Feb;109(2):236-47. Full text (https://onlinelibrary.wiley.com/doi/10.1111/apa.14988) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31454101?tool=bestpractice.bmj.com)
- 140. Webster J, Osborne S, Rickard CM, et al. Clinically-indicated replacement versus routine replacement of peripheral venous catheters. Cochrane Database Syst Rev. 2019 Jan 23;1(1):CD007798. Full text (https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD007798.pub5/full) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30671926?tool=bestpractice.bmj.com)
- 141. Simundic AM, Bölenius K, Cadamuro J, et al. Joint EFLM-COLABIOCLI recommendation for venous blood sampling. Clin Chem Lab Med 2018;56(12):2015-38. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/30004902?tool=bestpractice.bmj.com)

- 142. Zouari M, Louati H, Abid I, et al. C-reactive protein value is a strong predictor of acute appendicitis in young children. Am J Emerg Med. 2018 Jul;36(7):1319-20. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/29217177?tool=bestpractice.bmj.com)
- 143. Dahabreh IJ, Adam GP, Halladay CW, et al. Diagnosis of right lower quadrant pain and suspected acute appendicitis. Dec 2015 [internet publication]. Full text (https://www.ncbi.nlm.nih.gov/books/NBK355441) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/27054223?tool=bestpractice.bmj.com)
- 144. College of American Pathologists. Urinalysis checklist. Apr 2014 [internet publication]. Full text (https://webapps.cap.org/apps/docs/education/OnlineCourseContent/2014/TLTM/URN04212014.PDF)
- 145. Downs SM. Technical report: urinary tract infections in febrile infants and young children. The Urinary Tract Subcommittee of the American Academy of Pediatrics Committee on Quality Improvement. Pediatrics. 1999 Apr;103(4):e54. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/10103346? tool=bestpractice.bmj.com)
- 146. Whiting P, Westwood M, Watt I, et al. Rapid tests and urine sampling techniques for the diagnosis of urinary tract infection (UTI) in children under five years: a systematic review. BMC Pediatr. 2005 Apr 5;5(1):4. Full text (https://bmcpediatr.biomedcentral.com/articles/10.1186/1471-2431-5-4) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/15811182?tool=bestpractice.bmj.com)
- 147. National Institute for Health and Care Excellence. Urinary tract infection in under 16s: diagnosis and management. Oct 2018 [internet publication].
- 148. Evers MB. Meckel's diverticulum. In: Townsend CM Jr, Beauchamp RD, Evers BM, et al, eds. Sabiston textbook of surgery. 18th ed. Philadelphia: Saunders Elsevier; 2007:1321-3.
- 149. Walsham NE, Sherwood RA. Fecal calprotectin in inflammatory bowel disease. Clin Exp Gastroenterol. 2016;9:21-9. Full text (https://www.dovepress.com/fecal-calprotectin-in-inflammatory-bowel-disease-peer-reviewed-fulltext-article-CEG) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26869808?tool=bestpractice.bmj.com)
- 150. Cohen RB, Olafson SN, Krupp J, et al. Timing of Gastrografin administration in the management of adhesive small bowel obstruction (ASBO): does it matter? Surgery. 2021 Aug;170(2):596-602. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/33836900?tool=bestpractice.bmj.com)
- 151. Maung AA, Johnson DC, Piper GL, et al. Evaluation and management of small-bowel obstruction: an Eastern Association for the Surgery of Trauma practice management guideline. J Trauma Acute Care Surg. 2012 Nov;73(5 suppl 4):S362-9. Full text (https://journals.lww.com/jtrauma/Fulltext/2012/11004/Evaluation_and_management_of_small_bowel.14.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23114494?tool=bestpractice.bmj.com)
- 152. Maheshwari A. Role of platelets in neonatal necrotizing enterocolitis. Pediatr Res. 2021

 Apr;89(5):1087-93. Full text (https://www.nature.com/articles/s41390-020-1038-8) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/32601461?tool=bestpractice.bmj.com)

- 153. Chan B, Gordon S, Yang M, et al. Abdominal ultrasound assists the diagnosis and management of necrotizing enterocolitis. Adv Neonatal Care. 2021 Oct 1;21(5):365-70. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34469367?tool=bestpractice.bmj.com)
- 154. Rubio-Tapia A, Hill ID, Kelly CP, et al. ACG clinical guidelines: diagnosis and management of celiac disease. Am J Gastroenterol. 2013 May;108(5):656-76; quiz 677. Full text (https://journals.lww.com/ajg/fulltext/2013/05000/acg_clinical_guidelines__diagnosis_and_management.7.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/23609613?tool=bestpractice.bmj.com)
- 155. Husby S, Koletzko S, Korponay-Szabó I, et al. European Society Paediatric Gastroenterology, Hepatology and Nutrition guidelines for diagnosing coeliac disease 2020. J Pediatr Gastroenterol Nutr. 2020 Jan;70(1):141-56. Full text (https://journals.lww.com/jpgn/Fulltext/2020/01000/ European_Society_Paediatric_Gastroenterology,.24.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/31568151?tool=bestpractice.bmj.com)
- 156. Rompianesi G, Hann A, Komolafe O, et al. Serum amylase and lipase and urinary trypsinogen and amylase for diagnosis of acute pancreatitis. Cochrane Database Syst Rev. 2017 Apr 21;(4):CD012010. Full text (https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012010.pub2/full) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/28431198?tool=bestpractice.bmj.com)
- 157. Goldstein B, Giroir B, Randolph A, et al; International Consensus Conference on Pediatric Sepsis. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. Pediatr Crit Care Med. 2005 Jan;6(1):2-8. Abstract (http://www.ncbi.nlm.nih.gov/pubmed/15636651?tool=bestpractice.bmj.com)
- 158. Mahajan P, Grzybowski M, Chen X, et al. Procalcitonin as a marker of serious bacterial infections in febrile children younger than 3 years old. Acad Emerg Med. 2014 Feb;21(2):171-9. Full text (https://onlinelibrary.wiley.com/doi/10.1111/acem.12316) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/24673673?tool=bestpractice.bmj.com)
- 159. Milcent K, Faesch S, Gras-Le Guen C, et al. Use of procalcitonin assays to predict serious bacterial infection in young febrile infants. JAMA Pediatr. 2016 Jan;170(1):62-9. Full text (https://jamanetwork.com/journals/jamapediatrics/fullarticle/2472718) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/26595253?tool=bestpractice.bmj.com)
- 160. Pantell RH, Roberts KB, Adams WG, et al; Subcommittee on Febrile Infants. Evaluation and management of well-appearing febrile infants 8 to 60 days old. Pediatrics. 2021 Aug;148(2):e2021052228. Full text (https://publications.aap.org/pediatrics/article/148/2/e2021052228/179783/Clinical-Practice-Guideline-Evaluation-and) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/34281996?tool=bestpractice.bmj.com)
- 161. Qin KR, Qu LG. Diagnosing with a TWIST: systematic review and meta-analysis of a testicular torsion risk score. J Urol. 2022 Jul;208(1):62-70. Full text (https://journals.lww.com/ajg/fulltext/2013/05000/acg_clinical_guidelines__diagnosis_and_management.7.aspx) Abstract (http://www.ncbi.nlm.nih.gov/pubmed/35238603?tool=bestpractice.bmj.com)

162. American College of Radiology. ACR appropriateness criteria: acute onset of scrotal pain - without trauma, without antecedent mass. 2018 [internet publication]. Full text (https://acsearch.acr.org/docs/69363/Narrative)

Images




Figure 1: Necrotic appendix




Figure 2: CT scan demonstrating faecalith (white arrow) outside the lumen of the appendix consistent with perforated appendix




Figure 3: Intussusception: blood vessels become trapped between layers of intestine, leading to reduced blood supply, oedema, strangulation of bowel, and gangrene. Sepsis, shock, and death may eventually occur

Created by the BMJ Knowledge Centre




Figure 4: Intraoperative photo of Meckel's diverticulum



Figure 5: Abdominal x-ray of a neonate with abnormal stooling pattern and constipation. The dilated transverse and descending colon is suggestive of Hirschsprung's disease




Figure 6: Abdominal x-ray demonstrating double bubble gas pattern consistent with duodenal atresia




Figure 7: Infant with right groin bulge consistent with incarcerated inguinal hernia. The lack of overlying skin oedema and erythema does not rule out strangulation of the small intestine



Figure 8: Gallbladder ultrasound demonstrating cholelithiasis with characteristic shadowing



Figure 9: Abdominal x-ray with opacities in the RUQ consistent with gallstones




Figure 10: CT scan of teenage girl presenting with mid-epigastric abdominal pain as a result of gallstone pancreatitis. The large fluid collection in the pancreatic bed (white arrow) and lack of pancreatic enhancement suggest liquefactive necrosis of the pancreas



Figure 11: CT scan demonstrating fluid-filled cyst within the spleen




Figure 12: Intraoperative photo of large splenic cyst



Figure 13: Young boy with right testicular pain. The testicle is swollen, tender, and erythematous as a result of torsion of the appendix testes. The clinical signs and symptoms mimic those of testicular torsion



Figure 14: Infant boy with swollen, tender, and erythematous left testicle. The testicle is retracted consistent with testicular torsion



Figure 15: Torsion of an appendix testis resulting in acute infarction




Figure 16: Intraoperative photo of ovarian mass that presented as ovarian torsion




Figure 17: CT scan of a young girl presenting with ovarian torsion. The large pelvic cystic lesion contains calcifications (white arrow) consistent with a teratoma or dermoid cyst



Figure 18: Contrast enema demonstrating ileocolic intussusception (black arrow)



Figure 19: Abdominal x-ray of a young boy with acute, severe abdominal pain, demonstrating stool throughout the colon and rectum




Figure 20: CT scan demonstrating intra-abdominal abscess consistent with perforated appendix



Figure 21: Upper GI contrast study demonstrating malrotation with volvulus. The duodenum fails to develop the normal anatomical C-loop. There is failure of contrast to pass, resulting in a characteristic bird beak consistent with acute mid-gut volvulus

Disclaimer

BMJ Best Practice is intended for licensed medical professionals. BMJ Publishing Group Ltd (BMJ) does not advocate or endorse the use of any drug or therapy contained within this publication nor does it diagnose patients. As a medical professional you retain full responsibility for the care and treatment of your patients and you should use your own clinical judgement and expertise when using this product.

This content is not intended to cover all possible diagnosis methods, treatments, follow up, drugs and any contraindications or side effects. In addition, since such standards and practices in medicine change as new data become available, you should consult a variety of sources. We strongly recommend that you independently verify specified diagnosis, treatments and follow-up and ensure it is appropriate for your patient within your region. In addition, with respect to prescription medication, you are advised to check the product information sheet accompanying each drug to verify conditions of use and identify any changes in dosage schedule or contraindications, particularly if the drug to be administered is new, infrequently used, or has a narrow therapeutic range. You must always check that drugs referenced are licensed for the specified use and at the specified doses in your region.

Information included in BMJ Best Practice is provided on an "as is" basis without any representations, conditions or warranties that it is accurate and up to date. BMJ and its licensors and licensees assume no responsibility for any aspect of treatment administered to any patients with the aid of this information. To the fullest extent permitted by law, BMJ and its licensors and licensees shall not incur any liability, including without limitation, liability for damages, arising from the content. All conditions, warranties and other terms which might otherwise be implied by the law including, without limitation, the warranties of satisfactory quality, fitness for a particular purpose, use of reasonable care and skill and non-infringement of proprietary rights are excluded.

Where BMJ Best Practice has been translated into a language other than English, BMJ does not warrant the accuracy and reliability of the translations or the content provided by third parties (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages). BMJ is not responsible for any errors and omissions arising from translation and adaptation or otherwise. Where BMJ Best Practice lists drug names, it does so by recommended International Nonproprietary Names (rINNs) only. It is possible that certain drug formularies might refer to the same drugs using different names.

Please note that recommended formulations and doses may differ between drug databases drug names and brands, drug formularies, or locations. A local drug formulary should always be consulted for full prescribing information.

Treatment recommendations in BMJ Best Practice are specific to patient groups. Care is advised when selecting the integrated drug formulary as some treatment recommendations are for adults only, and external links to a paediatric formulary do not necessarily advocate use in children (and vice-versa). Always check that you have selected the correct drug formulary for your patient.

Where your version of BMJ Best Practice does not integrate with a local drug formulary, you should consult a local pharmaceutical database for comprehensive drug information including contraindications, drug interactions, and alternative dosing before prescribing.

Interpretation of numbers

Regardless of the language in which the content is displayed, numerals are displayed according to the original English-language numerical separator standard. For example 4 digit numbers shall not include a comma nor a decimal point; numbers of 5 or more digits shall include commas; and numbers stated to be less than 1 shall be depicted using decimal points. See Figure 1 below for an explanatory table.

BMJ accepts no responsibility for misinterpretation of numbers which comply with this stated numerical separator standard.

This approach is in line with the guidance of the International Bureau of Weights and Measures Service.

Figure 1 – BMJ Best Practice Numeral Style

5-digit numerals: 10,000

4-digit numerals: 1000

numerals < 1: 0.25

Our full website and application terms and conditions can be found here: Website Terms and Conditions.

Contact us

+ 44 (0) 207 111 1105 support@bmj.com

BMJ BMA House Tavistock Square London WC1H 9JR UK



Contributors:

// Authors:

Charles L. Snyder, MD

Professor of Surgery

University of Missouri, Kansas City Children's Mercy Hospital, Kansas City, MO

DISCLOSURES: CLS declares that he has no competing interests.

Jeong Hyun, MD

Assistant Professor of Surgery

Stanford University, Lucile Packard Children's Hospital, Stanford, CA

DISCLOSURES: JH has several research grants for translational work with Crohn's disease for Stanford University.

// Acknowledgements:

Professor Charles L. Snyder and Dr Jeong Hyun would like to gratefully acknowledge Dr KuoJen Tsao, Dr Kathryn Tinsley Anderson, Dr Luke R. Putnam, Dr Shinil K. Shah, Dr Nathan D. Allison, and Dr Peter A. Walker, previous contributors to this topic.

DISCLOSURES: KT, KTA, LRP, SKS, NDA, and PAW declare that they have no competing interests.

// Peer Reviewers:

Kimberley Chien, MD

Associate Professor of Clinical Pediatrics Weill Cornell Medicine, New York, NY

DISCLOSURES: KC declares that she has no competing interests.

Ann O. Scheimann, MD, MBA

Associate Professor of Pediatrics

Johns Hopkins School of Medicine, Baltimore, MD

DISCLOSURES: AOS declares that she has no competing interests.

Karla Au. Yeung, MD

Assistant Professor

Pediatric Gastroenterology and Nutrition, The Johns Hopkins Hospital, Baltimore, MD DISCLOSURES: KAY declares that she has no competing interests.

Raymond G. Buick, MB BCh, BAO, FRCS (Edin), FRCS (Eng)

Consultant Paediatric Surgeon

Department of Paediatric Surgery, Birmingham Children's Hospital, Birmingham, UK

DISCLOSURES: RGB declares that he has no competing interests.