

BMJ Best Practice

Acute aspiration

Straight to the point of care



Last updated: Mar 25, 2025

Table of Contents

Overview	3
Summary	3
Definition	3
Theory	4
Epidemiology	4
Etiology	6
Pathophysiology	7
Case history	9
Diagnosis	12
Approach	12
History and exam	19
Risk factors	20
Tests	25
Differentials	30
Management	34
Approach	34
Treatment algorithm overview	37
Treatment algorithm	38
Primary prevention	41
Secondary prevention	42
Patient discussions	43
Follow up	44
Monitoring	44
Complications	44
Prognosis	45
Guidelines	46
Diagnostic guidelines	46
References	47
Images	60
Disclaimer	64

Summary

Acute aspiration is the inhalation of foreign material into the airways beyond the vocal cords with variable manifestations that result in aspiration bronchiolitis, aspiration pneumonitis, aspiration pneumonia, and acute respiratory distress syndrome (ARDS).

Usually occurs in patients with altered level of consciousness, dysphagia, or impaired cough reflex.

Screening for dysphagia is recommended for patients at risk of, or following, aspiration (including all patients after an acute stroke).

Clinical presentation and severity depends on the volume and type of the aspirated material as well as the host defenses, such as adequate cough reflex, intact mucociliary barrier, and respiratory reserve.

Anesthesia-related aspiration of gastric contents can be prevented by identifying patients susceptible to vomiting and reflux, minimizing gastric contents before surgery, avoiding drugs that reduce gastric emptying, minimizing emetic stimuli, and avoiding complete loss of protective reflexes from oversedation.

Antibiotics are not indicated early after aspiration of gastric contents but they should be considered if the pneumonitis does not resolve after 48 hours or if there is clear risk or susceptibility for infection.

Patients with neurologic deficits, infants, older patients (age >70 years), and debilitated patients with dysphagia may also aspirate barium sulfate during radiologic procedures, which can result in severe bronchiolitis and pneumonitis.

Definition

Aspiration is the inhalation of liquid or solid particles, particularly food or gastric contents, into the airways below the vocal cords.[1] It can result in aspiration pneumonitis (chemical injury) or aspiration pneumonia (infection), both of which can lead to acute respiratory distress syndrome (ARDS). See Aspiration pneumonia and Acute respiratory distress syndrome .[1] [2] If the central airways are completely occluded, asphyxiation may quickly ensue. See Foreign body aspiration . Awareness is increasing for silent or more subclinical chronic micro-aspiration, which has been implicated as a potential aggravating mechanism across a variety of disorders, such as interstitial lung disease, asthma, COPD, and chronic cough. Chronic micro-aspiration is not covered in this topic.[3]

Epidemiology

One of the common settings in which pulmonary aspiration is known to take place is the perioperative period. Various risk factors including reduced consciousness, prolonged supine positioning, and illness acuity can predispose an individual to aspiration of gastric contents. In the US, reported incidence of perioperative pulmonary aspiration in the adult population ranges from 1 in 3216 (in 1993) to 1 in 7103 (in 2006).^{[5] [6]}

Studies published in 1986 and 1993 reported that aspiration occurred in 1 in every 3000 cases of anesthesia and accounted for 10% to 30% of deaths associated with anesthesia.^{[2] [5] [7]} Although the incidence of aspiration is low, the extent of surgery has increased in people with comorbidities and the older population, and these patients are expected to have a higher incidence of aspiration.^[8]

In the pediatric population, one UK study published in 2013 notes an incidence of perioperative pulmonary aspiration of 1 in 4932 anesthetics in the elective setting and 1 in 4498 in the emergency setting.^[9] This suggests an improvement compared with two prior US landmark studies in children, which reported an incidence of 1 in 2632 aspirations during general anesthesia between 1985 and 1997, and 1 in 978 between 1988 and 1993.^{[10] [11]}

Despite a low incidence, perioperative pulmonary aspiration carries a risk of morbidity and mortality. In US anesthesia malpractice claims data, 5% were attributed to aspiration, with a 57% mortality directly attributable to anesthesia.^[12] Of those who aspirated, 61% had gastrointestinal obstruction or an acute abdomen.^[12]

The body of literature regarding acute aspiration outside of the perioperative time frame is smaller, and limited by the lack of specific diagnostic testing for aspiration when it is not witnessed. Between 5% and 15% of community-acquired pneumonia has been attributed to aspiration.^[2] Pneumonia due to aspiration events is more frequent in patients ages >80 years (10%) compared with patients ages <80 years (5%).^[13] Predisposing factors for aspiration included dysphagia, and impaired consciousness and gag reflex.^[13] Another study noted that of a cohort of patients admitted with pneumonia ages >70 years, about 55% were found to have dysphagia and aspiration on testing with a water swallow test.^[14] In the Lung Injury Prediction Score cohort, aspiration was the third leading cause of acute lung injury among at-risk patients admitted to the US hospitals.^[15]

Dysphagia may cause "acute on chronic" aspirations. According to one systematic review from 2005, the incidence of dysphagia (swallowing difficulties) following acute stroke was reported to vary from 37% to 78% depending on the site of the stroke and screening tools used to identify dysphagia.^{[16] [17]} A review from 2010 reported that up to 30% of older patients with dysphagia present with aspiration.^[18]

Among patients admitted to the hospital who are receiving enteral nutrition, the reported prevalence varies widely from 4.4% to nearly 90%, depending on how aspiration is defined (silent versus symptomatic), the method of diagnosis, the position of the feeding tube in the gastrointestinal tract (nasogastric or nasojejunal), and the type of feeding tube (nasogastric or gastrostomy).^{[19] [20]} One meta-analysis conducted in 2019 including 41 studies and involving 3248 participants suggests there is a lower incidence of pulmonary aspiration with postpyloric feeding tube positioning compared with gastric positioning.^[21] Clinically evident aspiration, however, is rare and is seen in <1% of patients given enteral nutrition.^[20]

In the setting of severe trauma, the incidence of gross aspiration can be as high as 38%.^[22]

Accidental aspiration of barium contrast medium during radiologic investigations is rare but can occur in up to 8% of children with GERD.^[23] The severity of pulmonary damage depends on the density of the suspension,

with high-density barium sulfate causing the most damage and being potentially fatal, especially in older patients.[4] [24] [25]



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood



Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed

From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Etiology

Physiologic mechanisms that normally reduce the risk of aspiration include the gastroesophageal junction, upper esophageal sphincter, and protective laryngeal reflexes. These protective mechanisms can be affected by several conditions, age, or drugs.[26]

Aspiration of food and liquids is more common in:[27]

- Patients with oropharyngeal dysphagia, especially when it is due to stroke, Parkinson disease, dementia, or cervical spine surgery[28][29][30]

- Older patients
- Patients who are taking sedatives
- Patients being fed by a gastric tube[29]
- Patients who depend on others for feeding
- Current smokers
- Patients taking >8 drugs
- Pregnant women.

Among critically ill patients, the major risk factors for aspiration include:[31]

- Documented previous episode of aspiration
- Decreased level of consciousness (Glasgow coma scale score <9 or a high level of sedation)
- Neuromuscular disease, or congenital or acquired structural abnormalities of the aerodigestive tract
- Endotracheal intubation
- Vomiting
- Persistently high gastric residual volume
- Supine positioning.

Additional risk factors include the presence of a nasoenteric tube, intermittent feeding, abdominal/thoracic surgery or trauma, delayed gastric emptying, older age, inadequate nursing staff, large size or diameter of feeding tube in children, malpositioning of the feeding tube, and transport.

In the perioperative period, factors that increase the likelihood of aspiration include:[8]

- A high urgency of surgery
- Difficult airways
- Inadequate depth of anesthesia
- Use of the lithotomy position
- Gastrointestinal problems such as delayed gastric emptying, gastroesophageal reflux, ileus, acute abdomen, or bowel obstruction[12]
- Depressed consciousness
- Increased severity of illness
- Obesity
- Use of drugs that reduce lower esophageal sphincter pressure or delay gastric emptying.
 - Certain drugs reduce the lower esophageal sphincter pressure and promote gastroesophageal reflux in anesthesia and disease states, and thereby increase the risk for aspiration. These drugs include atropine, glycopyrrolate, dopamine, nitroprusside, ganglion blockers, thiopental, tricyclic antidepressants, beta-adrenergic stimulants, halothane, opioids, and propofol.
 - Certain drugs delay gastric emptying. Opioids can significantly delay gastric emptying and promote ileus. Glucagon-like peptide-1 (GLP-1) receptor agonists and dual glucose-dependent insulinotropic polypeptide (GIP)/GLP-1 receptor agonists are known to delay gastric emptying, and because of retained gastric contents, their use is a risk factor for aspiration during procedures requiring general anesthesia or deep sedation.[32][33][34]

Pathophysiology

A higher prevalence of cerebrovascular and degenerative neurologic disease in people ages >70 years explains why increased age is a risk factor for aspiration. These conditions result in dysphagia and an impaired cough reflex, both of which increase the risk for aspiration of foreign bodies, food, liquids, and barium contrast.[4] [35]

Anticholinergics, antipsychotics, or anxiolytics may also impair the cough reflex and/or swallowing.

Tachypnea, resulting from various medical conditions, alters the coordination between deglutition and respiration, and increases the risk for aspiration.[36]

The lower and upper esophageal sphincter tone is decreased in GERD, by certain drugs (anticholinergics, antipsychotics, and anxiolytics), and by indwelling endotracheal, tracheostomy, gastric, or nasogastric tubes. This decrease in sphincter tone increases the risk for gastric content aspiration.[8]

The loss or impairment of protective laryngeal reflexes during the perioperative period contributes to the increased risk for anesthesia-related pulmonary aspiration. In addition to the obtunded reflexes from anesthesia and sedation, the supine position during diagnostic, surgical, and dental procedures aligns the trachea and the oropharynx, thereby increasing the risk for aspiration.[37]

Altered physiologic states such as pregnancy, gastrointestinal disorders, and diabetes mellitus are associated with delays in the rate of gastric emptying, which increase the gastric volume. Opioids can significantly delay gastric emptying and promote ileus. Glucagon-like peptide-1 (GLP-1) receptor agonists and dual glucose-dependent insulinotropic polypeptide (GIP)/GLP-1 receptor agonists have been shown to delay gastric emptying.[32] [33]

Pregnancy increases the risk of aspiration because progesterone decreases the tone of the lower esophageal sphincter and delays gastric emptying, in addition to the increase in intra-abdominal pressure secondary to the gravid uterus. Gastric juice aspiration and the resulting pneumonitis occurring in the context of pregnancy and anesthesia was first described by the eponymous Mendelson and is the basis for the wide practice of preoperative fasting.[38]

Aspiration of different gastric materials may have different effects. Low pH (acidic) material leads to neutrophil-predominant lung injury. There is apoptosis of type I alveolar epithelium caused by direct acid contact, and release of proinflammatory mediators secondary to activation of capsaicin-sensitive (TRPV-1) receptors. Alternatively, aspiration of small nonacidified gastric particles (SNAP) or bacteria directly stimulates alveolar macrophages to release innate immune/proinflammatory mediators through activation of polyanionic scavenger receptors or Toll-like receptors. Even without particulate or acid-related injury, gastric juice can contain digestive enzymes such as pepsin and bile salts that are caustic to the respiratory epithelium and likely to influence the type and severity of the respiratory presentation.[39] Most patients aspirate complex gastric contents, which are a combination of gastric food particles, bacterial products, cytokines, and acid, referred to as CASP (combined acid and small food particles). Aspiration of CASP may exacerbate damage due to the synergistic effect of acid and small gastric particles. The volume, viscosity, force of inhalation during aspiration, and host defenses contribute to the variable presentation that can manifest as bronchitis, bronchiolitis, pneumonitis/pneumonia, and ultimately acute respiratory distress syndrome.

Regardless of the type of aspiration material, there is an acute inflammation in the lung characterized by neutrophil infiltration, alveolar hemorrhage, intra-alveolar and interstitial edema, and impairment of alveolar fluid clearance. This is followed by a repair process characterized by scavenging of alveolar detritus by macrophages and proliferation of type II alveolar epithelial cells.[40]

Case history

Case history #1

A 62-year-old woman with systemic lupus erythematosus undergoes a head magnetic resonance imaging for acute mental status changes suggesting lupus cerebritis. The patient has been taking prednisone each day for several months. She has a diagnosis of GERD, for which she takes a proton-pump inhibitor. During the imaging study in the supine position, the patient vomits and aspirates gastric contents consisting of yellowish-greenish fluid. Severe respiratory distress and hypoxemia develop, and she requires endotracheal intubation, mechanical ventilation, and admission to the intensive care unit. Physical exam reveals bilateral crackles and wheezes.

Case history #2

An 80-year-old male with a past medical history of ischemic stroke, COPD, obstructive sleep apnea (noncompliant with noninvasive ventilation), and diabetes mellitus presents after being found by his partner with altered mental status. He had undergone a recent dental procedure and had been managing his pain with opioids. Physical exam reveals severe hypoxemia and the patient requires immediate intubation for airway protection. Right-sided crackles are identified on auscultation. Imaging reveals dense right lower lobe consolidation.

Other presentations

Aspiration is more common in older patients who have comorbid conditions that may lead to altered mental status. This population includes patients with swallowing dysfunction, disruption of the gastroesophageal junction, or anatomic abnormalities of the upper airway or digestive tract. Additionally, patients undergoing upper airway or endoscopic procedures are at risk. Any situation in which mental status is altered and airway protection mechanisms are lost increases the risk of aspiration. Limited protection is provided by a nasogastric tube, a percutaneous endoscopic gastrostomy, or an endotracheal or tracheostomy tube, and in fact they may increase the risk.^[2] Patients with aspiration of gastric contents can develop aspiration bronchiolitis, pneumonitis, pneumonia, and acute respiratory distress syndrome. Patients with neurologic deficits, infants, older patients, and debilitated patients with dysphagia may aspirate barium sulfate during radiologic procedures, which can result in severe pneumonitis and death.^[4]



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood



Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed

From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Approach

The diagnosis of acute aspiration is primarily by history, bearing in mind the presence of risk factors, with confirmation when necessary by imaging studies. At times it can be witnessed: for example, during intubation or in a patient who is vomiting and then aspirates. The signs, symptoms, and imaging studies can be indistinguishable from pneumonia and other causes of acute respiratory distress syndrome (ARDS), so the possibility of gastric aspiration must be considered in high-risk patients.

History, physical exam, and lab tests

Aspiration usually takes place in the presence of risk factors, typically in patients ages >70 years who have comorbid conditions that may alter mental status, or in patients with a depressed level of consciousness due to intoxication, illness, or drugs. A sudden onset of fever, cough paroxysms, wheeze, laryngospasm, or cyanosis in at-risk patients should prompt consideration for acute aspiration. Aspiration in the setting of general anesthesia or intoxication, or in the intensive care unit, is also common, but may not present with overt symptoms. It may be silent and may only be apparent from the consequences of aspiration such as bronchospasm, hypoxia, bronchiolitis/bronchitis, pneumonia, atelectasis, fever, and even respiratory failure from noncardiogenic pulmonary edema.[\[37\]](#) [\[41\]](#)

A complete blood count (CBC) and blood gas analysis can determine leukocytosis and hypoxia.

If pneumonia develops after the aspiration of gastric contents, hypoxia, leukocytosis, and other features of systemic inflammatory response may ensue, though these findings may not distinguish infection from more chemical pneumonitis. See Aspiration pneumonia .

Aspiration pneumonitis caused by barium sulfate should be suspected in patients who develop respiratory distress after upper gastrointestinal studies with barium. This is often directly visualized radiographically given the high-density material.



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood



Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed


From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Chest x-ray

When aspiration of gastric contents results in aspiration pneumonitis or pneumonia, the chest x-ray reveals patchy, bilateral airspace consolidations with a perihilar and basilar distribution.[48] However, an estimated 25% of cases may not be apparent on chest x-rays relative to computed tomography (CT) imaging.[68] The right lung may be involved more frequently because of the obtuse angle between the trachea and the right main bronchus. The most commonly involved segments are the superior and posterobasal segments of the right lower lobe and the posterior segment of the right upper lobe because of their dependent location in the supine position. If the patient is in a different position, other segments can be involved.

Chest x-ray findings of aspiration pneumonitis usually develop within 2 hours of aspiration and sometimes resolve quickly. However, if pneumonia develops, opacities can become apparent days later and can take weeks to resolve.

Barium aspiration should be confirmed by chest x-ray, which may reveal hyperdense opacities in the mid and lower lungs, sometimes outlining the tracheobronchial tree, and more distally, with a bronchiolocentric or centrilobular pattern.^{[70] [71] [72]}



A. Portable upright chest x-ray before aspiration; B. Chest x-ray 1 hour after aspiration, showing bilateral diffuse alveolar infiltrates, worse at the bases on the right side

From the collection of Dr Henri Colt




Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed

From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Chest CT

In cases of aspirated gastric content, chest CT may show opacities in the posterior segments of the upper lobes and the superior segments of the lower lobes. In advanced cases, the findings may be indistinguishable from those of ARDS. CT scanning precisely delineates the location of the lobar and segmental opacities. Aspiration of fat (exogenous lipoid pneumonia) or contrast material can sometimes be determined by measuring the tissue attenuation on CT scans. Lung abscess and empyema are potential complications of aspiration that are visualized better with CT than with plain chest x-ray.^[73] See Lung abscess and Empyema .

Although chest x-ray is sufficient in most cases of aspiration, a CT chest should be ordered if foreign body aspiration is suspected (to plan extraction) or if the patient fails to improve with initial therapy (to rule out empyema or lung abscess). Chest CT is superior compared with chest x-ray in the detection of aspiration pneumonia, and if initial chest x-ray is inconclusive, CT should be ordered.[68]



Lipoid pneumonia. A 77-year-old woman with dysphagia and achalasia following a stroke presented with recurrent lung infiltrates, including a persisting right middle lobe lesion. Hounsfield units (HU) measurement was -157, consistent with lipoid pneumonia. Comparative subcutaneous fat and aorta (blood/tissue) HU are shown

From the collection of Dr Augustine Lee, used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Bronchoscopy and bronchoalveolar lavage

Bronchoscopy can play both a diagnostic and therapeutic role. If the aspirated material is particulate or potentially obstructive, or if there is radiographic evidence of lobar or segmental collapse, the airway can be cleared and further evolution to ARDS may be avoided. In cases of aspiration of barium sulfate,

especially if a patient has hypoxia or respiratory distress, early bronchoscopy may be helpful, but incurs some risk of disseminating the contrast to unaffected airways.[24] [74]



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood

Another occasional therapeutic role besides airway and lung lavage is the potential placement of tracheal or bronchial stents in cases of tracheoesophageal fistula, as well as the placement of percutaneous tracheostomy for those with severe dysphagia.

Aside from potential therapeutic roles, bronchoscopy is also important diagnostically. Bronchoalveolar lavage techniques or protected brush specimens can be used to identify, confirm, or refute the presence of an infection, which will have direct impact on the patient's antibiotic management.[2] In addition to infection, bronchoalveolar lavage can diagnose acute eosinophilic pneumonia and diffuse alveolar hemorrhage, both of which can present similarly to acute aspiration with acute respiratory distress and lung infiltrates.[75] Occasionally, cytologic analysis can pick up unusual tumors that can appear pneumonic on imaging. When the differential is broader, a bronchoscopic biopsy may occasionally be employed for interstitial pneumonias and other considerations, though not often in the acute setting.[75]

Finally, a careful bronchoscopic exam may be important to identify a tracheoesophageal fistula that may be complicating cancer, surgery, radiation, or trauma.

Biomarkers of aspiration syndromes

Multiple biomarkers are used in the research setting and at some centers. These have included the evaluation of lipid laden macrophages, pepsin, amylase, bile, and cytokines directly from lower airway samples.[40] [76] The sensitivity and specificity of these markers remain unclear, and they lack clinical validation.[40]

Tests to identify need for antibiotics

Tests to identify a bacterial component to the pneumonic presentation depend on the clinical scenario and may include culture and sensitivities of samples obtained from bronchoalveolar lavage and brushing, venepuncture, or thoracentesis. See Aspiration pneumonia , Hospital-acquired pneumonia , Community-acquired pneumonia , Pleural effusion , Empyema , Lung abscess , and Sepsis in adults .

History and exam

Key diagnostic factors

acute or chronic cough (common)

- Seen in 36% of patients with aspiration of gastric contents.[77] However, this presumes that the patient's cough reflex is intact. Those with absent or diminished cough reflex (e.g., due to opioid intoxication or respiratory muscle weakness related to amyotrophic lateral sclerosis or critical illness myopathy) may have more consequential aspiration.[18] [42]

fever (common)

- Extremely common after aspiration of gastric contents, occurring in 94% of cases.[77]

dyspnea (common)

- Extremely common after aspiration of gastric contents, seen in 78% of cases.[77] In severe cases, respiratory failure may result from noncardiogenic pulmonary edema.

wheezing (common)

- Occurs in 32% of patients after aspiration of gastric contents.[77]

crackles (common)

- Presence on lung auscultation is common after aspiration of gastric contents, occurring in 72% of cases.[77]

Other diagnostic factors

laryngospasm (uncommon)

- May be induced by laryngopharyngeal reflux, and serves as a protective mechanism against aspiration, though may lead to other dire consequences. In one series about 8% of patients with confirmed aspiration also had laryngospasm.[78]

Risk factors

Strong

decreased level of consciousness (Glasgow coma scale score <9)

- Associated with trauma, use of sedatives, alcohol, general anesthesia, or neurologic disorders (e.g., brain tumor, seizure, Parkinson disease, mental retardation, stroke). Impairs the protective airway mechanisms and increases the risk for aspiration of foreign objects.[\[20\]](#) [\[35\]](#)

increased severity of illness

- Critical illness and higher anesthesia risk classification (defined as class III, IV, or V in the American Society for Anesthesiology risk classification system, which is based on physical status) indicates higher risk of aspiration.[\[5\]](#) [\[41\]](#) [\[42\]](#)

general anesthesia

- Decreased level of consciousness results in loss of protective reflexes. Anesthetic drugs may decrease lower and upper esophageal sphincter tones, and promote ileus and gastroparesis, increasing risk for gastrolaryngeal reflux. Supine positioning and unsecured airways may add to the increased risk for reflux and subsequent aspiration.[\[8\]](#)

age >70 years

- Risk of aspiration is higher in older patients, especially during and after the seventh decade, probably because of higher prevalence of aging-associated degenerative neurologic and cerebrovascular disorders that can cause dysphagia and/or impaired cough reflex.[\[27\]](#) [\[35\]](#)[\[41\]](#) Over 50% of patients with acute food asphyxiation are 71-90 years of age.[\[43\]](#)

head trauma

- Can result in altered mental status and impaired airway reflexes. Also increases risk for gastric content aspiration because the elevated intracranial pressure that may result from head trauma delays gastric emptying.[\[44\]](#)

cerebrovascular disease

- Over one third of patients with acute stroke have aspiration documented on radiologic studies.[\[27\]](#) Abnormal swallowing increases risk for aspiration.[\[27\]](#) Patients with dysphagia have delayed triggering of the pharyngeal motor response and decreased laryngeal elevation, resulting in poor coordination and timing of oral, pharyngeal, and laryngeal events during swallowing.[\[45\]](#)

endotracheal or tracheostomy tube

- Indwelling endotracheal or tracheostomy tube raises risk for aspiration. Cuffed tubes are not completely protective and mechanically interrupt glottic closure or the lower esophageal sphincter.[\[41\]](#) [\[46\]](#) [\[47\]](#)[\[48\]](#) Endotracheal intubation that lasts >8 hours can result in laryngotracheal complications, which increase the risk for aspiration after extubation.[\[49\]](#)
- Aspiration has been documented in up to 45% of critically ill trauma patients after extubation. The mechanisms are multiple and include changes in glottis anatomy caused by vocal cord ulceration and laryngeal edema, and disruption of the swallowing reflex caused by muscle atrophy, incoordination, and diminished sensory abilities of the larynx.[\[50\]](#) Traumatic intubations (blood in the endotracheal tube, esophageal intubation, and multiple intubation attempts) increase the risk for aspiration after

extubation. Patients are often sedated and critical illness further impairs swallow and cough reflexes. Muscle weakness associated with critical illness can further diminish effective cough that might otherwise offset aspiration events.[42]

dysphagia

- Abnormal swallowing increases risk for aspiration.[27] Patients with dysphagia have delayed triggering of the pharyngeal motor response and decreased laryngeal elevation, resulting in poor coordination and timing of oral, pharyngeal, and laryngeal events during swallowing.[45]
- High-risk conditions resulting in oropharyngeal dysphagia include neurologic disorders (cerebrovascular disease, head trauma, closed head injury, cervical spine injury, anoxia, seizure disorder, vocal cord paralysis, Parkinson disease, amyotrophic lateral sclerosis, Alzheimer disease), certain surgeries (for head and neck cancer, anterior and posterior cervical spine surgery, brain surgery, coronary artery bypass grafting, esophagogastrectomy), structural abnormalities (oropharyngeal tumors, tracheoesophageal fistula), gastrointestinal disorders (laryngopharyngeal reflux), tracheostomy, and adverse effects from sedatives and antipsychotics.

airway difficulties

- Difficult intubation and laryngospasm are reported risk factors for aspiration during the perioperative period.[5] [7]

barium meal

- Can be aspirated during a barium radiographic exam.



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood



Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed
From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Weak

male sex

- Risk for aspiration in males is almost twice that in females, probably because of the higher rates of neurologic and cardiovascular disorders in males.

gastroesophageal reflux disease

- Independent risk factor for aspiration in critically ill patients and during perioperative period in patients undergoing general anesthesia.[8] [20] Conditions that increase the risk for gastroesophageal reflux

include esophageal disease (collagen vascular disease, cancer, achalasia, dysmotility), hiatal hernia, peptic ulcer disease, gastritis, bowel obstruction, ileus, and elevated intracranial pressure.[41]

feeding tubes

- Nasoenteric feeding tubes may stimulate the pharynx and increase risk for aspiration by decreasing lower esophageal sphincter tone.[51] Gastric placement of feeding tubes increases risk for aspiration, and postpyloric placement of tube may have a protective effect.[20] Large feeding tubes increase risk, especially in infants and young children. It is unclear whether gastrostomy feeding tubes have lower risk for aspiration than nasogastric feeding.[44]

supine position

- Supine positioning during dental, medical, or radiologic procedures, transport, and general anesthesia aligns the trachea and oropharynx, and facilitates gravitational flow of gastric content in the oropharynx. Also associated with difficulty swallowing. Sedation and anesthesia enhance risk of aspiration by obtunding the protective reflexes.[37]

delayed gastric emptying

- High gastric residual volume is common in critically ill patients who receive enteral nutrition. This increases risk for vomiting or regurgitation of feeds and makes patients prone to aspiration.[44] Associated factors include abdominal surgery, sepsis, metabolic abnormalities (hyperglycemia, diabetes mellitus, hypokalemia), renal failure, increased intracranial pressure, and some drugs (e.g., opioids).
- Glucagon-like peptide-1 (GLP-1) receptor agonists and dual glucose-dependent insulinotropic polypeptide (GIP)/GLP-1 receptor agonists are known to delay gastric emptying, and because of retained gastric contents, their use is a risk factor for aspiration during procedures requiring general anesthesia or deep sedation.[32] [33][34]

obesity


- Independent risk factor for aspiration related to general anesthesia.[52] This association may be partly influenced by the presence of obstructive sleep apnea.[53]

drugs that reduce esophageal sphincter tone

- Drugs used in anesthesia and in the intensive care unit may reduce the lower and upper esophageal sphincter tone and increase risk for aspiration. Drugs include atropine, glycopyrrolate, dopamine, nitroprusside, ganglion blockers, thiopental, beta-adrenergic stimulants, halothane, and propofol. Tricyclic antidepressants also have this effect.[8]

Tests

1st test to order

Test	Result
chest x-ray - In pneumonitis or pneumonia resulting from aspirated gastric contents, there are patchy, bilateral airspace consolidations with a perihilar and basilar distribution.[48] The most commonly involved areas are the superior and posterobasal segments of the right lower lobe and the posterior segment of the right upper lobe because of their dependent location in the supine position. If the patient is in a different position, other segments can be involved. - X-ray changes are usually apparent within 2 hours of aspiration but if pneumonia develops, opacities can become apparent days later and take weeks to resolve. See Aspiration pneumonia . - Barium aspiration should be confirmed by CXR.[70] [71] [72]  <i>A. Portable upright chest x-ray before aspiration; B. Chest x-ray 1 hour after aspiration, showing bilateral diffuse alveolar infiltrates, worse at the bases on the right side</i> <i>From the collection of Dr Henri Colt</i>	patchy, bilateral airspace consolidations; barium may be identified by hyperdense opacities in the mid and lower lungs, sometimes outlining the tracheobronchial tree, and more distally, with a bronchiolocentric or centrilobular pattern

Test

Result





Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed

From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

- An estimated 25% of cases may not be apparent on CXRs relative to CT imaging.[68]

Other tests to consider

Test	Result
chest CT - In cases of aspirated gastric contents, opacities may be seen in dependent segments of the lung. Chest CT scanning precisely delineates the location of the lobar and segmental opacities. In advanced cases, the findings may be indistinguishable from those of acute respiratory distress syndrome. See Acute respiratory distress syndrome . - Aspiration of fat (exogenous lipid pneumonia) or contrast material can sometimes be determined by measuring the tissue attenuation on CT scans. Lung abscess and empyema are potential complications of aspiration that are visualized better with CT than with plain chest x-ray.[73] See Lung abscess and Empyema . - Although CXR is sufficient in most cases of aspiration, a CT chest should be ordered if foreign body aspiration is suspected (to plan extraction) or if the patient fails to improve with initial therapy (to rule out empyema or lung abscess). - Chest CT is superior compared with CXR in the detection of aspiration pneumonia, and if initial CXR is inconclusive, CT should be ordered.[68]	opacities in dependent segments
 <i>Lipoid pneumonia. A 77-year-old woman with dysphagia and achalasia following a stroke presented with recurrent lung infiltrates, including a persisting right middle lobe lesion. Hounsfield units (HU) measurement was -157, consistent with lipoid pneumonia. Comparative subcutaneous fat and aorta (blood/tissue) HU are shown</i> <i>From the collection of Dr Augustine Lee, used with permission of Mayo Foundation for Medical Education and Research, all rights reserved</i>	

Test	Result
bronchoscopy with bronchoalveolar lavage - Bronchoscopy is indicated if aspirated material is particulate or potentially obstructive, or if there is radiographic evidence of lobar or segmental collapse, in order to clear the airway. In addition, bronchoscopy can be used to collect quantitative cultures on bronchoalveolar lavage or protected specimen brush, which can be used to guide antibiotic therapy, particularly in patients who fail to respond to empiric antibiotic treatment. - Cytologic analysis can pick up unusual tumors that can appear pneumonic on imaging. When the differential is broader, a bronchoscopic biopsy may occasionally be employed for interstitial pneumonias and other considerations, though not often in the acute setting.[75] Finally, a careful bronchoscopic exam may be important to identify a tracheoesophageal fistula that may be complicating cancer, surgery, radiation, or trauma. - In cases of aspiration of barium sulfate, especially if a patient has hypoxia or respiratory distress, early bronchoscopy should be performed to remove the barium from the airway, but there is some risk of disseminating the contrast to unaffected airways. Deaths have been reported in patients who aspirated barium and for whom bronchoscopy was not performed for airway clearance.[24] [74]  <i>Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study</i> From the collection of Dr Kamran Mahmood	airway erythema; particulate matter, barium may be seen; protected brush specimens can be used to identify, confirm, or refute the presence of an infection; eosinophils or tumor cells may be present
CBC Performed if pneumonia develops after the aspiration of gastric contents.	leukocytosis
arterial blood gases Aspiration may cause hypoxia.	reduced oxygen tension
blood culture In the context of small bowel obstruction, where there is a high risk of bacterial translocation, bacterial contamination of gastric juice, and sepsis, the initiation of antibiotics should be considered up front and	growth of causative bacterial species

Test	Result
should be adjusted when pathogens are identified.[2] [79] See Small bowel obstruction and Sepsis in adults .	
thoracentesis • Thoracentesis for microscopy, culture, and sensitivity should be considered if there is concern for a complicated parapneumonic effusion. See Pleural effusion and Empyema .	growth of causative bacterial species

Differentials

Condition	Differentiating signs / symptoms	Differentiating tests
Acute respiratory distress syndrome	Seen in the setting of sepsis, chemical exposure, drug reactions, toxic inhalation, or trauma. No difference in signs and symptoms from aspiration pneumonitis, which can itself lead to acute respiratory distress syndrome.^[80]	- Bilateral alveolar infiltrates can be indistinguishable from those seen in aspiration pneumonitis. - Aspiration pneumonitis is usually distinguished from aspiration pneumonia on the history of a witnessed large-volume aspiration. In aspiration pneumonitis, the patient often has a reduced level of consciousness.
Asthma exacerbation	Wheezing is paroxysmal and intermittent and usually diffuse. It decreases after bronchodilators, is polyphonic, and is characterized by many different pitches. Cough is triggered by exercise, cold, sleep, and allergens. Patient may have history and/or family history of atopy or asthma.^[81]	Pulmonary function tests usually show reversible obstructive ventilatory impairment.^[81]
Cystic fibrosis with exacerbation	Wheezing presents early in life, with accompanying poor weight gain, diarrhea, and recurrent sinus and pulmonary infections. Nasal exam may reveal polyps. Cough is productive and wet, suggesting a suppurative process such as bronchiectasis. Patient may have family history of bronchiectasis.^[81]	Sweat chloride test shows elevated level of chloride. Chest x-ray may reveal bronchiectasis, but CT scan is more sensitive.^[82]
COPD exacerbation	Wheezing is diffuse and associated with increased mucus production and history of progressive dyspnea.	Chest x-ray shows peribronchial cuffing and hyperinflation.
Community-acquired pneumonia	No difference in signs and symptoms.	Chest x-ray may show lobar consolidation.
Hospital-acquired pneumonia	No difference in signs and symptoms.	Chest x-ray may show lobar consolidation.
Congestive heart failure	Orthopnea, paroxysmal nocturnal dyspnea, and right	Chest x-ray may reveal enlarged pulmonary

Condition	Differentiating signs / symptoms	Differentiating tests
	upper quadrant discomfort may be features. Sputum, if present, is usually frothy. Fever is usually absent.[83] The third heart sound gallop rhythm has up to 50% sensitivity but 90% specificity. Pulsus alternans, characterized by evenly spaced strong and weak peripheral pulses, is pathognomonic of severe left ventricular failure.[83] [84]	vessels, cardiomegaly, and pulmonary edema; the infiltrates improve quickly after diuresis. Brain natriuretic peptide >100 picograms/mL makes a diagnosis of heart failure with sensitivity of 90%, specificity of 76%, and predictive accuracy of 83%.[85]
Acute exacerbation of interstitial lung disease (ILD)	No differentiating symptoms or signs, but they develop over a few days to several weeks. In some cases it is idiopathic (Hamman-Rich syndrome) or related to other ILD including idiopathic pulmonary fibrosis.[75] Some connective tissue disease-ILDs such as systemic sclerosis are prone to esophageal dysmotility and consequent reflux and aspiration. Drug and radiation toxicity should also be considered in the appropriate clinical context.	- Bilateral alveolar infiltrates can be indistinguishable from those seen in aspiration pneumonitis or acute respiratory distress syndrome (ARDS). However, evidence of background fibrotic lung disease such as honeycombing may be present. Comparison to prior imaging when available is critical. - Bronchoalveolar lavage cell count reveals >10% neutrophils, and lung biopsy shows diffuse alveolar damage.[75]
Acute bronchiolitis obliterans organizing pneumonia	No differentiating symptoms or signs. Can be idiopathic or due to collagen vascular disorders, drugs, radiation, or infection.[75]	Bilateral alveolar infiltrates can be indistinguishable from those seen in aspiration pneumonitis. Bronchoalveolar lavage cell count in organizing pneumonia may reveal neutrophilia and sometimes lymphocytosis (but <25% lymphocytes) with eosinophilia (but <25% eosinophils).[75]
Acute eosinophilic pneumonia	No differentiating symptoms or signs, but duration of illness is usually <1 week. Can be idiopathic or caused by drugs.[75]	Bilateral alveolar infiltrates can be indistinguishable from those seen in aspiration pneumonitis. Bronchoalveolar lavage cell count reveals >25% eosinophils. Eosinophilic pleural effusion is rare. Eosinophilic infiltration and

Condition	Differentiating signs / symptoms	Differentiating tests
		diffuse alveolar damage may be seen on lung biopsy.[86]
Acute hypersensitivity pneumonitis	No differentiating symptoms or signs, but usually develops within 4-6 hours after inhalation of an organic agent. Caused by environmental and work-related antigens.[75]	Bilateral alveolar infiltrates can be indistinguishable from those of aspiration pneumonitis. Granulomatous and cellular pneumonitis with diffuse alveolar damage may be seen on lung biopsy. Bronchoalveolar lavage cell count shows lymphocytosis (>25%) and sometimes neutrophilia (<10%).[75]
Diffuse alveolar hemorrhage	Hemoptysis is absent in 33% of patients. Causes include vasculitis, collagen vascular disorders, ant basement membrane antibody disease, coagulopathies, antiphospholipid antibody syndrome, and diffuse infections.[75]	- Bilateral alveolar infiltrates can be indistinguishable from those of aspiration pneumonitis. Pulmonary capillaritis, bland hemorrhage, and diffuse alveolar damage seen on lung biopsy. - Bronchoalveolar lavage shows progressively bloodier return. Cytology shows red blood cells and hemosiderin-laden macrophages. - Urinalysis may show proteinuria, hematuria, and red cell casts in cases of pulmonary-renal syndromes.[75]
Neurogenic pulmonary edema	No differentiating symptoms or signs. Usually develops within minutes to hours after acute central nervous system injury such as seizures, head injury, or cerebral hemorrhage. Resolves within 48-72 hours.[87]	Bilateral alveolar infiltrates may be indistinguishable from those seen in aspiration pneumonitis.[87]

Screening

The most commonly used diagnostic test for aspiration is the bedside swallow evaluation. This test includes a patient interview, a physical exam, and the assessment for signs of aspiration. While this is usually performed by a speech-language pathologist, to streamline the diagnostic evaluation some of the more sensitive components of the bedside swallow evaluation can be performed individually or in combination as screening tests.[88] These screening tests include a patient's attempts to swallow small quantities of water or ice chips while being observed by a nurse, a speech-language pathologist, or a physician. Screening tests should be considered whenever aspiration is suspected based on the presence of risk factors. Furthermore, all patients suffering from an acute stroke should undergo a bedside swallow evaluation by a speech-language pathologist, largely because between 40% and 78% of stroke patients have dysphagia.[16] [68]

Dysphagia program studies applied as a result of these screening methods could substantially reduce the rates of pneumonia in stroke patients.[89]

Bedside swallow evaluation

Patients thought to have any risk factors for aspiration should undergo a complete neurologic evaluation before feeding.[90] This includes assessing cortical functions, bulbar muscles, gag reflex, and cough reflex. Alert patients with cough who are at high risk of aspiration should be observed drinking small amounts of water. If the patient coughs or shows signs of aspiration, the patient should be referred for a detailed swallowing evaluation.[27] Early bedside swallow screening (within 1 day of admission and before any oral intake) and dysphagia management in patients with acute stroke reduces the risk of aspiration pneumonia, may be cost-effective, and may assure quality care with optimal outcome.[91] If doubt exists or if silent aspiration is suspected, imaging studies should be performed.[16] Phonetic assessment independent of and during the swallow provocation tests may be a useful adjunct to bedside tests for aspiration risk, but it is not validated.[88] [92] Although a multitude of other screening tests are available, they have variable sensitivity.[93] In the opinion of the authors, however, there may be a benefit to obtaining an assessment of aspiration risk from more than one test.

Flexible endoscopic evaluation of swallowing

Flexible endoscopic evaluation of swallowing (FEES) can be performed by speech pathologists at the bedside. With the use of a flexible fiberoptic scope, direct evidence of aspiration is documented by detecting the presence of food or a thick liquid over the vocal cords. The test also assesses the vocal cord function. In a study of critically ill trauma patients, FEES performed within 24 hours after extubation showed aspiration in 45% of the patients, of whom almost half were silent aspirators. These patients resumed an oral diet at a mean of 5 days after extubation and had no pulmonary complications.[50] FEES avoids exposure to radiation; however, modified barium swallow is noninvasive and more widely available.[56][94]

Videofluoroscopic swallowing study and modified barium swallow

For these tests, the patient swallows barium under fluoroscopic imaging. Passage or retention of the radiopaque material in the respiratory tract is seen.[16] On videofluoroscopic swallowing study, 38% of patients with acute stroke had overt aspiration and 67% had silent aspiration.[95] Modified barium swallow performed in patients with indwelling tracheostomies on positive-pressure ventilation showed an incidence of aspiration of 50%, and 77% of these patients were silent aspirators.[46]

Emerging assessments for dysphagia

Interventions targeting improved respiratory sensation and respiratory mechanics may prove to be beneficial in patients (such as those with Parkinson disease) who have impaired airway protection.[96] Various tests of respiratory mechanics have been employed to determine aspiration risk or at least the ability to compensate for a respiratory event. For example, a reduction in the peak cough flow was an independent risk factor for pneumonia.[97] Phonetic screening is a novel technique that may be used to identify dysphagia and aspiration risk at the bedside.[92]

Approach

Patients that aspirate will often have a reduced level of consciousness and require close observation for at least 48 hours, either in the hospital or in another care facility. Although some patients may have dramatic signs and symptoms, many are asymptomatic, and infectious complications and acute respiratory distress syndrome (ARDS) can be delayed.[2] A high index of suspicion and prompt action are required if optimum outcomes are to be obtained.

Aspiration pneumonitis

Aspiration of gastric contents

Patients with recent suspected or witnessed aspiration of gastric contents should, whenever possible, be immediately placed semi-prone and tilted to a 30° head-down position. This positions the larynx at a higher level than the oropharynx and allows the gastric content to drain externally. The oropharynx should be gently suctioned, taking care to avoid initiating a gag reflex that may worsen aspiration.[2]

Once the oropharynx has been suctioned, the airway should be secured by endotracheal intubation after rapid sequence induction with cricoid pressure if the patient is deemed to be at risk of further aspiration, is unable to protect their own airway (regurgitation, poor cough reflex), or shows signs of respiratory failure (tachypnea, dyspnea, confusion, cyanosis).[62] [63] Once the airway is secured, a nasogastric tube should be inserted to empty the stomach, and where possible the patient can be tilted to a 45° head-up position to help prevent further aspiration.

If a substantial amount of gastric content (>20-25 mL in an adult) is likely to have been aspirated, prompt (within a few hours) bronchoscopy and suctioning can remove aspirated gastric fluid and solid material from the central airways, thereby helping reduce inflammatory reaction, prevent lung collapse, and lessen the risk of subsequent infection.[98] A volume of gastric aspirate >0.3 mL per kilogram of body weight (i.e., 20-25 mL in adults) with a pH <2.5 is believed necessary for the development of aspiration pneumonitis, although aspiration of particulate food matter can cause severe pulmonary damage, even if the pH of the aspirate is above 2.5.[2] [99] Animal studies have shown a biphasic pattern to injury, with an initial peak at 1-2 hours after aspiration (direct burn effects) and a second peak at 4-6 hours (related to neutrophil infiltration).[1] [2]

Gastric aspirate is sterile under normal conditions due to the low pH, so bacterial infection does not have an important role in the early stages of acute lung injury.[2] Consequently, immediate routine antibiotic therapy is not recommended for aspiration pneumonitis.[2] Even if a patient has fever, leukocytosis, or pulmonary infiltrates, immediate use of antibiotics may be unnecessary and may select resistant organisms in an uncomplicated case of chemical pneumonitis.

The empiric use of antibiotics should, however, be considered depending on the clinical situation (e.g., severe respiratory involvement; severe preexisting lung pathology such as advanced COPD or interstitial lung disease; immunocompromise, including corticosteroid or immunomodulator use, post-transplant immunosuppression, chemotherapy, neutropenia; poor dentition; or the development of lung abscess, necrotizing pneumonia, or empyema).[100] In the context of small bowel obstruction, where there is a high risk of bacterial translocation, bacterial contamination of gastric juice, and sepsis, the initiation of antibiotics should be considered up front and should be adjusted when pathogens are identified.[2] See Small bowel obstruction and Sepsis in adults .

In practice, antibiotics should be initiated based on clinical concern and if the patient is not responding to supportive care. Local microbiology, resistance patterns, and risks for anaerobic infection should be considered when choosing the antibiotic regimen. Culture and sensitivities of samples obtained via bronchoscopy with bronchoalveolar lavage or protected brush specimens, thoracentesis, or blood taken for culture should guide antimicrobial management.[101] Empiric therapy with broad-spectrum antibiotics is recommended if the pneumonitis does not resolve 48 hours after aspiration.[2] [102] Antibiotics should be stopped if concern for infection drops and narrowed quickly if a specific pathogen is identified. See Aspiration pneumonia , Community-acquired pneumonia , and Hospital-acquired pneumonia .


Computed tomography (CT) imaging can identify related pathologies indicating a need for antibiotic therapy (e.g., empyema, abscess, necrosis). See Lung abscess , Empyema , Pleural effusion , and Sepsis in adults .

Positive-pressure ventilation with positive end-expiratory pressure can be used in patients who are intubated for airway protection or respiratory failure. However, positive pressure ventilation without a secure endotracheal tube is generally contraindicated in patients with known or suspected aspiration causing respiratory failure. Positive pressure helps to prevent atelectasis and improve the ventilation-perfusion ratio in patients who have aspirated gastric content.[37] However, it is essential that endotracheal suctioning, and if needed, post-intubation bronchoscopy, is performed before positive-pressure ventilation is employed, to avoid forcing aspirated material deeper into the lungs.[35] [37] Mechanical ventilatory support should follow lung-protective strategies as per ARDS guidelines. See Acute respiratory distress syndrome .

Respiratory cultures should be obtained from patients on mechanical ventilation, and antibiotics should be initiated immediately because of a high risk of developing ventilator-associated pneumonia.[1] See Hospital-acquired pneumonia .

Aspiration of barium

This usually occurs in the context of upper gastrointestinal radiologic studies.



Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood

Bronchoscopy with suctioning, if considered, should be done immediately post-aspiration with the aim to remove any residual barium from the proximal airway and to reduce hypoxemia. Care should be taken to minimize potential spread of barium to unaffected areas from washing or lavage techniques. Owing to the inert nature of barium sulfate, usually no severe long-term harm is to be expected; however, severe pneumonitis and death has been reported in infants, and in older or debilitated patients.[4] [24] Infants, older adult patients, and those of any age with significant symptoms, should be admitted for observation if not already hospitalized.

There is no evidence for routinely prescribing antibiotics after barium aspiration, but antibiotics for possible pneumonia are usually considered if infiltrates do not resolve 48 hours post-aspiration.[70] See Aspiration pneumonia .

Corticosteroids

Human studies have shown no improvement in mortality, and the rate of gram-negative pneumonia 5 days after aspiration was higher in patients receiving corticosteroids.[103] Although the infiltrates improve more quickly in the patients given corticosteroids than in those given placebo, patients given corticosteroids might have a longer intensive care unit (ICU) stay.[104] [105]

Overall, because of the increased risk for gram-negative bacterial pneumonia and a prolonged stay in the ICU, together with the lack of any mortality benefit, corticosteroids are not indicated in the initial or empiric

management of aspiration pneumonitis. They are also not indicated with aspiration pneumonitis that is complicated by acute respiratory distress syndrome.[2] However, corticosteroids may be considered if other indications develop such as in select patients with severe ARDS.[106] [107] See Acute respiratory distress syndrome .

Treatment algorithm overview

Please note that formulations/routes and doses may differ between drug names and brands, drug formularies, or locations. Treatment recommendations are specific to patient groups: [see disclaimer](#)

Acute (summary)	
pneumonitis due to aspiration of gastric contents	
1st	immediate positional drainage
plus	oropharyngeal suctioning
adjunct	endotracheal intubation + nasogastric tube
adjunct	bronchoscopy + endotracheal suctioning
adjunct	antibiotics
adjunct	positive-pressure ventilation
pneumonitis due to aspiration of barium	
1st	immediate positional drainage
plus	bronchoscopy + endotracheal suctioning

Treatment algorithm

Please note that formulations/routes and doses may differ between drug names and brands, drug formularies, or locations. Treatment recommendations are specific to patient groups: [see disclaimer](#)

Acute

pneumonitis due to aspiration of gastric contents

1st immediate positional drainage

» Patients with recent suspected or witnessed aspiration of gastric contents should, whenever possible, be immediately placed semi-prone and tilted to a 30° head-down position. This positions the larynx at a higher level than the oropharynx and allows the gastric content to drain externally.

plus oropharyngeal suctioning

Treatment recommended for ALL patients in selected patient group

» The oropharynx should be suctioned gently, taking care to avoid initiating a gag reflex that may worsen aspiration.[2]

adjunct endotracheal intubation + nasogastric tube

Treatment recommended for SOME patients in selected patient group

» Once the oropharynx has been suctioned, if the patient is deemed to be at risk of further aspiration, is unable to protect their own airway (regurgitation, poor cough reflex), or shows signs of respiratory failure (tachypnea, dyspnea, confusion, cyanosis), their airway should be secured by endotracheal intubation via rapid sequence induction with cricoid pressure.[62] [63] Once the airway is secured, a nasogastric tube should be inserted to empty the stomach, and where possible the patient can be tilted to a 45° head-up position to help prevent further aspiration.

adjunct bronchoscopy + endotracheal suctioning

Treatment recommended for SOME patients in selected patient group

» If a substantial amount of gastric content (>20-25 mL in an adult) is likely to have been aspirated, prompt (within a few hours) bronchoscopy and suctioning can remove aspirated gastric fluid and solid material from the central airways, thereby helping reduce inflammatory reaction, prevent lung collapse, and lessen risk of subsequent infection.[98]

Acute

» A volume of gastric aspirate >0.3 mL per kilogram of body weight (i.e., 20-25 mL in adults) with a pH <2.5 is believed necessary for the development of aspiration pneumonitis, although aspiration of particulate food matter can cause severe pulmonary damage, even if the pH of the aspirate is above 2.5.[2] [99] Animal studies have shown a biphasic pattern to injury, with an initial peak at 1-2 hours after aspiration (direct burn effects) and a second peak at 4-6 hours (related to neutrophil infiltration).[1] [2]

adjunct antibiotics

Treatment recommended for SOME patients in selected patient group

» Gastric aspirate is sterile under normal conditions due to the low pH, so bacterial infection does not have an important role in the early stages of acute lung injury in most cases.[2] Consequently, immediate routine antibiotic therapy is not recommended for aspiration pneumonitis.[2] Even if a patient has fever, leukocytosis, or pulmonary infiltrates, immediate use of antibiotics may be unnecessary as this may select resistant organisms in an uncomplicated case of chemical pneumonitis.

» The empiric use of antibiotics should, however, be considered depending on the clinical situation (e.g., severe respiratory involvement; severe preexisting lung pathology such as advanced COPD or interstitial lung disease; immunocompromise, including corticosteroid or immunomodulator use, post-transplant immunosuppression, chemotherapy, neutropenia; poor dentition; or the development of lung abscess, necrotizing pneumonia, or empyema).[100] In practice, antibiotics should be initiated based on clinical concern and if the patient is not responding to supportive care.

» Culture and sensitivities of samples obtained via bronchoscopy with bronchoalveolar lavage or protected brush specimens, thoracentesis, or blood taken for culture should guide antimicrobial management.[101] Empiric therapy with broad-spectrum antibiotics is recommended if the pneumonitis does not resolve 48 hours after aspiration.[2] [102] Antibiotics should be stopped if concern for infection drops and narrowed quickly if a specific pathogen is identified. See Aspiration pneumonia , Community-acquired pneumonia , and Hospital-acquired pneumonia .

» Computed tomography (CT) imaging can identify related pathologies indicating a need

Acute

for antibiotic therapy (e.g., empyema, abscess, necrosis). See Lung abscess , Empyema , Pleural effusion , and Sepsis in adults .

» In the context of small bowel obstruction, where there is a high risk of bacterial translocation, bacterial contamination of gastric juice, and sepsis, the initiation of antibiotics should be considered up front.[2] See Small bowel obstruction and Sepsis in adults .

» Local microbiology, resistance patterns, and risks for anaerobic infection should be considered when choosing the antibiotic regimen.

adjunct positive-pressure ventilation

Treatment recommended for SOME patients in selected patient group

» Positive-pressure ventilation with positive end-expiratory pressure can be used in patients who are intubated for airway protection or respiratory failure. However, positive pressure ventilation without a secure endotracheal tube should generally be considered contraindicated in patients with known or suspected aspiration causing respiratory failure. Positive pressure helps to prevent atelectasis and improve the ventilation-perfusion ratio in patients who have aspirated gastric content.[37] However, it is essential that endotracheal suctioning, and if needed, post-intubation bronchoscopy, is performed before positive-pressure ventilation is employed, to avoid forcing aspirated material deeper into the lungs.[35] [37]

» Mechanical ventilatory support should follow lung-protective strategies as per acute respiratory distress syndrome guidelines. See Acute respiratory distress syndrome .

» Respiratory cultures should be obtained from patients on mechanical ventilation, and antibiotics should be initiated immediately because of a high risk of developing ventilator-associated pneumonia.[1] See Hospital-acquired pneumonia .

pneumonitis due to aspiration of barium**1st immediate positional drainage**

» Patients with recent suspected or witnessed aspiration of barium should, whenever possible, be immediately placed semi-prone and tilted to a 30° head-down position. This positions the

Acute

plus

larynx at a higher level than the oropharynx and allows the gastric content to drain externally.

bronchoscopy + endotracheal suctioning

Treatment recommended for ALL patients in selected patient group

» Bronchoscopy with suctioning, if considered, should be performed immediately post-aspiration to remove any residual barium from the airway and reduce hypoxemia. Care should be taken to minimize potential spread of barium to unaffected areas from washing or lavage techniques.

» Infants, older adult patients, and those of any age with significant symptoms should be admitted for observation if not already hospitalized.

» There is no evidence for routinely prescribing antibiotics after barium aspiration, but antibiotics for possible pneumonia are usually considered if infiltrates do not resolve 48 hours after aspiration.[70]

» See Aspiration pneumonia .

Primary prevention

Patients with comorbidities or neurologic disorders, or in critical care

Measures to identify gastroesophageal reflux disease, oropharyngeal dysphagia, gastrointestinal motility, and metabolic disorders (including diabetes mellitus) and to reduce oropharyngeal load of pathogenic organisms, particularly in patients with neurologic disease, patients with comorbidities, and those who are intubated, can potentially help to reduce aspiration and associated complications such as aspiration pneumonia.[54] [55]

Patients with stroke or other disorders who have impaired swallowing should be managed using a multidisciplinary approach.[56] Patients thought to have any risk factors for aspiration should be carefully evaluated before attempting feeding. This includes neurologic evaluation with assessment of cortical function, bulbar muscle gag, and cough reflex. A speech pathologist should evaluate swallowing in unclear cases.[48] Modified diets (nectar-thick instead of thin liquids) may be easier to swallow. These patients may benefit from swallow rehabilitation, including training like upright posture, chin tuck, and slow swallowing.

Keeping the head of the patient's bed higher than 30° to 45° reduces the risk of aspiration, especially in critically ill or mechanically ventilated patients.[57] [58] Other potentially useful measures to prevent aspiration in critically ill patients include oral decontamination with antiseptic solutions; a chin-down position while feeding for patients with dysphagia; percutaneous endoscopic gastrostomy tube or percutaneous endoscopic jejunostomy tube for feeding chronically debilitated patients; feeding by hand rather than inserting feeding tubes for geriatric patients; feeding a soft mechanical diet and thickened liquids; using capsaicin to sensitize the gag reflex; suctioning subglottic secretions in patients with endotracheal tubes; suppressing gastric acid with drugs; minimizing use of sedative drugs; monitoring gastric residual volumes as a marker of aspiration risks; and placing a postpyloric feeding tube.[37] [59]

Anesthesia

Anesthesia-related aspiration of gastric contents can be prevented by identifying patients susceptible to vomiting, minimizing gastric contents before surgery, minimizing emetic stimuli, and avoiding complete loss of protective reflexes from oversedation.[37] Antiemetics may be considered for patients at increased risk of postoperative nausea and vomiting.[54]

Aspiration of gastric contents during anesthesia can be prevented by adhering to national anesthesiology societies' guidelines. Water and other clear liquids (e.g., tea, coffee, soda water, apple juice, pulp-free orange juice) are allowed up to 2 hours before anesthesia in otherwise healthy adults (including pregnant women not in labor) and children who are scheduled for elective surgery.[54] [60] The fasting period after intake of solids should not be less than 6 hours. However, in the emergency procedural setting when the opportunity to use preventive strategies may not be available, there are no data to suggest that the risk of aspiration is increased due to lack of fasting.[61] Cricoid pressure (Sellick maneuver) is commonly employed to minimize the risk of aspiration during endotracheal intubation in patients at high risk of aspiration; however, its routine use is of uncertain benefit.[62][63]

A nasogastric tube is commonly inserted prophylactically in patients at high risk for aspiration during general anesthesia.[8]

Gastrointestinal stimulants or proton-pump inhibitors (or nonparticulate antacids) may be given to patients preoperatively only if they are at increased risk of aspiration.[54] Use of perioperative H₂ antagonists may maintain gastric pH >2.5 in unfasted patients, providing a theoretical benefit in preventing lung injury from aspiration.[64] [65][66] However, no data exist that demonstrate improvement in outcome with this approach. One meta-analysis comparing the effectiveness of H₂ antagonists with proton-pump inhibitors (PPIs) concluded that a single oral dose of an H₂ antagonist before surgery is more effective than PPI. However, when given as two oral doses preoperatively or using the intravenous route, both classes are equally effective.[67] Pregnant patients requiring anesthesia should receive a preoperative H₂ antagonist to increase gastric pH and undergo early intubation with cricoid pressure.[38] Routine preoperative use of H₂ antagonists, PPIs, antacids, antiemetics, or anticholinergics is not recommended for patients with no apparent increased risk for pulmonary aspiration.[54]

Pharmacologic management options to prevent aspiration are limited, and depend on the mechanism of aspiration risk. ACE inhibitors have been shown to increase the cough reflex in Chinese and Japanese patients at risk of aspiration after stroke.[68]

Barium studies

Postural techniques during barium studies can reduce or eliminate the risk of aspiration when small volumes of barium sulfate are used.[69] Radiologists should try to avoid barium entering the tracheobronchial tree by assessing the likelihood of aspiration in patients at risk before starting the exam.[4] Patients at high risk should swallow a contrast medium such as iopydol that is less likely to harm the lungs. Gastrografin can cause pulmonary edema and should be avoided if aspiration is suspected.[24]

Secondary prevention

Patients with ≥ 2 risk factors or with documented aspiration, persistent feeding intolerance, or both can be treated with a prokinetic drug and/or fed with tubes placed with the tip at or below the ligament of Treitz (suspensory ligament of the duodenum).[44] [116] [117] Relatively small, single-center trials have suggested reduction in pneumonia with both metoclopramide and postpyloric feeding.[116] [117] These findings need to be confirmed in larger randomized controlled studies. Improved oral hygiene may also decrease the risk of aspiration pneumonia.[16] [118]

Patients with stroke or other disorders who have impaired swallowing should be managed using a multidisciplinary approach.[56] Although swallowing function may return in most patients within 6 months of a stroke, tube feeding may be indicated in the acute phase. The Feed or Ordinary Diet (FOOD) study, a 3-part randomized controlled trial, found no significant benefit from nutritional supplements in patients with stroke.[119] [120] Early nutrition within the first week decreased mortality. In this study, percutaneous endoscopic gastrostomy tubes, when compared with nasogastric tubes, were associated with higher mortality or adverse outcomes at 6 months. The data available on the optimal type of feeding tube are conflicting. Postpyloric placement of feeding tubes has been shown to decrease pneumonia risk in small,

single-center studies, though no difference has been shown in other outcomes such as the duration of mechanical ventilation, vomiting, or mortality.[\[116\]](#) [\[121\]](#) [\[122\]](#) [\[123\]](#)[\[124\]](#)

Tracheostomy or laryngectomy may be options to be considered in highly select cases such as patients with neurodegenerative disorders (e.g., amyotrophic lateral sclerosis), or those with irreversible dysphagia and recurrent aspiration (e.g., previous radiation and surgery for throat cancer).

Patient discussions

All patients undergoing general anesthesia should fast preoperatively. Clear fluids of 400 mL or less (water, clear juices, coffee, or tea) up to 2 hours preoperatively, light meals up to 6 hours, and up to 150 mL of clear fluid up to 1 hour before surgery with oral drugs are recommended.[\[52\]](#) [\[54\]](#)[\[115\]](#)

These recommendations do not apply to milk, fat-containing fluids, or solids. For solids, a fasting period of at least 6 hours is recommended. Use of chewing gum and any form of tobacco is discouraged during the last 2 hours before induction of anesthesia because they can increase the gastric content. There are no clear recommendations regarding patients at high risk for aspiration.

Monitoring

Monitoring

Measuring gastric residual volume (GRV) in excess of a threshold value has been used to monitor the risk for aspiration in tube-fed patients. Because the cutoff values for GRV that clearly indicate aspiration risks vary among studies, these assessments should be combined with clinical evaluation of feeding intolerance, which includes auscultation of bowel sounds and evaluation of abdominal distention.[44] Measures to reduce aspirations should be initiated when a GRV is >200 mL, and aspiration of gastric contents from any in-place nasogastric tube should be performed if anesthesia is imminent.[109]

The quality of evidence regarding measurement of GRV and its effect on parameters such as mortality, pneumonia, and length of stay is uncertain, and it is difficult to draw conclusions about the value of GRV in the clinical setting.[110] The practice of not monitoring GRV was not inferior to routine monitoring in regard to the outcome of ventilator-associated pneumonia in 452 intensive care unit (ICU) patients randomized in a noninferiority, open-label, multicenter trial conducted in adults requiring invasive mechanical ventilation for more than 2 days and given enteral nutrition within 36 hours after intubation at nine French ICUs.[111] Furthermore, in one study of 61 patients receiving mechanical ventilation and continuous enteral feeding, measurement of residual gastric volume by suctioning was inaccurate.[112]

The Society of Critical Care Medicine (SCCM) and the American Society for Parenteral and Enteral Nutrition (ASPEN) do not endorse using GRV as part of routine monitoring in the ICU.[113] The European Society for Clinical Nutrition and Metabolism (ESPEN) does suggest delaying commencement of feeding if GRV is >500 mL in a 6-hour window but also acknowledges SCCM/ASPEN guidelines regarding routine monitoring.[114] Clinicians should consider these guidelines within the context of the individual patient, particularly in those at high risk for aspiration, who have ileus, esophageal dysfunction, or gastroparesis.

Because of the inert character of barium, long-term reactions and late toxicities are not usually expected, and complete radiologic clearance is the norm. There are no extensive data on long-term complications from massive barium aspiration, but case reports suggest that abnormalities can be seen on high-resolution computed tomography scan up to 1 year later.[108] Therefore, it is reasonable to obtain follow-up chest imaging during the year after barium aspiration.

Complications

Complications	Timeframe	Likelihood
pneumonia related to aspiration of gastric contents	short term	medium
One quarter of patients with aspiration of gastric contents develop expanding infiltrates, probably because of superimposed bacterial infection, which should be treated with antibiotics that cover for gram-negative bacteria and gram-positive cocci.[48] [77]		
acute respiratory distress syndrome related to aspiration of gastric contents	short term	low
May be a complication of pulmonary aspiration of gastric contents. About 12% die, usually shortly after the aspiration.[77]		
fibrosis related to aspiration of barium sulfate	long term	low
Mild fibrotic changes have been described on high-resolution computed tomography scans of the chest 1 year after barium aspiration.[108]		

Prognosis

Aspiration of gastric contents results in limited disease in most patients. Aspiration can be asymptomatic, but it can result in chemical pneumonitis and acute respiratory distress syndrome (ARDS).[2] Mortality is high, however, if patients progress to ARDS. ARDS usually develops within 2-5 hours of aspiration, but can be delayed.[15] The clinical course is variable: in one small retrospective study, 12% of patients had a fulminant course and died shortly after aspiration of gastric contents; in 62% the chest x-ray improved rapidly within 5 days after aspiration; however, in 26% this initial improvement was followed by the development of new infiltrates on chest x-ray, probably indicating complicating bacterial pneumonia, and of this group over 60% died as a consequence.[77] The aspiration injury usually begins to resolve within 72 hours, if pneumonia or ARDS has not developed. Death as a result of aspiration of gastric contents as a consequence of anesthetic practice is rare, with rates varying from 0% to 5%, but is a leading cause of malpractice claims with high mortality in these reports.[8] [12]

Because of the inert character of barium, long-term reactions and late toxicities are not usually expected, and complete radiologic clearance is the norm. There are no extensive data on long-term complications from massive barium aspiration, but case reports suggest that abnormalities can be seen on high-resolution computed tomography scan up to 1 year later.[108] Therefore, it is reasonable to obtain follow-up chest imaging during the year after barium aspiration.

Diagnostic guidelines

International

ACR appropriateness criteria: dysphagia (<https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria>) [94]

Published by: American College of Radiology

Last published: 2018

Key articles

- Marik PE. Aspiration pneumonitis and aspiration pneumonia. *N Engl J Med*. 2001 Mar 1;344(9):665-71. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11228282?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11228282?tool=bestpractice.bmj.com)
- Smith Hammond CA, Goldstein LB. Cough and aspiration of food and liquids due to oral-pharyngeal dysphagia: ACCP evidence-based clinical practice guidelines. *Chest*. 2006 Jan;129(1 suppl):154S-68S. [Full text \(http://journal.chestnet.org/article/S0012-3692\(15\)52844-4/fulltext\)](http://journal.chestnet.org/article/S0012-3692(15)52844-4/fulltext) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16428705?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16428705?tool=bestpractice.bmj.com)
- American Society of Anesthesiologists. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures - an updated report by the American Society of Anesthesiologists task force on preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration. *Anesthesiology*. 2017 Mar;126(3):376-93. [Full text \(https://journals.lww.com/anesthesiology/fulltext/2017/03000/practice_guidelines_for_preoperative_fasting_and.14.aspx\)](https://journals.lww.com/anesthesiology/fulltext/2017/03000/practice_guidelines_for_preoperative_fasting_and.14.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/28045707?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/28045707?tool=bestpractice.bmj.com)

References

1. Raghavendran K, Nemzek J, Napolitano LM, et al. Aspiration-induced lung injury. 2011 Apr;39(4):818-26. [Full text \(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3102154\)](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3102154) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/21263315?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/21263315?tool=bestpractice.bmj.com)
2. Marik PE. Aspiration pneumonitis and aspiration pneumonia. *N Engl J Med*. 2001 Mar 1;344(9):665-71. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11228282?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11228282?tool=bestpractice.bmj.com)
3. Lee AS, Lee JS, He Z, et al. Reflux-aspiration in chronic lung disease. *Ann Am Thorac Soc*. 2020 Feb;17(2):155-64. [Full text \(https://www.atsjournals.org/doi/10.1513/AnnalsATS.201906-427CME\)](https://www.atsjournals.org/doi/10.1513/AnnalsATS.201906-427CME) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/31697575?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/31697575?tool=bestpractice.bmj.com)
4. Gray C, Sivaloganathan S, Simpkins KC. Aspiration of high-density barium contrast medium causing acute pulmonary inflammation: report of two fatal cases in elderly women with disordered swallowing. 1989 Jul;40(4):397-400. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2758750?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2758750?tool=bestpractice.bmj.com)
5. Warner MA, Warner ME, Weber JG. Clinical significance of pulmonary aspiration during the perioperative period. *Anesthesiology*. 1993 Jan;78(1):56-62. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/8424572?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/8424572?tool=bestpractice.bmj.com)
6. Sakai T, Planinsic RM, Quinlan JJ, et al. The incidence and outcome of perioperative pulmonary aspiration in a university hospital: a 4-year retrospective analysis. *Anesth Analg*. 2006 Oct;103(4):941-7. [Full text \(https://journals.lww.com/anesthesia-analgesia/fulltext/2006/10000/the_incidence_and_outcome_of_perioperative.28.aspx\)](https://journals.lww.com/anesthesia-analgesia/fulltext/2006/10000/the_incidence_and_outcome_of_perioperative.28.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/17000809?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/17000809?tool=bestpractice.bmj.com)

7. Olsson GL, Hallen B, Hambraeus-Jonzon K. Aspiration during anaesthesia: a computer-aided study of 185,358 anaesthetics. *Acta Anaesthesiol Scand*. 1986 Jan;30(1):84-92. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/3754372?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/3754372?tool=bestpractice.bmj.com)
8. Ng A, Smith G. Gastroesophageal reflux and aspiration of gastric contents in anesthetic practice. *Anesth Analg*. 2001 Aug;93(2):494-513. [Full text \(http://www.anesthesia-analgesia.org/content/93/2/494.full\)](http://www.anesthesia-analgesia.org/content/93/2/494.full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11473886?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11473886?tool=bestpractice.bmj.com)
9. Walker RW. Pulmonary aspiration in pediatric anesthetic practice in the UK: a prospective survey of specialist pediatric centers over a one-year period. *Paediatr Anaesth*. 2013 Aug;23(8):702-11. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/23763657?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/23763657?tool=bestpractice.bmj.com)
10. Warner MA, Warner ME, Warner DO, et al. Perioperative pulmonary aspiration in infants and children. *Anesthesiology*. 1999 Jan;90(1):66-71. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9915314?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9915314?tool=bestpractice.bmj.com)
11. Borland LM, Sereika SM, Woelfel SK, et al. Pulmonary aspiration in pediatric patients during general anesthesia: incidence and outcome. *J Clin Anesth*. 1998 Mar;10(2):95-102. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9524892?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9524892?tool=bestpractice.bmj.com)
12. Warner MA, Meyerhoff KL, Warner ME, et al. Pulmonary aspiration of gastric contents: a closed claims analysis. *Anesthesiology*. 2021 Aug 1;135(2):284-91. [Full text \(https://journals.lww.com/anesthesiology/fulltext/2021/08000/pulmonary_aspiration_of_gastric_contents__a_closed.20.aspx\)](https://journals.lww.com/anesthesiology/fulltext/2021/08000/pulmonary_aspiration_of_gastric_contents__a_closed.20.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/34019629?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/34019629?tool=bestpractice.bmj.com)
13. Fernández-Sabé N, Carratalà J, Rosón B, et al. Community-acquired pneumonia in very elderly patients: causative organisms, clinical characteristics, and outcomes. *Medicine (Baltimore)*. 2003 May;82(3):159-69. [Full text \(https://journals.lww.com/md-journal/fulltext/2003/05000/community_acquired_pneumonia_in_very_elderly.2.aspx\)](https://journals.lww.com/md-journal/fulltext/2003/05000/community_acquired_pneumonia_in_very_elderly.2.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/12792302?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/12792302?tool=bestpractice.bmj.com)
14. Cabre M, Serra-Prat M, Palomera E, et al. Prevalence and prognostic implications of dysphagia in elderly patients with pneumonia. *Age Ageing*. 2010 Jan;39(1):39-45. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/19561160?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/19561160?tool=bestpractice.bmj.com)
15. Gajic O, Dabbagh O, Park PK, et al. Early identification of patients at risk of acute lung injury: evaluation of lung injury prediction score in a multicenter cohort study. *Am J Respir Crit Care Med*. 2011 Feb 15;183(4):462-70. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC3056224\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC3056224) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/20802164?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/20802164?tool=bestpractice.bmj.com)
16. Intercollegiate Stroke Working Party. National clinical guideline for stroke for the UK and Ireland. May 2023 [internet publication]. [Full text \(https://www.strokeguideline.org\)](https://www.strokeguideline.org)
17. Martino R, Foley N, Bhogal S, et al. Dysphagia after stroke: incidence, diagnosis, and pulmonary complications. *Stroke*. 2005 Dec;36(12):2756-63. [Full text \(https://www.ahajournals.org/doi/10.1161/01.STR.0000190056.76543.eb\)](https://www.ahajournals.org/doi/10.1161/01.STR.0000190056.76543.eb) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16269630?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16269630?tool=bestpractice.bmj.com)

18. Rofes L, Arreola V, Almirall J, et al. Diagnosis and management of oropharyngeal dysphagia and its nutritional and respiratory complications in the elderly. *Gastroenterol Res Pract*. 2011;2011:818979. [Full text \(https://onlinelibrary.wiley.com/doi/10.1155/2011/818979\)](https://onlinelibrary.wiley.com/doi/10.1155/2011/818979) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/20811545?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/20811545?tool=bestpractice.bmj.com)
19. Mullan H, Roubenoff RA, Roubenoff R. Risk of pulmonary aspiration among patients receiving enteral nutrition support. *JPEN J Parenter Enteral Nutr*. Mar-Apr 1992;16(2):160-4. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/1556813?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/1556813?tool=bestpractice.bmj.com)
20. Metheny NA, Clouse RE, Chang YH, et al. Tracheobronchial aspiration of gastric contents in critically ill tube-fed patients: frequency, outcomes, and risk factors. *Crit Care Med*. 2006 Apr;34(4):1007-15. [Full text \(http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2396145\)](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2396145) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16484901?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16484901?tool=bestpractice.bmj.com)
21. Liu Y, Wang Y, Zhang B, et al. Gastric-tube versus post-pyloric feeding in critical patients: a systematic review and meta-analysis of pulmonary aspiration- and nutrition-related outcomes. *Eur J Clin Nutr*. 2021 Sep;75(9):1337-48. [Full text \(https://www.nature.com/articles/s41430-021-00860-2\)](https://www.nature.com/articles/s41430-021-00860-2) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/33536570?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/33536570?tool=bestpractice.bmj.com)
22. Lockey DJ, Coats T, Parr MJ. Aspiration in severe trauma: a prospective study. *Anaesthesia*. 1999 Nov;54(11):1097-8. [Full text \(https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-2044.1999.00754.x?sid=nlm%3Apubmed\)](https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-2044.1999.00754.x?sid=nlm%3Apubmed) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/10540100?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/10540100?tool=bestpractice.bmj.com)
23. Fung KP, Seagram G, Pasioka J, et al. Investigation and outcome of 121 infants and children requiring Nissen fundoplication for the management of gastroesophageal reflux. *Clin Invest Med*. 1990 Oct;13(5):237-46. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2276217?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2276217?tool=bestpractice.bmj.com)
24. Tamm I, Kortsik C. Severe barium sulfate aspiration into the lung: clinical presentation, prognosis and therapy. *Respiration*. 1999;66(1):81-4. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9973698?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9973698?tool=bestpractice.bmj.com)
25. Ansell G. Alimentary tract. In: Ansell G, Bettmann MA, Kaufman JA, et al, eds. *Complications in diagnostic imaging and interventional radiology*. 2nd ed. Oxford, UK: Blackwell Scientific; 1987:218-9.
26. Robinson M, Davidson A. Aspiration under anaesthesia: risk assessment and decision-making. *Contin Educ Anaesth Crit Care Pain*. 2014 Aug;14(4):171-5. [Full text \(https://academic.oup.com/bjaed/article/14/4/171/293792\)](https://academic.oup.com/bjaed/article/14/4/171/293792)
27. Smith Hammond CA, Goldstein LB. Cough and aspiration of food and liquids due to oral-pharyngeal dysphagia: ACCP evidence-based clinical practice guidelines. *Chest*. 2006 Jan;129(1 suppl):154S-68S. [Full text \(http://journal.chestnet.org/article/S0012-3692\(15\)52844-4/fulltext\)](http://journal.chestnet.org/article/S0012-3692(15)52844-4/fulltext) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16428705?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16428705?tool=bestpractice.bmj.com)
28. Umemoto G, Furuya H. Management of dysphagia in patients with Parkinson's disease and related disorders. *Intern Med*. 2020 Jan 1;59(1):7-14. [Full text \(https://www.jstage.jst.go.jp/article/internalmedicine/59/1/59_2373-18/_article\)](https://www.jstage.jst.go.jp/article/internalmedicine/59/1/59_2373-18/_article) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/30996170?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/30996170?tool=bestpractice.bmj.com)

29. Marin-Corral J, Pascual-Guardia S, Amati F, et al. Aspiration risk factors, microbiology, and empiric antibiotics for patients hospitalized with community-acquired pneumonia. *Chest*. 2021 Jan;159(1):58-72. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/32687909?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/32687909?tool=bestpractice.bmj.com)
30. Yamamoto T, Kobayashi Y, Murata M. Risk of pneumonia onset and discontinuation of oral intake following videofluorography in patients with Lewy body disease. *Parkinsonism Relat Disord*. 2010 Sep;16(8):503-6. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/20584622?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/20584622?tool=bestpractice.bmj.com)
31. McClave SA, Lukan JK, Stefater JA, et al. Poor validity of residual volumes as a marker for risk of aspiration in critically ill patients. *Crit Care Med*. 2005 Feb;33(2):324-30. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15699835?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15699835?tool=bestpractice.bmj.com)
32. Silveira SQ, da Silva LM, de Campos Vieira Abib A, et al. Relationship between perioperative semaglutide use and residual gastric content: a retrospective analysis of patients undergoing elective upper endoscopy. *J Clin Anesth*. 2023 Aug;87:111091. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/36870274?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/36870274?tool=bestpractice.bmj.com)
33. Nersessian RSF, da Silva LM, Carvalho MAS, et al. Relationship between residual gastric content and peri-operative semaglutide use assessed by gastric ultrasound: a prospective observational study. *Anaesthesia*. 2024 Dec;79(12):1317-24. [Full text \(https://associationofanaesthetists-publications.onlinelibrary.wiley.com/doi/10.1111/anae.16454\)](https://associationofanaesthetists-publications.onlinelibrary.wiley.com/doi/10.1111/anae.16454) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/39435967?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/39435967?tool=bestpractice.bmj.com)
34. El-Boghdady K, Dhesi J, Fabb P, et al. Elective peri-operative management of adults taking glucagon-like peptide-1 receptor agonists, glucose-dependent insulintropic peptide agonists and sodium-glucose cotransporter-2 inhibitors: a multidisciplinary consensus statement: a consensus statement from the Association of Anaesthetists, Association of British Clinical Diabetologists, British Obesity and Metabolic Surgery Society, Centre for Perioperative Care, Joint British Diabetes Societies for Inpatient Care, Royal College of Anaesthetists, Society for Obesity and Bariatric Anaesthesia, and UK Clinical Pharmacy Association. *Anaesthesia*. 2025 Apr;80(4):412-24. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC11885194\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC11885194) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/39781571?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/39781571?tool=bestpractice.bmj.com)
35. Boyd M, Chatterjee A, Chiles C, et al. Tracheobronchial foreign body aspiration in adults. *South Med J*. 2009 Feb;102(2):171-4. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/19139679?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/19139679?tool=bestpractice.bmj.com)
36. Shaker R, Li Q, Ren J, et al. Coordination of deglutition and phases of respiration: effect of aging, tachypnea, bolus volume, and chronic obstructive pulmonary disease. *Am J Physiol*. 1992 Nov;263(5 Pt 1):G750-5. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/1443150?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/1443150?tool=bestpractice.bmj.com)
37. Vaughan GG, Grycko RJ, Montgomery MT. The prevention and treatment of aspiration of vomitus during pharmacosedation and general anesthesia. *J Oral Maxillofac Surg*. 1992 Aug;50(8):874-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/1634979?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/1634979?tool=bestpractice.bmj.com)

38. Royal College of Obstetricians and Gynaecologists. Maternal collapse in pregnancy and the puerperium. Green-top Guideline No. 56. Dec 2019 [internet publication]. [Full text \(https://obgyn.onlinelibrary.wiley.com/doi/10.1111/1471-0528.15995\)](https://obgyn.onlinelibrary.wiley.com/doi/10.1111/1471-0528.15995)
39. Hunt EB, Sullivan A, Galvin J, et al. Gastric aspiration and its role in airway inflammation. *Open Respir Med J*. 2018;12:1-10. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC5806178\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC5806178) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/29456774?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/29456774?tool=bestpractice.bmj.com)
40. Abou Jaoude P, Knight PR, Ohtake P, et al. Biomarkers in the diagnosis of aspiration syndromes. *Expert Rev Mol Diagn*. 2010 Apr;10(3):309-19. [Full text \(http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2882092\)](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2882092) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/20370588?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/20370588?tool=bestpractice.bmj.com)
41. Green SM, Krauss B. Pulmonary aspiration risk during emergency department procedural sedation: an examination of the role of fasting and sedation depth. *Acad Emerg Med*. 2002 Jan;9(1):35-42. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11772667?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11772667?tool=bestpractice.bmj.com)
42. Zuercher P, Moret CS, Dziewas R, et al. Dysphagia in the intensive care unit: epidemiology, mechanisms, and clinical management. *Crit Care*. 2019 Mar 28;23(1):103. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC6438038\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC6438038) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/30922363?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/30922363?tool=bestpractice.bmj.com)
43. Wick R, Gilbert JD, Byard RW. Café coronary syndrome - fatal choking on food: an autopsy approach. *J Clin Forensic Med*. 2006 Apr;13(3):135-8. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16356749?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16356749?tool=bestpractice.bmj.com)
44. Mizock BA. Risk of aspiration in patients on enteral nutrition: frequency, relevance, relation to pneumonia, risk factors, and strategies for risk reduction. *Curr Gastroenterol Rep*. 2007 Aug;9(4):338-44. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/17883984?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/17883984?tool=bestpractice.bmj.com)
45. Lundy DS, Smith C, Colangelo L, et al. Aspiration: cause and implications. *Otolaryngol Head Neck Surg*. 1999 Apr;120(4):474-8. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/10187936?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/10187936?tool=bestpractice.bmj.com)
46. Elpern EH, Scott MG, Petro L, et al. Pulmonary aspiration in mechanically ventilated patients with tracheostomies. *Chest*. 1994 Feb;105(2):563-6. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/8306764?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/8306764?tool=bestpractice.bmj.com)
47. Elpern EH, Jacobs ER, Bone RC. Incidence of aspiration in tracheally intubated adults. *Heart Lung*. 1987 Sep;16(5):527-31. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/3115917?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/3115917?tool=bestpractice.bmj.com)
48. Paintal HS, Kuschner WG. Aspiration syndromes: 10 clinical pearls every physician should know. *Int J Clin Pract*. 2007 May;61(5):846-52. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/17493092?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/17493092?tool=bestpractice.bmj.com)
49. Fernández-Carmona A, Peñas-Maldonado L, Yuste-Orsio E, et al. Exploration and approach to artificial airway dysphagia. *Med Intensiva*. Aug-Sep 2012;36(6):423-33. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/22055775?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/22055775?tool=bestpractice.bmj.com)

50. Leder SB, Cohn SM, Moller BA. Fiberoptic endoscopic documentation of the high incidence of aspiration following extubation in critically ill trauma patients. *Dysphagia*. Fall 1998;13(4):208-12. [Full text \(https://link.springer.com/article/10.1007/PL00009573\)](https://link.springer.com/article/10.1007/PL00009573) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9716751?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9716751?tool=bestpractice.bmj.com)
51. Langmore SE, Terpenning MS, Schork A, et al. Predictors of aspiration pneumonia: how important is dysphagia? *Dysphagia*. Spring 1998;13(2):69-81. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9513300?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9513300?tool=bestpractice.bmj.com)
52. Soreide E, Eriksson LI, Hirlekar G, et al. Pre-operative fasting guidelines: an update. *Acta Anaesthesiol Scand*. 2005 Sep;49(8):1041-7. [Full text \(https://onlinelibrary.wiley.com/doi/full/10.1111/j.1399-6576.2005.00781.x\)](https://onlinelibrary.wiley.com/doi/full/10.1111/j.1399-6576.2005.00781.x) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16095440?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16095440?tool=bestpractice.bmj.com)
53. Beal M, Chesson A, Garcia T, et al. A pilot study of quantitative aspiration in patients with symptoms of obstructive sleep apnea: comparison to a historic control group. *Laryngoscope*. 2004 Jun;114(6):965-8. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15179196?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15179196?tool=bestpractice.bmj.com)
54. American Society of Anesthesiologists. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures - an updated report by the American Society of Anesthesiologists task force on preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration. *Anesthesiology*. 2017 Mar;126(3):376-93. [Full text \(https://journals.lww.com/anesthesiology/fulltext/2017/03000/practice_guidelines_for_preoperative_fasting_and.14.aspx\)](https://journals.lww.com/anesthesiology/fulltext/2017/03000/practice_guidelines_for_preoperative_fasting_and.14.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/28045707?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/28045707?tool=bestpractice.bmj.com)
55. Sarin J, Balasubramaniam R, Corcoran AM, et al. Reducing the risk of aspiration pneumonia among elderly patients in long-term care facilities through oral health interventions. *J Am Med Dir Assoc*. 2008 Feb;9(2):128-35. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/18261707?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/18261707?tool=bestpractice.bmj.com)
56. Australian and New Zealand Society for Geriatric Medicine. Position statement abstract: dysphagia and aspiration in older people. *Australas J Ageing*. 2020 Mar;39(1):85. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/31313474?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/31313474?tool=bestpractice.bmj.com)
57. Healthcare Infection Control Practices Advisory Committee; Centers for Disease Control and Prevention (US). Guidelines for preventing health-care-associated pneumonia, 2003 recommendations of the CDC and the Healthcare Infection Control Practices Advisory Committee. *Respir Care*. 2004 Aug;49(8):926-39. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15320333?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15320333?tool=bestpractice.bmj.com)
58. van Nieuwenhoven CA, Vandenbroucke-Grauls C, van Tiel FH, et al. Feasibility and effects of the semirecumbent position to prevent ventilator-associated pneumonia: a randomized study. *Crit Care Med*. 2006 Feb;34(2):396-402. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16424720?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16424720?tool=bestpractice.bmj.com)

59. El Solh AA, Saliba R. Pharmacologic prevention of aspiration pneumonia: a systematic review. *Am J Geriatr Pharmacother*. 2007 Dec;5(4):352-62. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/18179994?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/18179994?tool=bestpractice.bmj.com)
60. Smith I, Kranke P, Murat I, et al. Perioperative fasting in adults and children: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol*. 2011 Aug;28(8):556-69. [Full text \(https://journals.lww.com/ejanaesthesiology/Fulltext/2011/08000/Perioperative_fasting_in_adults_and_children_4.aspx\)](https://journals.lww.com/ejanaesthesiology/Fulltext/2011/08000/Perioperative_fasting_in_adults_and_children_4.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/21712716?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/21712716?tool=bestpractice.bmj.com)
61. Thorpe RJ, Bengner J. Pre-procedural fasting in emergency sedation. *Emerg Med J*. 2010 Apr;27(4):254-61. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/20385672?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/20385672?tool=bestpractice.bmj.com)
62. Zdravkovic M, Rice MJ, Brull SJ. The clinical use of cricoid pressure: first, do no harm. *Anesth Analg*. 2021 Jan;132(1):261-7. [Full text \(https://journals.lww.com/anesthesia-analgnesia/fulltext/2021/01000/the_clinical_use_of_cricoid_pressure_first_do_no.35.aspx\)](https://journals.lww.com/anesthesia-analgnesia/fulltext/2021/01000/the_clinical_use_of_cricoid_pressure_first_do_no.35.aspx) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/31397697?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/31397697?tool=bestpractice.bmj.com)
63. Birenbaum A, Hajage D, Roche S, et al. Effect of cricoid pressure compared with a sham procedure in the rapid sequence induction of anesthesia: the IRIS randomized clinical trial. *JAMA Surg*. 2019 Jan 1;154(1):9-17. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC6439856\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC6439856) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/30347104?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/30347104?tool=bestpractice.bmj.com)
64. Paranjothy S, Griffiths JD, Broughton HK, et al. Interventions at caesarean section for reducing the risk of aspiration pneumonitis. *Cochrane Database Syst Rev*. 2014 Feb 5;(2):CD004943. [Full text \(http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD004943.pub4/full\)](http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD004943.pub4/full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/24497372?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/24497372?tool=bestpractice.bmj.com)
65. Gyte GM, Richens Y. Routine prophylactic drugs in normal labour for reducing gastric aspiration and its effects. *Cochrane Database Syst Rev*. 2006 Jul 19;(3):CD005298. [Full text \(https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD005298.pub2/full\)](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD005298.pub2/full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16856089?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16856089?tool=bestpractice.bmj.com)
66. Clark K, Lam LT, Gibson S, et al. The effect of ranitidine versus proton pump inhibitors on gastric secretions: a meta-analysis of randomised control trials. *Anaesthesia*. 2009 Jun;64(6):652-7. [Full text \(http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2044.2008.05861.x/full\)](http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2044.2008.05861.x/full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/19453319?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/19453319?tool=bestpractice.bmj.com)
67. Puig I, Calzado S, Suárez D, Sánchez-Delgado J, López S, Calvet X. Meta-analysis: comparative efficacy of H2-receptor antagonists and proton pump inhibitors for reducing aspiration risk during anaesthesia depending on the administration route and schedule. *Pharmacol Res*. 2012 Apr;65(4):480-90. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/22289674?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/22289674?tool=bestpractice.bmj.com)
68. Simpson AJ, Allen JL, Chatwin M, et al. BTS clinical statement on aspiration pneumonia. *Thorax*. 2023 Feb;78(suppl 1):s3-21. [Full text \(https://thorax.bmj.com/content/78/Suppl_1/s3.long\)](https://thorax.bmj.com/content/78/Suppl_1/s3.long) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/36863772?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/36863772?tool=bestpractice.bmj.com)

69. Rasley A, Logemann JA, Kahrilas PJ, et al. Prevention of barium aspiration during videofluoroscopic swallowing studies: value of change in posture. *AJR Am J Roentgenol*. 1993 May;160(5):1005-9. Full text (<http://www.ajronline.org/doi/pdf/10.2214/ajr.160.5.8470567>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/8470567?tool=bestpractice.bmj.com>)
70. Chiu CY, Wong KS, Tsai MH. Massive aspiration of barium sulfate during an upper gastrointestinal examination in a child with dysphagia. *Int J Pediatr Otorhinolaryngol*. 2005 Apr;69(4):541-4. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15763294?tool=bestpractice.bmj.com>)
71. Basu S, Kumar A, Das BK. Accidental aspiration of barium sulphate in an infant. *Pediatr Radiol*. 2009 Jul;39(7):762. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/19399489?tool=bestpractice.bmj.com>)
72. Fuentes Santos C, Steen B. Aspiration of barium contrast. *Case Rep Pulmonol*. 2014;2014:215832. Full text (<https://pmc.ncbi.nlm.nih.gov/articles/PMC4189518>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/25309769?tool=bestpractice.bmj.com>)
73. Franquet T, Gimenez A, Roson N, et al. Aspiration diseases: findings, pitfalls, and differential diagnosis. *Radiographics*. May-Jun 2000;20(3):673-85. Full text (<http://radiographics.rsna.org/content/20/3/673.full>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/10835120?tool=bestpractice.bmj.com>)
74. Yan GW, Deng JF, Bhetuwal A, et al. A case report and literature review of barium sulphate aspiration during upper gastrointestinal examination. *Medicine (Baltimore)*. 2017 Nov;96(47):e8821. Full text (https://journals.lww.com/md-journal/fulltext/2017/11270/a_case_report_and_literature_review_of_barium.78.aspx) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/29381987?tool=bestpractice.bmj.com>)
75. Schwarz MI, Albert RK. "Imitators" of the ARDS: implications for diagnosis and treatment. *Chest*. 2004 Apr;125(4):1530-5. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15078770?tool=bestpractice.bmj.com>)
76. Bandorski D, Tello K, Erdal H, et al. Clinical utility of pepsin and bile acid in tracheal secretions for accurate diagnosis of aspiration in ICU patients. *J Clin Med*. 2023 Aug 23;12(17):5466. Full text (<https://pmc.ncbi.nlm.nih.gov/articles/PMC10487459>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/37685534?tool=bestpractice.bmj.com>)
77. Bynum LJ, Pierce AK. Pulmonary aspiration of gastric contents. *Am Rev Respir Dis*. 1976 Dec;114(6):1129-36. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/1008348?tool=bestpractice.bmj.com>)
78. Kluger MT, Visvanathan T, Myburgh JA, et al. Crisis management during anaesthesia: regurgitation, vomiting, and aspiration. *Qual Saf Health Care*. 2005 Jun;14(3):e4. Full text (<https://qualitysafety.bmj.com/content/14/3/e4.long>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15933301?tool=bestpractice.bmj.com>)
79. Light RW. Parapneumonic effusions and empyema. *Proc Am Thorac Soc*. 2006;3(1):75-80. Full text (<https://www.atsjournals.org/doi/pdf/10.1513/pats.200510-113JH>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/16493154?tool=bestpractice.bmj.com>)

80. Bernard GR. Acute respiratory distress syndrome: a historical perspective. *Am J Respir Crit Care Med*. 2005 Oct 1;172(7):798-806. [Full text \(http://www.atsjournals.org/doi/full/10.1164/rccm.200504-663OE#.UtiD1tLFLWQ\)](http://www.atsjournals.org/doi/full/10.1164/rccm.200504-663OE#.UtiD1tLFLWQ) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/16020801?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/16020801?tool=bestpractice.bmj.com)
81. Martinati LC, Boner AL. Clinical diagnosis of wheezing in early childhood. *Allergy*. 1995 Sep;50(9):701-10. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/8546263?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/8546263?tool=bestpractice.bmj.com)
82. Fireman P. The wheezing infant. *Pediatr Rev*. 1986 Feb;7(8):247-54. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/3331439?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/3331439?tool=bestpractice.bmj.com)
83. Davie AP, Francis CM, Caruana L, et al. Assessing diagnosis in heart failure: which features are any use? *QJM*. 1997 May;90(5):335-9. [Full text \(http://qjmed.oxfordjournals.org/cgi/reprint/90/5/335\)](http://qjmed.oxfordjournals.org/cgi/reprint/90/5/335) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9205668?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9205668?tool=bestpractice.bmj.com)
84. Marcus GM, Gerber IL, McKeown BH, et al. Association between phonocardiographic third and fourth heart sounds and objective measures of left ventricular function. *JAMA*. 2005 May 11;293(18):2238-44. [Full text \(http://jama.ama-assn.org/cgi/content/full/293/18/2238\)](http://jama.ama-assn.org/cgi/content/full/293/18/2238) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15886379?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15886379?tool=bestpractice.bmj.com)
85. Maisel AS, Krishnaswamy P, Nowak RM, et al. Rapid measurement of B-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med*. 2002 Jul 18;347(3):161-7. [Full text \(http://www.nejm.org/doi/full/10.1056/NEJMoa020233#t=article\)](http://www.nejm.org/doi/full/10.1056/NEJMoa020233#t=article) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/12124404?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/12124404?tool=bestpractice.bmj.com)
86. National Safety Council. Leading causes of death: an excerpt of injury facts. Itasca, IL: National Safety Council Press; 2003:10-15.
87. Colice GL, Matthay MA, Bass E, et al. Neurogenic pulmonary edema. *Am Rev Respir Dis*. 1984 Nov;130(5):941-8. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/6388446?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/6388446?tool=bestpractice.bmj.com)
88. Brodsky MB, Suiter DM, González-Fernández M, et al. Screening accuracy for aspiration using bedside water swallow tests: a systematic review and meta-analysis. *Chest*. 2016 Jul;150(1):148-63. [Full text \(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4980548\)](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4980548) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/27102184?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/27102184?tool=bestpractice.bmj.com)
89. Doggett DL, Tappe KA, Mitchell MD, et al. Prevention of pneumonia in elderly stroke patients by systematic diagnosis and treatment of dysphagia: an evidence-based comprehensive analysis of the literature. *Dysphagia*. Fall 2001;16(4):279-95. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11720404?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11720404?tool=bestpractice.bmj.com)
90. Marik PE, Kaplan D. Aspiration pneumonia and dysphagia in the elderly. *Chest*. 2003 Jul;124(1):328-36. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/12853541?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/12853541?tool=bestpractice.bmj.com)
91. Odderson IR, Keaton JC, McKenna BS. Swallow management in patients on an acute stroke pathway: quality is cost effective. *Arch Phys Med Rehabil*. 1995 Dec;76(12):1130-3. [Full text \(https://\)](https://)

- [www.archives-pmr.org/article/S0003-9993\(95\)80121-9/pdf](http://www.archives-pmr.org/article/S0003-9993(95)80121-9/pdf) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/8540789?tool=bestpractice.bmj.com>)
92. Festic E, Soto JS, Pitre LA, et al. Novel bedside phonetic evaluation to identify dysphagia and aspiration risk. *Chest*. 2016 Mar;149(3):649-59. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/26203916?tool=bestpractice.bmj.com>)
 93. Boaden E, Burnell J, Hives L, et al. Screening for aspiration risk associated with dysphagia in acute stroke. *Cochrane Database Syst Rev*. 2021 Oct 18;10(10):CD012679. Full text (<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012679.pub2/full>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/34661279?tool=bestpractice.bmj.com>)
 94. American College of Radiology. ACR appropriateness criteria: dysphagia. 2018 [internet publication]. Full text (<https://acsearch.acr.org/docs/69471/Narrative>)
 95. Daniels SK, Brailey K, Priestly DH, et al. Aspiration in patients with acute stroke. *Arch Phys Med Rehabil*. 1998 Jan;79(1):14-9. Full text ([https://www.archives-pmr.org/article/S0003-9993\(98\)90200-3/pdf](https://www.archives-pmr.org/article/S0003-9993(98)90200-3/pdf)) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/9440410?tool=bestpractice.bmj.com>)
 96. Troche MS, Curtis JA, Sevitz JS, et al. Rehabilitating cough dysfunction in Parkinson's disease: a randomized controlled trial. *Mov Disord*. 2023 Feb;38(2):201-11. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/36345090?tool=bestpractice.bmj.com>)
 97. Choi J, Baek S, Kim G, et al. Peak voluntary cough flow and oropharyngeal dysphagia as risk factors for pneumonia. *Ann Rehabil Med*. 2021 Dec;45(6):431-9. Full text (<https://www.e-arm.org/journal/view.php?doi=10.5535/arm.21068>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/35000368?tool=bestpractice.bmj.com>)
 98. Dines DE, Titus JL, Sessler AD. Aspiration pneumonitis. *Mayo Clin Proc*. 1970 May;45(5):347-60. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/5443233?tool=bestpractice.bmj.com>)
 99. Hackett AP, Trinick RE, Rose K, et al. Weakly acidic pH reduces inflammatory cytokine expression in airway epithelial cells. *Respir Res*. 2016 Jul 15;17(1):82. Full text (<https://respiratory-research.biomedcentral.com/articles/10.1186/s12931-016-0399-3>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/27422381?tool=bestpractice.bmj.com>)
 100. Metlay JP, Waterer GW, Long AC, et al. Diagnosis and treatment of adults with community-acquired pneumonia. An official clinical practice guideline of the American Thoracic Society and Infectious Diseases Society of America. *Am J Respir Crit Care Med*. 2019 Oct 1;200(7):e45-67. Full text (<https://www.atsjournals.org/doi/10.1164/rccm.201908-1581ST>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/31573350?tool=bestpractice.bmj.com>)
 101. Lascarrou JB, Lissonde F, Le Thuaut A, et al. Antibiotic therapy in comatose mechanically ventilated patients following aspiration: differentiating pneumonia from pneumonitis. *Crit Care Med*. 2017 Aug;45(8):1268-75. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/28594680?tool=bestpractice.bmj.com>)
 102. Mandell LA, Niederman MS. Aspiration pneumonia. *N Engl J Med*. 2019 Feb 14;380(7):651-63.

103. Wolfe JE, Bone RC, Ruth WE. Effects of corticosteroids in the treatment of patients with gastric aspiration. *Am J Med.* 1977 Nov;63(5):719-22. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/930946?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/930946?tool=bestpractice.bmj.com)
104. Sukumaran M, Granada MJ, Berger HW, et al. Evaluation of corticosteroid treatment in aspiration of gastric contents: a controlled clinical trial. *Mt Sinai J Med.* Jul-Aug 1980;47(4):335-40. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/6997729?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/6997729?tool=bestpractice.bmj.com)
105. Lee M, Sukumaran M, Berger HW, et al. Influence of corticosteroid treatment on pulmonary function after recovery from aspiration of gastric contents. *Mt Sinai J Med.* Jul-Aug 1980;47(4):341-6. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/6997730?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/6997730?tool=bestpractice.bmj.com)
106. Zhao JN, Liu Y, Li HC. Corticosteroids in treatment of aspiration-related acute respiratory distress syndrome: results of a retrospective cohort study. *BMC Pulm Med.* 2016 Feb 10;16:29. [Full text \(https://bmcpulmed.biomedcentral.com/articles/10.1186/s12890-016-0194-4\)](https://bmcpulmed.biomedcentral.com/articles/10.1186/s12890-016-0194-4) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/26864571?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/26864571?tool=bestpractice.bmj.com)
107. Qadir N, Sahetya S, Munshi L, et al. An update on management of adult patients with acute respiratory distress syndrome: an official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med.* 2024 Jan 1;209(1):24-36. [Full text \(https://pmc.ncbi.nlm.nih.gov/articles/PMC10870893\)](https://pmc.ncbi.nlm.nih.gov/articles/PMC10870893) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/38032683?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/38032683?tool=bestpractice.bmj.com)
108. Voloudaki A, Ergazakis N, Gourtsoyiannis N. Late changes in barium sulfate aspiration: HRCT features. *Eur Radiol.* 2003 Sep;13(9):2226-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/12928969?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/12928969?tool=bestpractice.bmj.com)
109. McClave SA, Snider HL. Clinical use of gastric residual volumes as a monitor for patients on enteral tube feeding. *JPEN J Parenter Enteral Nutr.* Nov-Dec 2002;26(6 suppl):S43-8;discussion S49-50. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/12405622?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/12405622?tool=bestpractice.bmj.com)
110. Yasuda H, Kondo N, Yamamoto R, et al. Monitoring of gastric residual volume during enteral nutrition. *Cochrane Database Syst Rev.* 2021 Sep 27;9(9):CD013335. [Full text \(https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013335.pub2/full\)](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013335.pub2/full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/34596901?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/34596901?tool=bestpractice.bmj.com)
111. Reignier J, Mercier E, Le Gouge A, et al. Effect of not monitoring residual gastric volume on risk of ventilator-associated pneumonia in adults receiving mechanical ventilation and early enteral feeding: a randomized controlled trial. *JAMA.* 2013 Jan 16;309(3):249-56. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/23321763?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/23321763?tool=bestpractice.bmj.com)
112. Bouvet L, Zieleskiewicz L, Loubradou E, et al. Reliability of gastric suctioning compared with ultrasound assessment of residual gastric volume: a prospective multicentre cohort study. *Anaesthesia.* 2020 Mar;75(3):323-30. [Full text \(https://associationofanaesthetists-publications.onlinelibrary.wiley.com/doi/10.1111/anae.14915\)](https://associationofanaesthetists-publications.onlinelibrary.wiley.com/doi/10.1111/anae.14915) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/31802485?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/31802485?tool=bestpractice.bmj.com)
113. McClave SA, Taylor BE, Martindale RG, et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter*

Enteral Nutr. 2016 Feb;40(2):159-211. Full text (<https://aspenjournals.onlinelibrary.wiley.com/doi/10.1177/0148607115621863>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/26773077?tool=bestpractice.bmj.com>)

114. Singer P, Blaser AR, Berger MM, et al. ESPEN practical and partially revised guideline: clinical nutrition in the intensive care unit. Clin Nutr. 2023 Sep;42(9):1671-89. Full text ([https://www.clinicalnutritionjournal.com/article/S0261-5614\(23\)00230-3/fulltext](https://www.clinicalnutritionjournal.com/article/S0261-5614(23)00230-3/fulltext)) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/37517372?tool=bestpractice.bmj.com>)
115. Brady M, Kinn S, Stuart P, et al. Preoperative fasting for adults to prevent perioperative complications. Cochrane Database Syst Rev. 2003 Oct 20;(4):CD004423. Full text (<https://mrw.interscience.wiley.com/cochrane/clsystrev/articles/CD004423/frame.html>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/14584013?tool=bestpractice.bmj.com>)
116. Wang D, Zheng SQ, Chen XC, et al. Comparisons between small intestinal and gastric feeding in severe traumatic brain injury: a systematic review and meta-analysis of randomized controlled trials. J Neurosurg. 2015 Nov;123(5):1194-201. Full text (<https://thejns.org/view/journals/j-neurosurg/123/5/article-p1194.xml>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/26024007?tool=bestpractice.bmj.com>)
117. Warusevitane A, Karunatilake D, Sim J, et al. Safety and effect of metoclopramide to prevent pneumonia in patients with stroke fed via nasogastric tubes trial. Stroke. 2015 Feb;46(2):454-60. Full text (<http://stroke.ahajournals.org/content/46/2/454.long>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/25516196?tool=bestpractice.bmj.com>)
118. Pace CC, McCullough GH. The association between oral microorganisms and aspiration pneumonia in the institutionalized elderly: review and recommendations. Dysphagia. 2010 Dec;25(4):307-22. Full text (<https://link.springer.com/article/10.1007%2Fs00455-010-9298-9>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/20824288?tool=bestpractice.bmj.com>)
119. Dennis MS, Lewis SC, Warlow C; FOOD Trial Collaboration. Routine oral nutritional supplementation for stroke patients in hospital (FOOD): a multicentre randomised controlled trial. Lancet. 2005 Feb 26-Mar 4;365(9461):755-63. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15733716?tool=bestpractice.bmj.com>)
120. Dennis MS, Lewis SC, Warlow C; FOOD Trial Collaboration. Effect of timing and method of enteral tube feeding for dysphagic stroke patients (FOOD): a multicentre randomised controlled trial. Lancet. 2005 Feb 26-Mar 4;365(9461):764-72. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15733717?tool=bestpractice.bmj.com>)
121. Park RH, Allison MC, Lang J, et al. Randomised comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding in patients with persisting neurological dysphagia. BMJ. 1992 May 30;304(6839):1406-9. Full text (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1882203>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/1628013?tool=bestpractice.bmj.com>)
122. Strong RM, Condon SC, Solinger MR, et al. Equal aspiration rates from postpylorus and intragastric-placed small-bore nasoenteric feeding tubes: a randomized, prospective study. JPEN J Parenter

Enteral Nutr. Jan-Feb 1992;16(1):59-63. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/1738222?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/1738222?tool=bestpractice.bmj.com)

123. Spain DA, DeWeese RC, Reynolds MA, et al. Transpyloric passage of feeding tubes in patients with head injuries does not decrease complications. J Trauma. 1995 Dec;39(6):1100-2. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/7500401?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/7500401?tool=bestpractice.bmj.com)
124. Alkhawaja S, Martin C, Butler RJ, et al. Post-pyloric versus gastric tube feeding for preventing pneumonia and improving nutritional outcomes in critically ill adults. Cochrane Database Syst Rev. 2015 Aug 4;(8):CD008875. [Full text \(http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008875.pub2/full\)](http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008875.pub2/full) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/26241698?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/26241698?tool=bestpractice.bmj.com)

Images



Figure 1: Bronchoscopy showing barium aspiration in a lung transplant patient in the right mainstem bronchus after a barium swallow study

From the collection of Dr Kamran Mahmood



Figure 2: Barium aspiration. A barium swallow was conducted in a 53-year-old woman. Imaging revealed hyperdense airway-centered material in the left lower lobe consistent with barium aspiration bronchiolitis. A tracheoesophageal fistula was confirmed

From the collection of Dr Augustine Lee; used with permission of Mayo Foundation for Medical Education and Research, all rights reserved




Figure 3: A. Portable upright chest x-ray before aspiration; B. Chest x-ray 1 hour after aspiration, showing bilateral diffuse alveolar infiltrates, worse at the bases on the right side

From the collection of Dr Henri Colt




Figure 4: Lipoid pneumonia. A 77-year-old woman with dysphagia and achalasia following a stroke presented with recurrent lung infiltrates, including a persisting right middle lobe lesion. Hounsfield units (HU) measurement was -157, consistent with lipoid pneumonia. Comparative subcutaneous fat and aorta (blood/tissue) HU are shown

From the collection of Dr Augustine Lee, used with permission of Mayo Foundation for Medical Education and Research, all rights reserved

Disclaimer

BMJ Best Practice is intended for licensed medical professionals. BMJ Publishing Group Ltd (BMJ) does not advocate or endorse the use of any drug or therapy contained within this publication nor does it diagnose patients. As a medical professional you retain full responsibility for the care and treatment of your patients and you should use your own clinical judgement and expertise when using this product.

This content is not intended to cover all possible diagnosis methods, treatments, follow up, drugs and any contraindications or side effects. In addition, since such standards and practices in medicine change as new data become available, you should consult a variety of sources. We strongly recommend that you independently verify specified diagnosis, treatments and follow-up and ensure it is appropriate for your patient within your region. In addition, with respect to prescription medication, you are advised to check the product information sheet accompanying each drug to verify conditions of use and identify any changes in dosage schedule or contraindications, particularly if the drug to be administered is new, infrequently used, or has a narrow therapeutic range. You must always check that drugs referenced are licensed for the specified use and at the specified doses in your region.

Information included in BMJ Best Practice is provided on an “as is” basis without any representations, conditions or warranties that it is accurate and up to date. BMJ and its licensors and licensees assume no responsibility for any aspect of treatment administered to any patients with the aid of this information. To the fullest extent permitted by law, BMJ and its licensors and licensees shall not incur any liability, including without limitation, liability for damages, arising from the content. All conditions, warranties and other terms which might otherwise be implied by the law including, without limitation, the warranties of satisfactory quality, fitness for a particular purpose, use of reasonable care and skill and non-infringement of proprietary rights are excluded.

Where BMJ Best Practice has been translated into a language other than English, BMJ does not warrant the accuracy and reliability of the translations or the content provided by third parties (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages). BMJ is not responsible for any errors and omissions arising from translation and adaptation or otherwise. Where BMJ Best Practice lists drug names, it does so by recommended International Nonproprietary Names (rINNs) only. It is possible that certain drug formularies might refer to the same drugs using different names.

Please note that recommended formulations and doses may differ between drug databases drug names and brands, drug formularies, or locations. A local drug formulary should always be consulted for full prescribing information.

Treatment recommendations in BMJ Best Practice are specific to patient groups. Care is advised when selecting the integrated drug formulary as some treatment recommendations are for adults only, and external links to a paediatric formulary do not necessarily advocate use in children (and vice-versa). Always check that you have selected the correct drug formulary for your patient.

Where your version of BMJ Best Practice does not integrate with a local drug formulary, you should consult a local pharmaceutical database for comprehensive drug information including contraindications, drug interactions, and alternative dosing before prescribing.

Interpretation of numbers

Regardless of the language in which the content is displayed, numerals are displayed according to the original English-language numerical separator standard. For example 4 digit numbers shall not include a comma nor a decimal point; numbers of 5 or more digits shall include commas; and numbers stated to be less than 1 shall be depicted using decimal points. See Figure 1 below for an explanatory table.

BMJ accepts no responsibility for misinterpretation of numbers which comply with this stated numerical separator standard.

This approach is in line with the guidance of the [International Bureau of Weights and Measures Service](#).

Figure 1 – BMJ Best Practice Numeral Style

5-digit numerals: 10,000

4-digit numerals: 1000

numerals < 1: 0.25

Our full website and application terms and conditions can be found here: [Website Terms and Conditions](#).

Contact us

+ 44 (0) 207 111 1105

support@bmj.com

BMJ

BMA House

Tavistock Square

London

WC1H 9JR

UK

BMJ Best Practice

Contributors:

// Authors:

Augustine Lee, MD

Professor of Medicine

Division of Pulmonary and Critical Care Medicine, Mayo Clinic Florida, Jacksonville, FL

DISCLOSURES: AL declares that he has no competing interests.

Spencer Deleveaux, MBBS

Fellow

Division of Pulmonary and Critical Care Medicine, Mayo Clinic Florida, Jacksonville, FL

DISCLOSURES: SD declares that he has no competing interests.

// Acknowledgements:

Dr Augustine Lee and Dr Spencer Deleveaux would like to gratefully acknowledge Dr Madison Macht, Dr Kamran Mahmood, Dr Scott Shofer, Dr Septimiu Murgu, and Dr Henri Colt, previous contributors to this topic.

DISCLOSURES: MM, KM, SS, SM, and HC declare they have no competing interests.

// Peer Reviewers:

Andrew Parfitt, MBBS, FFAEM

Clinical Director

Acute Medicine, Associate Medical Director, Consultant Emergency Medicine, Guy's and St Thomas' NHS Foundation Trust, Clinical Lead and Consultant, Accident and Emergency Medicine, St Thomas' Hospital, London, UK

DISCLOSURES: AP declares that he has no competing interests.

Momen M. Wahidi, MD, MBA

Director

Interventional Pulmonology, Division of Pulmonary, Allergy, and Critical Care Medicine, Assistant Professor of Medicine, Duke University Medical Center, Durham, NC

DISCLOSURES: MMW declares that he has no competing interests.